

KU ScholarWorks

Trends in Alcohol Use Disorder Treatment Utilization and Location from 2008 to 2017

| | |
|---------------|--|
| Item Type | Article |
| Authors | Larsen, Aidan R.;Cummings, Janet R.;von Esenwein, Silke A.;Druss, Benjamin G. |
| Citation | Larsen AR, Cummings JR, von Esenwein SA, Druss BG. Trends in Alcohol Use Disorder Treatment Utilization and Setting From 2008 to 2017. Psychiatr Serv. 2022 Sep 1;73(9):991-998. doi: 10.1176/appi.ps.202000323. PMID: 35193376. |
| DOI | 10.1176/appi.ps.202000323 |
| Publisher | Psychiatry Online |
| Rights | Copyright 2022 Psychiatry Online |
| Download date | 2024-08-18 05:04:42 |
| Link to Item | https://hdl.handle.net/1808/33590 |

Trends in Alcohol Use Disorder Treatment Utilization and Location from 2008 to 2017

Aidan R. Larsen, MSPH^a, Janet R. Cummings, Ph.D^b., Silke A. von Esenwein, Ph.D^c,

Benjamin G. Druss, M.D, M.P.H^b

^a Mathematica Policy Research, Washington, D.C.

^b Department of Health Policy and Management, Rollins School of Public Health, Emory University, Atlanta

^c University of Kansas, Center for Public Partnerships and Research, Lawrence

Disclosures and acknowledgments: All authors have no financial interests or other conflicts of interest to disclose.

Word Count: 2,984

Highlights:

- Data from the 2008-2017 National Survey of Drug Use and Health were used to study trends in overall treatment utilization and changes in treatment location among those with alcohol use disorder.
- Treatment for alcohol use disorder declined by 1.1 percentage points from 6.9% in 2008-2010 to 5.8% between 2014-2017.
- After controlling for confounders, compared to the baseline of 2008-2010, the probability of receiving no treatment increased by 1.4 percentage points in 2011-2013 and 1.5 percentage points in 2014-2017.
- In adjusted models, compared to the baseline of 2008-2010, the likelihood of receiving treatment in any medical location declined by 1.0 percentage points and the probability of receiving only self-help treatment declined by 0.5 percentage points in 2014-2017.

Trends in Alcohol Use Disorder Treatment Utilization and Location from 2008 to 2017

Disclosures and acknowledgments: All authors have no financial interests or other conflicts of interest to disclose.

Word Count: 2,984

Highlights:

- Data from the 2008-2017 National Survey of Drug Use and Health were used to study trends in overall treatment utilization and changes in treatment location among those with alcohol use disorder.
- Treatment for alcohol use disorder declined by 1.1 percentage points from 6.9% in 2008-2010 to 5.8% between 2014-2017.
- After controlling for confounders, compared to the baseline of 2008-2010, the probability of receiving no treatment increased by 1.4 percentage points in 2011-2013 and 1.5 percentage points in 2014-2017.
- In adjusted models, compared to the baseline of 2008-2010, the likelihood of receiving treatment in any medical location declined by 1.0 percentage points and the probability of receiving only self-help treatment declined by 0.5 percentage points in 2014-2017.

Abstract

Objective: Little is known about recent trends in treatment for alcohol use disorder. This study used national data to examine treatment trends among individuals with alcohol use disorder.

Methods: A sample of non-elderly adults (18-64) with alcohol use disorder were identified from the National Survey of Drug Use and Health. Multinomial logistic regression analysis was conducted to examine trends in treatment for alcohol use disorder in 2008-2010, 2011-2013, and 2014-2017 in: (1) any medical setting (including hospitals, rehab centers, mental health centers, emergency room, and private doctors' offices); (2) self-help groups only (no medical setting); and (3) no setting (i.e., no treatment). Additional analyses investigated trends in mental health treatment. Regression models adjusted for predisposing, enabling, and need-related characteristics.

Results: Among those with an alcohol use disorder, the percentage that received any treatment was significantly lower in 2011-2013 (5.6%) compared to 2008-2010 (6.9%, $p < 0.05$). In adjusted analyses, the probability of receiving no treatment increased by 1.5 percentage points in 2014-2017 (95% CI = 0.5, 2.5), compared to the baseline of 2008-2010. Significant declines were observed in the receipt of any treatment in a medical setting (M.E. = -1.0%, 95% CI = -2.0, -0.0) and self-help treatment only (M.E. = -0.5%, 95% CI = -0.8, -0.1) in 2014-2017 compared to baseline. The probability of receiving any mental health treatment did not change during the study period.

Conclusions: Treatment among those with an alcohol use disorder declined from 2008-2017. Future studies should examine the mechanisms that may be responsible for the decline in treatment.

As of 2018, alcohol use disorder was the most common substance use disorder in the United States, with 5.4% of Americans over age twelve affected (1). Alcohol use disorder is associated with various medical problems, including cancer, diabetes, and cardiovascular disease, and it was associated with 95,158 deaths a year from 2011-2015 in the United States (2-5). Although efficacious treatments for alcohol use disorder exist, only about 10% of those with alcohol use disorder receive treatment in a given year and many individuals with alcohol use disorder seek mental health (MH) treatment instead of alcohol treatment (6, 7).

In the past decade, two significant laws were implemented with the potential to address the treatment gap for alcohol use and other behavioral health disorders -- the Mental Health Parity and Addiction Equity Act (MHPAEA) and the Patient Protection and Affordable Care Act (ACA) -- in 2008 and 2010, respectively. These laws increased the number of insured, increased coverage for behavioral health treatment among the insured, and more tightly integrated behavioral healthcare and medical healthcare (8-13). The MHPAEA established parity for substance use disorder benefits, which required that both treatment limits (e.g., caps on outpatient visits) and financing (e.g., co-pays) were no more restrictive for substance use treatment than for medical benefits (8). The ACA extended parity to individual and Medicaid plans and required covering substance use disorder benefits as a part of the essential health benefits (EHB) (9). The ACA also increased the number of insured individuals through the individual marketplace and Medicaid expansion (10, 12). Finally, payment reform and other ACA incentives were expected to lead to greater substance use disorder treatment integration with primary care, facilitating access to treatment (11, 13).

Several studies have examined treatment trends for substance use disorders since the MHPAEA and ACA were passed (14-18). These studies have typically found no change in the

treatment rate for substance use disorders. However, these studies aggregate treatment for all substance use disorders, potentially masking differential trends for illicit drug use disorders and alcohol use disorder. To date, only one known study has used national data (from 2010 to 2015) to investigate recent trends in alcohol use disorder treatment specifically (19). This study reported no change in alcohol use disorder treatment over that period. Because this study only included data through 2015 (the first full year after ACA implementation), more recent data are needed to understand how trends in treatment for alcohol use disorder have changed since its implementation. In addition, prior research examined trends in alcohol use disorder treatment across all settings in aggregate (including medical settings and self-help groups) instead of by treatment location.

To address this gap in the literature, we used national data from the 2008-2017 National Survey of Drug Use and Health to examine trends in treatment for alcohol use disorder. We hypothesized that alcohol use disorder treatment increased over that period due, in part, to the increase in insurance coverage because of the ACA. We examined three measures of alcohol use disorder treatment (no treatment, any treatment in a medical setting, self-help treatment only) and MH treatment for those with alcohol use disorder.

Methods

Data and Sample

We used pooled data from the 2008-2017 National Survey of Drug Use and Health (NSDUH). NSDUH is an annual, nationally-representative, cross-sectional survey of the US civilian, non-institutionalized population over 12 (20). NSDUH uses DSM-IV criteria to determine whether respondents have an alcohol use disorder (i.e., alcohol abuse or dependence). We identified 38,335 individuals with alcohol use disorder between 18-64 in the pooled sample,

of whom 36,707 had complete information on all of the model covariates for the analytic sample used to examine the three-alcohol use disorder treatment outcomes. When estimating a separate MH treatment model among those with alcohol use disorder, an additional 79 observations were missing on the measure of MH treatment and dropped from the analytic sample. Similar to prior research examining behavioral health services changes over time using NSDUH data, we examined trends in alcohol use disorder treatment across three periods: 2008-2010, 2011-2013, and 2014-2017 (14, 16, 18, 19).

Dependent Variables

The receipt of any alcohol use disorder treatment was measured using a categorical variable (received no treatment; received any medical treatment; received self-help treatment only). We derived this measure from a question in which respondents were asked whether they received treatment for alcohol or drugs in the last twelve months and whether that treatment was for alcohol, drugs, or both. Those who did not report treatment for alcohol or both alcohol and drugs were coded as not receiving alcohol treatment.

Respondents who received treatment were categorized based on the location in which they reported receiving treatment – either in a medical setting (hospital as an inpatient, rehab center, MH center, emergency room, private doctor’s office), or whether they received self-help only treatment such as Alcoholics Anonymous; we excluded from analyses those who only received treatment in jail or for whom treatment location could not be ascertained (n=654). Consistent with prior studies using NSDUH (21-23) we aggregated treatment locations into these categories to separate treatment locations where health insurance coverage is likely to facilitate financial access to services (medical settings) and where insurance is unlikely to make a difference (self-help treatment only).

The receipt of MH treatment was measured using a dichotomous variable (yes/no). Respondents who reported receiving inpatient treatment, outpatient treatment, or prescription medication for MH treatment were coded as receiving MH treatment. We include MH treatment as a separate dependent variable because previous research has found that individuals with alcohol use disorder are more likely to seek MH treatment than substance use disorder treatment and that outpatient MH treatment increased overall in the years after the ACA was passed (7, 24). However, it is unclear to what extent utilization of MH treatment changed for those with alcohol use disorder.

Independent Variables

Sociodemographic characteristics. Regression models controlled for predisposing and enabling characteristics (25). Predisposing characteristics included age category (18 to 25, 26 to 34, 35 to 49, and 50 to 64), race/ethnicity (Non-Hispanic white, Non-Hispanic black, Hispanic, and Non-Hispanic other), gender (male versus female), educational attainment (less than high school, high school graduate, some college/associate's degree, and college graduate and above), marital status (married, widowed or divorced, and never married), and employment status (full time employed, part-time employed, unemployed, and other). Due to small sample sizes, we combined four race/ethnicity categories into Non-Hispanic other for analysis: non-Hispanic Native Am/AK Native, non-Hispanic Native HI/Other Pac Islander, non-Hispanic Asian, and non-Hispanic more than one race. Enabling characteristics included income (less than \$20,000, \$20,000 to \$49,999, \$50,000 to \$74,999, \$75,000 or more), and insurance status (any private insurance, Medicaid [no private insurance], other insurance [no Medicaid or private, and other insurance], and uninsured). Income was imputed for 8.6% of the sample (3,167).

Health status. Regression models also adjusted for differences in need-related characteristics (25). Health status measures included a categorical measure of self-reported health status (excellent, very good, good, fair, and poor), dichotomous indicators for comorbid illicit substance use disorder, comorbid MH disorder, and alcohol use disorder severity. Comorbid substance use disorder was created from heroin abuse or dependence, cocaine abuse or dependence, and marijuana abuse or dependence; these were the measures of illicit substance abuse or dependence that were consistently measured during the study period (26). The measure of comorbid MH status was created using information from the NSDUH prediction model that determines the predicted probability of any mental illness based on responses to several questions; individuals with a predicted probability greater or equal to a specified cutoff value were considered to have a comorbid MH disorder (26). Alcohol use disorder severity was assessed with an indicator for whether the individual met DSM-IV criteria for alcohol dependence (versus alcohol abuse). In addition, the models included indicators for the following self-reported health conditions: diabetes, liver cirrhosis, hepatitis, asthma, HIV or AIDS, and high blood pressure. Each condition was assessed with an indicator for whether the individual reported ever being told by a doctor or other health professional they had each respective condition.

Analysis

Analysis was conducted in STATA using the SVY command to account for the complex survey design elements of NSDUH. We first conducted bivariate analyses using Adjusted Wald tests to compare treatment rates in 2011-2013 and 2014-2017 to the baseline period of 2008-2010. Next, we adjusted for predisposing, enabling, and need-related factors that may have changed over time using multinomial and binomial logistic regression models.

We estimated a multinomial logistic regression for the dependent variable of alcohol treatment: received no treatment, received any treatment in a medical location, and received self-help group treatment only (i.e., no medical treatment). We estimated a separate logistic regression for the dependent variable of any MH treatment. Marginal effects were reported for all models. Marginal effects can be interpreted as the percentage point change in the predicted probability that individuals fall into one outcome category compared to the other outcome categories combined, holding all other covariates constant. The alpha level for statistical significance was set to 0.05.

This study was exempt from IRB review in a letter of determination from Emory University's IRB because the data came from publicly available sources without Private Health Information identifiers.

Results

The prevalence of alcohol use disorder among adults declined over the period from 2008-2017. Bivariate analysis indicated a decline in alcohol use disorder among all NSDUH respondents from 8.9% in 2008-2010 to 8.1% in 2011-2013 ($p < 0.01$) to 7.3% in 2014-2017 ($p < 0.01$) (Table 1). However, among those with alcohol use disorder, a higher proportion had alcohol dependence, the more severe form of alcohol use disorder, in 2014-2017 (52.3%) than in 2008-2010 (46.8%, $p < 0.01$) (Table 2).

Among those with an alcohol use disorder, bivariate analysis indicated that there was an overall decline in any treatment for alcohol from 6.9% in 2008-2010 to 5.6% ($p < 0.05$) in 2011-2013 (Table 2). There was a significant decline in treatment in any medical setting from 5.6% in 2008-2010 to 4.4% in 2011-2013 ($p < 0.05$). Both any treatment and treatment in any medical settings had nonsignificant declines from 2008-2010 to 2014-2017. Self-help treatment only

declined from 1.2% in 2008-2010 to 0.8% ($p<0.05$) in 2014-2017. For MH treatment, there was a nonsignificant increase from 22.6% in 2008-2010 to 23.9% in 2014-2017.

Next, we examined differences in characteristics of those with an alcohol use disorder across these periods (Table 2). Notably, there were significant health insurance status changes for those with alcohol use disorder. A higher percentage of those with alcohol use disorder reported having Medicaid coverage (13.6%) and private insurance (64.5%) in 2014-2017 compared to 2008-2010 (8.0%, $p<0.01$; 61.4%, $p<0.01$). The percentage of those with alcohol use disorder that were uninsured declined from 25.1% in 2008-2010 to 16.0% in 2014-2017 ($p<0.01$). In addition to increases in insurance coverage, there were significant changes in other enabling characteristics. A higher percentage of those with alcohol use disorder had incomes above \$75,000 per year (35.9%) and obtained college degrees or higher (30.6%) in 2014-2017 compared to 2008-2010 (30.0%, $p<0.01$; 24.5%, $p<0.01$).

We estimated a multinomial logistic regression to assess whether there were significant changes in the probability of treatment after controlling for changes in predisposing, enabling, and need characteristics during the study period (Table 3). Compared to 2008-2010, the probability of not receiving treatment (versus receiving treatment in any setting) increased by 1.4 percentage points (95% CI= 0.4, 2.4) in 2011-2013 and 1.5 percentage points (95% CI= 0.5, 2.5) in 2014-2017, after adjusting for confounders. There were also significant declines in treatment in medical settings and self-help treatment only. Compared to baseline, the probability of receiving treatment in any medical location declined by 1.3 percentage points in 2011-2013 (95% CI= -2.2, -0.4) and 1.0 percentage points in 2014-2017 (95% CI= -2.0, -0.0). Moreover, the probability of self-help treatment only declined 0.5 percentage points in 2014-2017 (95% CI = -0.8, -0.1).

Insurance status was also associated with trends in alcohol treatment. Compared to being uninsured, those with Medicaid insurance had a lower probability of not receiving any treatment (marginal effect = -2.2 percentage points; 95% CI = -3.9, -0.4). Moreover, Medicaid increased the probability of receiving treatment in a medical location by 2.3 percentage points (95% CI = 0.5, 4.0).

Results from the logistic regression model for MH treatment (Table 4) as the dependent variable were similar to those from the bivariate analysis. After controlling for predisposing, enabling, and need factors, there were no significant differences in respondents' likelihood of receiving any MH treatment across the study period.

Discussion

Treatment among those with alcohol use disorder declined from 6.9% in 2008-2010 to 5.6% in 2011-2013. When predisposing, enabling, and need-related factors were controlled, significant decreases in alcohol use disorder treatment were observed in medical settings and self-help settings. Furthermore, during the study period, the probability of not receiving any treatment increased by 1.4 percentage points in 2011-2013 and 1.5 percentage points in 2014-2017 compared to the baseline 2008-2010.

Our results differ from previous studies of the trends in substance use disorder treatment. Earlier research found no increase in substance use or specialty substance use treatment among those with a substance use disorder from 2005-2014 (14) or 2011-2014 (16). The decline noted in this study may be due to the additional years of data available and/or to differences in the population with an alcohol use disorder compared to the broader population with any substance use disorder.

Although there was a decline in the proportion of those with alcohol use disorder who were uninsured during the study period, the probability of receiving no treatment increased. There are several potential explanations for the observed declines in treatment. First, even with expanded insurance coverage, the out-of-pocket cost may be high for individuals seeking treatment. Although MHPAEA intended to reduce the treatment limitations and cost-sharing patients face when seeking behavioral health care, recent research has questioned whether MHPAEA was effective. In 2017, substance use disorder care was more likely to be out-of-network than medical/surgical care for inpatient visits (10.1x), outpatient visits (8.5x), and office visits (9.5x). These disparities increased for all service types from 2013 to 2017 (27). Out-of-network services have higher cost-sharing than in-network services and are thus more expensive for individuals. Although many individuals with alcohol use disorder gained insurance, the out-of-pocket expenses for receiving alcohol treatment may have been high if seeking out-of-network services (28).

Another possible explanation is that the supply of alcohol treatment facilities is limited, preventing individuals with alcohol use disorder from accessing alcohol treatment. However, data from the National Survey of Substance Abuse Treatment Services (N-SSATS), an annual survey of substance use treatment facilities in the United States, shows a modest increase in the number of facilities offering various alcohol use disorder treatment types during the study period. (29, 30). Even so, the number of individuals who received treatment for alcohol alone or alcohol and drugs in these facilities has decreased every year since 2011, declining from 756,890 in 2011 to 712,480 in 2017 (29). Future research should examine whether an increased emphasis on or increased demand for other types of substance use disorder treatment, such as opioid use disorder, has reduced these facilities' capacity to treat those with alcohol use disorder.

In addition to declines in overall treatment, self-help treatment declined by 0.5 percentage points in 2014-2017 compared to the baseline of 2008-2010. This finding is consistent with data that has reported a steady decline in Alcoholics Anonymous members since 2001 (31). Future efforts to increase treatment rates may need to address attitudinal barriers such as stigma and awareness about the effectiveness of therapies for alcohol use disorder and the financial barriers that were the focus of the ACA and MHPAEA.

The current findings suggest that expanded insurance was not enough to increase rates of alcohol use disorder treatment. Potential interventions to increase demand for treatment among those with alcohol use disorder may be one way to increase treatment. Mass media public awareness campaigns can be a step in this direction, although they may need to be coupled with other demand side and targeted interventions such as advertising restrictions and the targeting of high-risk groups, as the overall evidence base for their efficacy at increasing treatment uptake is relatively weak (32).

It is also notable that there were no significant changes in the likelihood of receiving MH treatment during the study period after controlling for confounders. While the percentage of those with alcohol use disorder with a comorbid MH condition significantly increased during the study period, there was no commensurate increase in MH treatment. These findings suggest that individuals with alcohol use disorder are not likely to substitute for a decline in alcohol treatment with increases in MH treatment.

This study has several limitations. These analyses are descriptive; other unmeasured constructs may confound changes over time, and causality cannot be established in any reported associations. Thus, this study cannot provide reasons for the unexpected decline in alcohol use disorder treatment. Several potential explanations need to be explored in future studies to better

understand these trends, including changes in insurance benefits for alcohol use disorder treatment, changes in the geographic availability and capacity of substance use treatment facilities that offer alcohol use disorder services, changes in screening practices for alcohol use disorder, and changes in attitudes about alcohol use disorder treatment. In addition, the study only contains four years of post-ACA data. Another limitation is that survey design changes in 2015 restricted the measurement of comorbid substance use disorders in these analyses to those consistently measured during the study period (26). Lastly, NSDUH excludes the institutionalized and non-civilian US population. The estimates from this study cannot be generalized to those populations.

Conclusions

This study findings show unexpected declines in the treatment rates for alcohol use disorder in any setting, including medical and self-help settings, in both unadjusted and adjusted analyses. Given the gains in insurance coverage during this period, further research is needed to understand and address the factors responsible for the decline in treatment.

References

1. Key Substance Use and Mental Health Indicators in the United States: Results from the 2018 National Survey on Drug Use and Health. Rockville, MD, Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, 2019
2. Rehm J: The risks associated with alcohol use and alcoholism. *Alcohol Res Health* 2011; 34:135-43
3. Rehm J, Dawson D, Frick U, et al: Burden of disease associated with alcohol use disorders in the United States. *Alcohol Clin Exp Res* 2014; 38:1068-77
4. Esser MB, Sherk A, Liu Y, et al: Deaths and Years of Potential Life Lost From Excessive Alcohol Use - United States, 2011-2015. *MMWR Morb Mortal Wkly Rep* 2020; 69:1428-33
5. Stahre M, Roeber J, Kanny D, et al: Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis* 2014; 11:E109
6. Schmidt LA: Recent Developments in Alcohol Services Research on Access to Care. *Alcohol research : current reviews* 2016; 38:27-33
7. Edlund MJ, Booth BM, Han X: Who Seeks Care Where? Utilization of Mental Health and Substance Use Disorder Treatment in Two National Samples of Individuals with Alcohol Use Disorders. *Journal of Studies on Alcohol and Drugs* 2012; 73:635-46
8. Busch SH: Implications of the Mental Health Parity and Addiction Equity Act. *Am J Psychiatry* 2012; 169:1-3
9. Barry CL, Huskamp HA: Moving beyond parity--mental health and addiction care under the ACA. *N Engl J Med* 2011; 365:973-5
10. Mojtabai R, Feder KA, Kealhofer M, et al: State variations in Medicaid enrollment and utilization of substance use services: Results from a National Longitudinal Study. *J Subst Abuse Treat* 2018; 89:75-86
11. McLellan AT, Woodworth AM: The affordable care act and treatment for "Substance Use Disorders:" Implications of ending segregated behavioral healthcare. *Journal of Substance Abuse Treatment* 2014; 46:541-5
12. Tolbert J, Orgera K, Damico A: Key Facts about the Uninsured Population: Kaiser Family Foundation, 2020. <https://www.kff.org/uninsured/issue-brief/key-facts-about-the-uninsured-population/>
13. Buck JA: The Looming Expansion And Transformation Of Public Substance Abuse Treatment Under The Affordable Care Act. *Health Aff (Millwood)* 2011; 30:1402-10
14. Creedon TB, Cook BL: Access To Mental Health Care Increased But Not For Substance Use, While Disparities Remain. *Health Aff (Millwood)* 2016; 35:1017-21
15. Saloner B, Le Cook B: An ACA Provision Increased Treatment For Young Adults With Possible Mental Illnesses Relative To Comparison Group. *Health Aff (Millwood)* 2014; 33:1425-34
16. Saloner B, Bandara S, Bachhuber M, et al: Insurance Coverage and Treatment Use Under the Affordable Care Act Among Adults With Mental and Substance Use Disorders. *Psychiatr Serv* 2017; 68:542-8
17. Olfson M, Wall M, Barry CL, et al: Impact of Medicaid expansion on coverage and treatment of low-income adults with substance use disorders. *Health Affairs* 2018; 37:1208-15
18. Olfson M, Wall M, Barry CL, et al: Effects of the Affordable Care Act on Private Insurance Coverage and Treatment of Behavioral Health Conditions in Young Adults. *American journal of public health* 2018; 108:1352-4

19. Feder KA, Mojtabai R, Krawczyk N, et al: Trends in insurance coverage and treatment among persons with opioid use disorders following the Affordable Care Act. *Drug Alcohol Depend* 2017; 179:271-4
20. 2017 National Survey on Drug Use and Health Final Analytic File Codebook. Rockville, MD, Substance Abuse and Mental Health Services Administration, 2018
21. Bandara SN, Samples H, Crum RM, et al: Is screening and intervention associated with treatment receipt among individuals with alcohol use disorder? Evidence from a national survey. *Journal of Substance Abuse Treatment* 2018; 92:85-90
22. Johnson K, Rigg KK, Hopkins Eyles C: Receiving addiction treatment in the US: Do patient demographics, drug of choice, or substance use disorder severity matter? *International Journal of Drug Policy* 2020; 75:102583
23. Romo E, Ulbricht CM, Clark RE, et al: Correlates of specialty substance use treatment among adults with opioid use disorders. *Addictive Behaviors* 2018; 86:96-103
24. Olfson M, Wang S, Wall M, et al: Trends in Serious Psychological Distress and Outpatient Mental Health Care of US Adults. *JAMA Psychiatry* 2019; 76:152-61
25. Andersen RM, Davidson PL: Improving Access to Care in America: Individual and Contextual Indicators; in *Changing the US health care system: Key issues in health services policy and management*, 3rd ed. San Francisco, CA, US: Jossey-Bass, 2007
26. 2015 National Survey on Drug Use and Health Public Use Data File Codebook. Rockville, MD, Substance Abuse and Mental Health Services Administration, 2018
27. Davenport S, Gray TJ, Melek SP: Addiction and mental health vs. physical health: Widening disparities in network use and provider reimbursement. *Millman Research Report* 2019;
28. Thomas CP, Hodgkin D, Levit K, et al: Growth in spending on substance use disorder treatment services for the privately insured population. *Drug Alcohol Depend* 2016; 160:143-50
29. National Survey of Substance Abuse Treatment Services (N-SSATS): 2017. Data on Substance Abuse Treatment Facilities. Rockville, MD, Substance Abuse and Mental Health Services Administration, 2018.
30. National Survey of Substance Abuse Treatment Services (N-SSATS): 2011. Data on Substance Abuse Treatment Facilities. Rockville, MD, Substance Abuse and Mental Health Services Administration, 2012.
31. C J: AA Membership: Growth or Decline? *AA Agnostica*, 2015.
<https://aaagnostica.org/2015/09/06/aa-membership-growth-or-decline/>
32. Young B, Lewis S, Katikireddi SV, et al: Effectiveness of Mass Media Campaigns to Reduce Alcohol Consumption and Harm: A Systematic Review. *Alcohol and Alcoholism* 2018; 53:302-16

TABLE 1. Prevalence of adults with alcohol use disorder from 2008-2017 NSDUH^a

| Period | Total N | N | Weighted % |
|-----------|---------|--------|------------------|
| 2008-2010 | 107,893 | 13,432 | 8.9 |
| 2011-2013 | 107,713 | 11,733 | 8.1 ^b |
| 2014-2017 | 155,868 | 13,170 | 7.3 ^b |

^a NSDUH, National Survey of Drug Use and Health. P values from adjusted Wald test comparing 2011-2013 and 2014-2017 values to the baseline of 2008-2010

^b P-value less than .001

TABLE 2. Characteristics of individuals with alcohol use disorder from the 2008-2017 NSDUH^a

| Variables | 2008-2010 | | 2011-2013 | | | 2014-2017 | | |
|---|---|------------|---|------------|----------------|---|------------|----------------|
| | (N=12,810 ^b /12,788 ^c) | | (N=11,211 ^b /11,186 ^c) | | | (N=12,686 ^b /12,654 ^c) | | |
| | N | Weighted % | N | Weighted % | P ^a | N | Weighted % | P ^a |
| Alcohol Treatment | | | | | | | | |
| Any | 780 | 6.9 | 573 | 5.6 | .033 | 738 | 5.8 | .054 |
| Medical Location | 644 | 5.6 | 454 | 4.4 | .030 | 626 | 5.0 | .252 |
| Self-Help Only | 136 | 1.2 | 119 | 1.1 | .290 | 112 | 0.8 | .017 |
| Mental Health Treatment | 2,504 | 22.6 | 2,373 | 23.3 | .365 | 2,922 | 23.9 | .130 |
| Age | | | | | | | | |
| 18-25 | 8,731 | 32.7 | 7,415 | 30.5 | .019 | 5,756 | 26.6 | <.001 |
| 26-34 | 1,878 | 25.3 | 1,650 | 25.4 | .920 | 3,042 | 25.2 | .918 |
| 35-49 | 1,722 | 26.9 | 1,598 | 27.6 | .569 | 2,933 | 27.4 | .674 |
| 50-64 | 479 | 15.1 | 548 | 16.4 | .237 | 955 | 20.8 | <.001 |
| Male | 7,749 | 65.9 | 6,658 | 64.0 | .043 | 7,456 | 63.2 | .009 |
| Race/Ethnicity | | | | | | | | |
| NH-White | 8,602 | 69.6 | 7,274 | 67.9 | .115 | 8,002 | 65.6 | <.001 |
| NH-Black | 1,218 | 11.0 | 1,125 | 10.2 | .377 | 1,318 | 11.1 | .844 |
| Hispanic | 1,910 | 15.2 | 1,792 | 16.7 | .049 | 2,074 | 16.7 | .043 |
| NH-Other | 1,080 | 4.2 | 1,020 | 5.2 | .014 | 1,292 | 6.6 | <.001 |
| Marital Status | | | | | | | | |
| Married | 2,181 | 32.3 | 1,879 | 32.3 | .969 | 3,053 | 33.2 | .437 |
| Widowed or Divorced | 1,121 | 16.1 | 1,044 | 16.3 | .829 | 1,551 | 16.4 | .737 |
| Never Married | 9,508 | 51.6 | 8,288 | 51.5 | .901 | 8,082 | 50.4 | .285 |
| Income | | | | | | | | |
| Less than \$20,000 | 3,810 | 22.2 | 3,477 | 23.4 | .234 | 3,086 | 20.5 | .047 |
| \$20,000 - \$49,999 | 4,189 | 31.0 | 3,691 | 31.5 | .619 | 3,973 | 28.3 | .002 |
| \$50,000 - \$74,999 | 1,908 | 16.8 | 1,505 | 14.8 | .031 | 1,893 | 15.2 | .044 |
| \$75,000 or more | 2,903 | 30.0 | 2,538 | 30.3 | .863 | 3,734 | 35.9 | <.001 |
| Self-Reported Health Status | | | | | | | | |
| Excellent | 2,871 | 20.3 | 2,404 | 20.2 | .877 | 2,444 | 18.5 | .012 |
| Very Good | 5,249 | 39.3 | 4,626 | 38.5 | .553 | 5,044 | 38.5 | .447 |
| Good | 3,523 | 28.9 | 3,084 | 28.2 | .475 | 3,749 | 29.8 | .293 |
| Fair | 1,028 | 9.6 | 933 | 10.6 | .091 | 1,273 | 11.0 | .041 |
| Poor | 139 | 1.9 | 164 | 2.5 | .147 | 176 | 2.1 | .506 |
| Education | | | | | | | | |
| Less than high school | 2,191 | 15.8 | 1,690 | 14.1 | .050 | 1,537 | 11.7 | <.001 |
| High School Graduate | 4,021 | 29.8 | 3,366 | 27.2 | .025 | 3,280 | 24.3 | <.001 |
| Some College/Associate's Degree | 4,202 | 29.9 | 3,868 | 30.8 | .362 | 4,569 | 33.4 | <.001 |
| College Graduate and Above | 2,396 | 24.5 | 2,287 | 27.9 | .001 | 3,300 | 30.6 | <.001 |
| Employment Status | | | | | | | | |
| Full-Time | 6,425 | 58.7 | 5,633 | 57.9 | .514 | 7,442 | 61.3 | .021 |
| Part-Time | 2,753 | 16.9 | 2,410 | 15.9 | .244 | 2,199 | 15.1 | .013 |
| Unemployed | 1,514 | 9.9 | 1,286 | 9.3 | .337 | 1,089 | 7.6 | <.001 |
| Other | 2,118 | 14.5 | 1,882 | 17.0 | .005 | 1,956 | 16.0 | .023 |
| Comorbid Substance Use Disorder | 1,997 | 11.4 | 1,668 | 11.1 | .664 | 1,685 | 11.3 | .868 |
| Comorbid Mental Illness | 4,458 | 36.0 | 4,114 | 38.8 | .009 | 5,282 | 39.3 | .001 |
| Alcohol Dependence | 5,635 | 46.8 | 5,123 | 47.1 | .790 | 6,396 | 52.3 | <.001 |
| Number of Alcohol Use Disorder Criteria | | | | | | | | |
| 1-2 | 4,224 | 33.6 | 3,633 | 33.3 | .840 | 3,908 | 30.7 | .003 |
| 3-4 | 5,362 | 40.6 | 4,816 | 41.3 | .550 | 5,420 | 42.0 | .101 |
| 5+ | 3,224 | 25.8 | 2,762 | 25.4 | .649 | 3,358 | 27.3 | .061 |
| Insurance Status | | | | | | | | |
| Private | 7,445 | 61.4 | 6,681 | 60.6 | .558 | 8,009 | 64.5 | .004 |
| Medicaid | 1,189 | 8.0 | 1,127 | 8.3 | .585 | 1,837 | 13.6 | <.001 |
| Other Insured | 774 | 5.9 | 769 | 6.0 | .746 | 871 | 6.3 | .424 |
| Uninsured | 3,402 | 24.8 | 2,634 | 25.0 | .824 | 1,969 | 15.6 | <.001 |
| Health Indicators | | | | | | | | |
| Diabetes | 224 | 3.1 | 223 | 3.3 | .690 | 405 | 4.4 | .007 |
| Cirrhosis | 23 | 0.3 | 29 | 0.4 | .451 | 45 | 0.5 | .217 |

| | | | | | | | | |
|---------------------|-------|------|-------|------|------|-------|------|------|
| Hepatitis B or C | 96 | 1.3 | 94 | 1.7 | .121 | 157 | 1.9 | .028 |
| Asthma | 1,852 | 13.0 | 1,607 | 13.4 | .535 | 1,607 | 11.7 | .057 |
| HIV or AIDS | 32 | 0.4 | 22 | 0.2 | .340 | 40 | 0.4 | .801 |
| High Blood Pressure | 1,113 | 15.2 | 1,080 | 15.8 | .509 | 1,234 | 13.7 | .048 |

^aNSDUH, National Survey of Drug Use and Health. P values from adjusted Wald test comparing 2011-2013 and 2014-2017 values to the baseline of 2008-2010

^bSample for Alcohol treatment models

^cSample for Mental Health Treatment model

TABLE 3. Marginal effects (in percentages) for receiving treatment among those with alcohol use disorder from 2008-2017 NSDUH^a

| Variables | No Treatment (Baseline = 93.9%) | | Any Medical Location (Baseline = 5.1%) | | Self-Help Only (Baseline = 1.1%) | |
|---|------------------------------------|--------------|---|--------------|-------------------------------------|--------------|
| | ME. | 95% CI. | ME. | 95% CI. | ME. | 95% CI. |
| Year (reference = 2008-2010) | | | | | | |
| 2011-2013 | 1.4** | 0.4 to 2.4 | -1.3** | -2.2 to -0.4 | -0.1 | -0.5 to 0.3 |
| 2014-2017 | 1.5** | 0.5 to 2.5 | -1.0* | -2.0 to -0.0 | -0.5** | -0.8 to -0.1 |
| Male | -2.0** | -2.8 to -1.1 | 1.4** | 0.7 to 2.2 | 0.5** | 0.2 to 0.9 |
| Age (reference = 18-25) | | | | | | |
| 26-34 | -2.1** | -3.3 to -0.8 | 1.9** | 0.7 to 3.2 | 0.1 | -0.3 to 0.5 |
| 35-49 | -4.2** | -5.8 to -2.6 | 4.1** | 2.6 to 5.6 | 0.1 | -0.4 to 0.5 |
| 50-64 | -3.1* | -5.8 to -0.3 | 3.4* | 0.7 to 6.1 | -0.4 | -0.9 to 0.2 |
| Race/Ethnicity (ref = NH-White) | | | | | | |
| NH-Black | 1.9** | 0.9 to 2.9 | -1.4** | -2.4 to -0.5 | -0.5** | -0.8 to -0.1 |
| Hispanic | 1.7** | 0.7 to 2.7 | -1.8** | -2.7 to -0.9 | 0.1 | -0.4 to 0.5 |
| NH-Other | 1.5* | 0.3 to 2.7 | -1.3* | -2.3 to -0.2 | -0.2 | -0.7 to 0.3 |
| Marital Status (ref = Married) | | | | | | |
| Widowed or Divorced | -2.0** | -3.5 to -0.5 | 2.1** | 0.7 to 3.5 | -0.1 | -0.6 to 0.4 |
| Never Married | -1.3* | -2.6 to -0.1 | 1.5* | 0.3 to 2.7 | -0.1 | -0.6 to 0.3 |
| Income (ref = 75,000+) | | | | | | |
| Less than \$20,000 | -1.3 | -2.9 to 0.2 | 1.5 | 0.0 to 3.0 | -0.2 | -0.6 to 0.2 |
| \$20,000 - \$49,999 | -0.6 | -2.0 to 0.9 | 0.7 | -0.6 to 2.1 | -0.2 | -0.6 to 0.3 |
| \$50,000 - \$74,999 | -1.2 | -3.1 to 0.7 | 1.3 | -0.5 to 3.2 | -0.1 | -0.6 to 0.3 |
| Self-Reported Health Status (ref = Excellent) | | | | | | |
| Very Good | -0.3 | -1.6 to 1.1 | 0.3 | -1.1 to 1.6 | 0.0 | -0.5 to 0.5 |
| Good | -1.4 | -2.9 to 0.1 | 1.3 | -0.1 to 2.8 | 0.1 | -0.5 to 0.6 |
| Fair | 0.2 | -1.3 to 1.8 | -0.7 | -2.1 to 0.8 | 0.4 | -0.4 to 1.3 |
| Poor | 0.8 | -1.5 to 3.1 | -0.3 | -2.5 to 1.9 | -0.5 | -1.5 to 0.4 |
| Insurance Status (ref = Uninsured) | | | | | | |
| Private | -0.5 | -1.7 to 0.6 | 0.7 | -0.4 to 1.7 | -0.1 | -0.6 to 0.3 |
| Medicaid | -2.2* | -3.9 to -0.4 | 2.3* | 0.5 to 4.0 | -0.1 | -0.5 to 0.4 |
| Other Insured | -1.5 | -3.3 to 0.3 | 0.9 | -0.5 to 2.4 | 0.5 | -0.5 to 1.6 |
| Education (ref = High School Graduate) | | | | | | |
| Less than high school | -0.3 | -1.4 to 0.8 | 0.1 | -0.9 to 1.1 | 0.2 | -0.5 to 0.9 |
| Some College | -0.7 | -2.0 to 0.5 | 0.2 | -0.8 to 1.3 | 0.5 | -0.0 to 1.0 |
| College Graduate and above | 0.3 | -1.3 to 1.7 | -0.4 | -1.9 to 1.0 | 0.2 | -0.3 to 0.8 |
| Employment Status (ref = Full-Time) | | | | | | |
| Part-Time | -0.9 | -2.3 to 0.5 | 1.1 | -0.3 to 2.4 | -0.2 | -0.7 to 0.3 |
| Unemployed | -3.6** | -5.4 to -1.9 | 3.8** | 2.1 to 5.5 | -0.1 | -0.8 to 0.5 |
| Other | -2.8** | -4.4 to -1.2 | 3.0** | 1.5 to 4.5 | -0.2 | -0.7 to 0.3 |
| Comorbid substance use disorder | -5.1** | -6.5 to -3.8 | 4.4** | 3.1 to 5.6 | 0.8* | 0.1 to 1.4 |
| Comorbid Mental Illness | -3.6** | -4.5 to -2.7 | 3.1** | 2.3 to 4.0 | 0.5** | 0.2 to 0.9 |
| Alcohol Dependence (ref = Abuse) | -5.7** | -6.6 to -4.8 | 5.0** | 4.2 to 5.9 | 0.7** | 0.3 to 1.0 |
| Health Indicators | | | | | | |
| Diabetes | 0.4 | -1.3 to 2.1 | -0.1 | -1.6 to 1.5 | -0.3 | -1.2 to 0.6 |
| Cirrhosis | -5.0 | -10.8 to 0.8 | 6.1* | 0.3 to 11.8 | -1.1** | -1.2 to -0.9 |
| Hepatitis B or C | -3.7 | -7.6 to 0.1 | 1.1 | -1.4 to 3.6 | 2.7 | -0.2 to 5.5 |
| Asthma | -0.5 | -1.9 to 0.9 | 0.6 | -0.7 to 1.9 | -0.1 | -0.7 to 0.5 |
| HIV or AIDS | -0.0 | -5.6 to 5.6 | -1.9 | -4.6 to 0.9 | 1.9 | -4.0 to 7.8 |
| High Blood Pressure | -0.4 | -1.8 to 1.0 | 0.6 | -0.7 to 1.9 | -0.2 | -0.7 to 0.3 |

^aNSDUH, National Survey of Drug Use and Health

*p<.05, **p<.01

TABLE 4. Marginal effects (in percentages) for receiving mental health treatment among those with alcohol use disorder from 2008-2017 NSDUH^a

| Variables | Mental Health Treatment (Baseline = 23.3%) ^b | |
|---|--|---------------|
| | ME. | 95% CI. |
| Year (reference = 2008-2010) | | |
| 2011-2013 | -0.6 | -1.8 to 0.6 |
| 2014-2017 | -1.3 | -3.0 to 0.4 |
| Male | -11.6** | -13.3 to -9.8 |
| Age (reference = 18-25) | | |
| 26-34 | 4.3** | 2.4 to 6.3 |
| 35-49 | 7.9** | 5.5 to 10.2 |
| 50-64 | 8.1** | 4.9 to 11.3 |
| Race/Ethnicity (ref = NH-White) | | |
| NH-Black | -8.7** | -10.5 to -6.8 |
| Hispanic | -8.0** | -9.8 to -6.1 |
| NH-Other | -7.4** | -9.3 to -5.5 |
| Marital Status (ref = Married) | | |
| Widowed or Divorced | 2.4* | 0.1 to 4.7 |
| Never Married | 0.6 | -1.4 to 2.7 |
| Income (ref = 75,000+) | | |
| Less than \$20,000 | -0.8 | -3.1 to 1.6 |
| \$20,000 - \$49,999 | -2.3* | -4.1 to -0.6 |
| \$50,000 - \$74,999 | -0.3 | -2.4 to 1.9 |
| Self-Reported Health Status (ref = Excellent) | | |
| Very Good | 0.1 | -1.9 to 2.1 |
| Good | 3.0* | 0.7 to 5.2 |
| Fair | 3.2* | 0.3 to 6.1 |
| Poor | 3.1 | -2.9 to 9.0 |
| Insurance Status (ref = Uninsured) | | |
| Private | 6.2** | 4.6 to 7.8 |
| Medicaid | 10.8** | 7.8 to 13.8 |
| Other Insured | 9.7** | 5.4 to 14.1 |
| Education (ref = High School Graduate) | | |
| Less than high school | -2.7* | -5.0 to -0.5 |
| Some College | 1.9* | 0.1 to 3.7 |
| College Graduate and above | 6.0** | 3.6 to 8.4 |
| Employment Status (ref = Full-Time) | | |
| Part-Time | 2.8** | 0.7 to 4.8 |
| Unemployed | 5.7** | 3.2 to 8.2 |
| Other | 7.4** | 5.0 to 9.8 |
| Comorbid substance use disorder | 3.3** | 1.5 to 5.2 |
| Comorbid Mental Illness | 24.0** | 22.5 to 25.6 |
| Alcohol Dependence (ref = Abuse) | 4.1** | 2.6 to 5.7 |
| Health Indicators | | |
| Diabetes | 0.8 | -2.9 to 4.5 |
| Cirrhosis | 1.6 | -8.3 to 11.5 |
| Hepatitis B or C | 5.9 | -0.4 to 12.3 |
| Asthma | 1.5 | -0.5 to 3.6 |
| HIV or AIDS | 13.9* | 1.8 to 26.0 |
| High Blood Pressure | 3.9** | 1.7 to 6.2 |

^a NSDUH, National Survey of Drug Use and Health

^b Sample of 36,707 compared to 36,628 for alcohol treatment

* $p < .05$, ** $p < .01$