

## Observation of $\bar{B}^0 \rightarrow D^0 \pi^0$ and $\bar{B}^0 \rightarrow D^{*0} \pi^0$

T. E. Coan,<sup>1</sup> Y. S. Gao,<sup>1</sup> F. Liu,<sup>1</sup> Y. Maravin,<sup>1</sup> I. Narsky,<sup>1</sup> R. Stroynowski,<sup>1</sup> J. Ye,<sup>1</sup> M. Artuso,<sup>2</sup> C. Boulahouache,<sup>2</sup> K. Bukin,<sup>2</sup> E. Dambasuren,<sup>2</sup> R. Mountain,<sup>2</sup> T. Skwarnicki,<sup>2</sup> S. Stone,<sup>2</sup> J. C. Wang,<sup>2</sup> A. H. Mahmood,<sup>3</sup> S. E. Csorna,<sup>4</sup> I. Danko,<sup>4</sup> Z. Xu,<sup>4</sup> G. Bonvicini,<sup>5</sup> D. Cinabro,<sup>5</sup> M. Dubrovin,<sup>5</sup> S. McGee,<sup>5</sup> A. Bornheim,<sup>6</sup> E. Lipeles,<sup>6</sup> S. P. Pappas,<sup>6</sup> A. Shapiro,<sup>6</sup> W. M. Sun,<sup>6</sup> A. J. Weinstein,<sup>6</sup> G. Masek,<sup>7</sup> H. P. Paar,<sup>7</sup> R. Mahapatra,<sup>8</sup> R. J. Morrison,<sup>8</sup> H. N. Nelson,<sup>8</sup> R. A. Briere,<sup>9</sup> G. P. Chen,<sup>9</sup> T. Ferguson,<sup>9</sup> G. Tishvili,<sup>9</sup> H. Vogel,<sup>9</sup> N. E. Adam,<sup>10</sup> J. P. Alexander,<sup>10</sup> C. Bebek,<sup>10</sup> K. Berkelman,<sup>10</sup> F. Blanc,<sup>10</sup> V. Boisvert,<sup>10</sup> D. G. Cassel,<sup>10</sup> P. S. Drell,<sup>10</sup> J. E. Duboscq,<sup>10</sup> K. M. Ecklund,<sup>10</sup> R. Ehrlich,<sup>10</sup> L. Gibbons,<sup>10</sup> B. Gittelmann,<sup>10</sup> S. W. Gray,<sup>10</sup> D. L. Hartill,<sup>10</sup> B. K. Heltsley,<sup>10</sup> L. Hsu,<sup>10</sup> C. D. Jones,<sup>10</sup> J. Kandaswamy,<sup>10</sup> D. L. Kreinick,<sup>10</sup> A. Magerkurth,<sup>10</sup> H. Mahlke-Krüger,<sup>10</sup> T. O. Meyer,<sup>10</sup> N. B. Mistry,<sup>10</sup> E. Nordberg,<sup>10</sup> M. Palmer,<sup>10</sup> J. R. Patterson,<sup>10</sup> D. Peterson,<sup>10</sup> J. Pivarski,<sup>10</sup> D. Riley,<sup>10</sup> A. J. Sadoff,<sup>10</sup> H. Schwarthoff,<sup>10</sup> M. R. Shepherd,<sup>10</sup> J. G. Thayer,<sup>10</sup> D. Urner,<sup>10</sup> B. Valant-Spaight,<sup>10</sup> G. Viehhauser,<sup>10</sup> A. Warburton,<sup>11</sup> M. Weinberger,<sup>11</sup> S. B. Athar,<sup>11</sup> P. Avery,<sup>11</sup> C. Prescott,<sup>11</sup> H. Stoeck,<sup>11</sup> J. Yelton,<sup>11</sup> G. Brandenburg,<sup>12</sup> A. Ershov,<sup>12</sup> D. Y.-J. Kim,<sup>12</sup> R. Wilson,<sup>12</sup> K. Benslama,<sup>13</sup> B. I. Eisenstein,<sup>13</sup> J. Ernst,<sup>13</sup> G. D. Gollin,<sup>13</sup> R. M. Hans,<sup>13</sup> I. Karliner,<sup>13</sup> N. Lowrey,<sup>13</sup> M. A. Marsh,<sup>13</sup> C. Pflager,<sup>13</sup> C. Sedlack,<sup>13</sup> M. Selen,<sup>13</sup> J. J. Thaler,<sup>13</sup> J. Williams,<sup>13</sup> K. W. Edwards,<sup>14</sup> R. Ammar,<sup>15</sup> D. Besson,<sup>15</sup> X. Zhao,<sup>15</sup> S. Anderson,<sup>16</sup> V. V. Frolov,<sup>16</sup> Y. Kubota,<sup>16</sup> S. J. Lee,<sup>16</sup> S. Z. Li,<sup>16</sup> R. Poling,<sup>16</sup> A. Smith,<sup>16</sup> C. J. Stepaniak,<sup>16</sup> J. Urheim,<sup>16</sup> S. Ahmed,<sup>17</sup> M. S. Alam,<sup>17</sup> L. Jian,<sup>17</sup> M. Saleem,<sup>17</sup> F. Wappler,<sup>17</sup> E. Eckhart,<sup>18</sup> K. K. Gan,<sup>18</sup> C. Gwon,<sup>18</sup> T. Hart,<sup>18</sup> K. Honscheid,<sup>18</sup> D. Hufnagel,<sup>18</sup> H. Kagan,<sup>18</sup> R. Kass,<sup>18</sup> T. K. Pedlar,<sup>18</sup> J. B. Thayer,<sup>18</sup> E. von Toerne,<sup>18</sup> M. M. Zoeller,<sup>18</sup> S. J. Richichi,<sup>19</sup> H. Severini,<sup>19</sup> P. Skubic,<sup>19</sup> S. A. Dytman,<sup>20</sup> S. Nam,<sup>20</sup> V. Savinov,<sup>20</sup> S. Chen,<sup>21</sup> J. W. Hinson,<sup>21</sup> J. Lee,<sup>21</sup> D. H. Miller,<sup>21</sup> V. Pavlunin,<sup>21</sup> E. I. Shibata,<sup>21</sup> I. P. J. Shipsey,<sup>21</sup> D. Cronin-Hennessy,<sup>22</sup> A. L. Lyon,<sup>22</sup> C. S. Park,<sup>22</sup> W. Park,<sup>22</sup> and E. H. Thorndike<sup>22</sup>

(CLEO Collaboration)

<sup>1</sup>*Southern Methodist University, Dallas, Texas 75275*

<sup>2</sup>*Syracuse University, Syracuse, New York 13244*

<sup>3</sup>*University of Texas–Pan American, Edinburg, Texas 78539*

<sup>4</sup>*Vanderbilt University, Nashville, Tennessee 37235*

<sup>5</sup>*Wayne State University, Detroit, Michigan 48202*

<sup>6</sup>*California Institute of Technology, Pasadena, California 91125*

<sup>7</sup>*University of California, San Diego, La Jolla, California 92093*

<sup>8</sup>*University of California, Santa Barbara, California 93106*

<sup>9</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213*

<sup>10</sup>*Cornell University, Ithaca, New York 14853*

<sup>11</sup>*University of Florida, Gainesville, Florida 32611*

<sup>12</sup>*Harvard University, Cambridge, Massachusetts 02138*

<sup>13</sup>*University of Illinois, Urbana-Champaign, Illinois 61801*

<sup>14</sup>*Carleton University, Ottawa, Ontario, Canada K1S 5B6*

*and the Institute of Particle Physics, Canada*

<sup>15</sup>*University of Kansas, Lawrence, Kansas 66045*

<sup>16</sup>*University of Minnesota, Minneapolis, Minnesota 55455*

<sup>17</sup>*State University of New York at Albany, Albany, New York 12222*

<sup>18</sup>*The Ohio State University, Columbus, Ohio 43210*

<sup>19</sup>*University of Oklahoma, Norman, Oklahoma 73019*

<sup>20</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260*

<sup>21</sup>*Purdue University, West Lafayette, Indiana 47907*

<sup>22</sup>*University of Rochester, Rochester, New York 14627*

(Received 22 October 2001; published 24 January 2002)

We have studied the color-suppressed hadronic decays of neutral  $B$  mesons into the final states  $D^{(*)0} \pi^0$ . Using  $9.67 \times 10^6$   $B\bar{B}$  pairs collected with the CLEO detector, we observe the decays  $\bar{B}^0 \rightarrow D^0 \pi^0$  and  $\bar{B}^0 \rightarrow D^{*0} \pi^0$  with the branching fractions  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \pi^0) = (2.74_{-0.32}^{+0.36} \pm 0.55) \times 10^{-4}$  and  $\mathcal{B}(\bar{B}^0 \rightarrow D^{*0} \pi^0) = (2.20_{-0.52}^{+0.59} \pm 0.79) \times 10^{-4}$ . The first error is statistical and the second systematic. The statistical significance of the  $D^0 \pi^0$  signal is  $12.1\sigma$  ( $5.9\sigma$  for  $D^{*0} \pi^0$ ). Utilizing the  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  branching fractions we determine the strong phases  $\delta_{I,D^{(*)}}$  between isospin 1/2 and 3/2 amplitudes in the  $D\pi$  and  $D^*\pi$  final states to be  $\cos\delta_{I,D} = 0.89 \pm 0.08$  and  $\cos\delta_{I,D^*} = 0.89 \pm 0.08$ , respectively.

DOI: 10.1103/PhysRevLett.88.062001

PACS numbers: 13.25.Hw

The decay  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  proceeds predominantly through the internal spectator diagram shown in Fig. 1. This diagram is color suppressed, since the color of the quark pair originating from the  $W$  decay must match the color of the other quark pair. The rate of  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  relative to  $B^- \rightarrow D^{(*)0} \pi^-$  is suppressed, crudely, by one factor of  $(1/3)^2$ , and an additional  $(1/\sqrt{2})^2$  due to the projection of the  $d\bar{d}$  state onto the pion wave function [1]. This gives a total suppression of  $1/18$  compared to the color-favored decay modes. Detailed theoretical calculations [2] predict an even larger suppression of about a factor of  $1/50$ .

So far the only established color-suppressed decays are two-body  $B$  decays into charmonium plus neutral hadrons. A measurement of  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  is therefore a benchmark test for theoretical models of hadronic  $B$  decays [1,2]. An investigation of color-suppressed decays into a  $D$  meson and light neutral mesons other than a  $\pi^0$  is currently under way and will be addressed in a future publication.

The observation of  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  completes the measurement of  $D^{(*)} \pi$  final states and allows us to extract the strong phase difference between isospin  $1/2$  and  $3/2$  amplitudes [2,3].

In this Letter we present the observation of  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$ , superseding the limits from our previous publication [4]. Our new analysis has significantly increased statistics and is based on a data sample with improved calibration and track reconstruction. The background shapes and the signal separation power have also been significantly improved. Charge conjugates are implied throughout this Letter. Our analysis uses  $e^+e^-$  annihilation data recorded with the CLEO detector at the Cornell Electron Storage Ring. The integrated luminosity of our data sample is  $9.15 \text{ fb}^{-1}$  for data collected on the  $Y(4S)$  (on-resonance), corresponding to  $9.67 \times 10^6$   $B\bar{B}$  pairs, and  $4.35 \text{ fb}^{-1}$  collected 60 MeV below the  $B\bar{B}$  threshold (off-resonance), which is used for background studies.

CLEO is a general purpose solenoidal magnet detector. Data were recorded with two detector configurations, CLEO II and CLEO II.V [5,6]. Cylindrical drift chambers in a 1.5 T solenoidal magnetic field measure momentum and specific ionization ( $dE/dx$ ) of charged particles. Photons are detected using a CsI(Tl) crystal electromagnetic calorimeter, consisting of a barrel-shaped central part of




FIG. 1. Diagram for the color-suppressed decay  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$ .

6144 crystals and 1656 crystals in the forward regions of the detector (end caps). In the II.V configuration the innermost chamber was replaced by a three-layer, double-sided silicon microvertex detector, and the main drift chamber gas was changed from an argon-ethane to a helium-propane mixture. As a result of these modifications, the CLEO II.V part of the data ( $2/3$  of the total sample) has improved momentum resolution and particle identification.

$B$  mesons were reconstructed by selecting high-momentum  $D^{(*)0}$  and  $\pi^0$  mesons. Track quality requirements are imposed on charged tracks and the purity of pions and kaons is improved by using  $dE/dx$  information whenever available. The  $\pi^0$  candidates are reconstructed from isolated electromagnetic clusters of at least 30 MeV in the barrel region and 50 MeV in the end caps. The mass resolution is 8 MeV in the barrel region and 10 MeV in the end caps. We require that the candidate's mass is within 2.5 standard deviations ( $\sigma$ ) of the nominal  $\pi^0$  mass. Prompt  $\pi^0$ 's from  $B$  decays are required to have a momentum larger than 1.8 GeV.

$D^0$  mesons are selected in the decay modes  $D^0 \rightarrow K^- \pi^+$ ,  $D^0 \rightarrow K^- \pi^+ \pi^0$ , and  $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ . The invariant mass of the  $D$  daughter particles is required to be within  $2.5\sigma$  of the known  $D^0$  mass. The mass resolution depends on the decay mode and is between 6 and 12 MeV. The momentum of the  $D^0$  is required to be larger than 1.65 GeV. In the  $D^0 \rightarrow K^- \pi^+ \pi^0$  mode we suppress combinatorial background by using only certain regions of the Dalitz plane.

$D^{*0}$  mesons are selected in the decay modes  $D^{*0} \rightarrow D^0 \pi^0$  and  $D^{*0} \rightarrow D^0 \gamma$ . To reduce the combinatorial background in the decay mode  $D^{*0} \rightarrow D^0 \gamma$  we require that the  $D^0$  decays into  $K^- \pi^+$ . We require that the mass difference  $m_{D^{*0}} - m_{D^0}$  is within  $2.5\sigma$  of the known value and that the momentum of the  $D^{*0}$  is larger than 1.8 GeV. The kinematic resolution of  $\pi^0$  and  $D^{(*)0}$  candidates is improved by a mass-constrained kinematic fit.

$B$  decay candidates are selected from  $\pi^0$  and  $D^{(*)0}$  pairings that have no electromagnetic clusters in common.  $B$  candidates are identified using a beam-constrained mass  $M_B = \sqrt{E_{\text{beam}}^2 - p_B^2}$ , where  $E_{\text{beam}}$  is the beam energy and  $p_B$  is the  $B$  candidate momentum, and an energy difference  $\Delta E = E_D + E_{\pi^0} - E_{\text{beam}}$ , where  $E_D$  and  $E_{\pi^0}$  are the energies of the  $D^{(*)0}$  and  $\pi^0$ . The resolution in  $M_B$  depends on the  $D$  decay mode and is between 3.5 and 4 MeV. The resolution is dominated by the beam energy spread and the  $\pi^0$  energy resolution. The resolution in  $\Delta E$  is between 35 and 40 MeV. The energy resolution is slightly asymmetric due to the energy loss out of the back of the CsI crystals. The mass-constrained kinematic fit to the pion 4-momentum compensates for most of this effect. We accept  $B$  candidates with  $M_B$  above 5.24 GeV and  $|\Delta E| < 300$  MeV. To better suppress background from  $e^+e^- \rightarrow q\bar{q}$  events (continuum background), several event shape variables are combined into a Fisher discriminant  $\mathcal{F}_D$  [7]. For  $B\bar{B}$

events (continuum background), the  $\mathcal{F}_D$  distribution is almost a Gaussian and has its maximum at 0.42 (0.57). The standard deviation is 0.11 (0.12). The separation between the  $B\bar{B}$  and continuum distributions is  $1.3\sigma$ . We reject clear continuum events by requiring  $\mathcal{F}_D < 1$ . For each candidate we calculate the sphericity vectors [8] of the  $B$  daughter particles and of the rest of the event. We require the cosine of the angle between these two vectors to be within  $-0.8$  and  $0.8$ . The distribution of this angle is strongly peaked at  $\pm 1$  for continuum background and is nearly flat for  $B\bar{B}$  events.

The number of signal events in the sample is obtained from unbinned, extended maximum likelihood fits. The free parameters of the fits are the number of signal events, background from  $B$  decays ( $B\bar{B}$ ), and from continuum  $e^+e^-$  annihilation (continuum). Four variables are used as input to the maximum likelihood fit: the beam-constrained mass  $M_B$ , the energy difference  $\Delta E$ , the Fisher discriminant  $\mathcal{F}_D$ , and the cosine of the decay angle of the  $B$   $\cos\theta_{B\text{Hel}}$ , defined as the angle between the  $D^{(*)}$  momentum and the  $B$  flight direction calculated in the  $B$  rest frame.

In each of the fits, the likelihood of the  $B$  candidate is the sum of probabilities for the signal and two background hypotheses with relative weights maximizing the likelihood. The probability of a particular hypothesis is the product of probability density functions (PDFs) for each of the input variables. The PDFs for  $M_B$  are represented by a bifurcated Gaussian [9] for signal, an empirical shape,  $M_B\sqrt{(1-x^2)}\exp[-E_{\text{fact}}(1-x^2)]$ , with  $x = M_B/E_{\text{beam}}$ , for continuum and a Gaussian on top of the empirical background shape (the parameters of the empirical shapes for  $B\bar{B}$  and continuum background are different) for  $B\bar{B}$ ; the PDFs for  $\Delta E$  are the sum of two Gaussians with a common mean (signal), first-order polynomial (continuum) and a sum of two Gaussians plus a first-order polynomial ( $B\bar{B}$ ); the PDFs for  $\mathcal{F}_D$  are the sum of two Gaussians with a common mean; and the PDFs for  $\cos\theta_{B\text{Hel}}$  are second-order polynomials. The PDF parameters are determined from off-resonance CLEO data (continuum) and from high-statistics Monte Carlo (MC) samples (signal and  $B\bar{B}$ ).

Monte Carlo experiments are generated to test the fitting procedure and to obtain the relation between fit yield and signal branching fractions. The experiments are repeated several hundred times with different Monte Carlo test samples randomly selected from high-statistics MC samples.

We summarize the results of the fits to CLEO data in Table I. We give results for all  $B$  decay modes, corresponding  $D$  decay modes, and the combination of all  $D$  decay modes. We combine the results for different  $D$  decay modes by adding the log likelihood as a function of the branching fraction. Branching fractions for each mode are obtained via

$$\mathcal{B}(\bar{B}^0 \rightarrow D^{(*)0}\pi^0) = \frac{\text{Yield}_{\text{fit}}}{\varepsilon \mathcal{B}(D^{(*)})N(B^0, \bar{B}^0)}.$$

The number of  $B^0$  plus  $\bar{B}^0$ ,  $N(B^0, \bar{B}^0) = 9.67 \pm 0.10$  M, is derived assuming equal branching fractions for charged and neutral  $B$  meson decays [10]. The uncertainty in the branching fractions of the  $Y(4S)$  is taken into account in the systematics. The significances of the observed signals in the seven fits is determined from the change in  $-2\log\mathcal{L}$  when refit with the signal yield constrained to zero: significance =  $\sqrt{2(\log\mathcal{L} - \log\mathcal{L}_{N_{\text{sig}}=0})}$ . We obtain a total significance of  $12.1\sigma$  for  $\bar{B}^0 \rightarrow D^0\pi^0$  and  $5.9\sigma$  for  $\bar{B}^0 \rightarrow D^{*0}\pi^0$ . Varying the PDF shapes within the systematic errors to obtain the lowest signal yield, the statistical significance is reduced to  $9.4\sigma$  ( $\bar{B}^0 \rightarrow D^0\pi^0$ ) and  $4.2\sigma$  ( $\bar{B}^0 \rightarrow D^{*0}\pi^0$ ). We obtain branching fractions of  $\mathcal{B}(\bar{B}^0 \rightarrow D^0\pi^0) = (2.74_{-0.32}^{+0.36} \pm 0.55) \times 10^{-4}$  and  $\mathcal{B}(\bar{B}^0 \rightarrow D^{*0}\pi^0) = (2.20_{-0.52}^{+0.59} \pm 0.79) \times 10^{-4}$ . The first error is statistical and the second error is systematic. Our result for  $\bar{B}^0 \rightarrow D^0\pi^0$  is higher than the previous CLEO upper limit [4]. We ascribe this disagreement, which is of the order of  $3.1\sigma$ , partly to a statistical fluctuation and partly to the description of the  $\Delta E$  background in the old CLEO publication.

We consider sources of systematic uncertainties from the PDF shapes,  $D$  and  $Y(4S)$  branching ratios, luminosity, possible fit bias,  $B$  candidate reconstruction, and cross

TABLE I. Fit yields for all decay modes. Our results are based on the  $D$  branching ratios given in column 5 [10]. Our measurement of the  $B$  branching ratios is given in the last column.

| Mode                                                      |                                                              | Fit yield<br>(events)  | Significance ( $\sigma$ ) | $\varepsilon$<br>(%) | $\mathcal{B}(D^{(*)})$<br>(%) | $\mathcal{B}(D^{(*)0}\pi^0)$<br>( $10^{-4}$ ) |
|-----------------------------------------------------------|--------------------------------------------------------------|------------------------|---------------------------|----------------------|-------------------------------|-----------------------------------------------|
| $\bar{B}^0 \rightarrow D^0\pi^0$                          | $D^0 \rightarrow K^-\pi^+$                                   | $37.5_{-6.9}^{+7.2}$   | 8.5                       | 37.1                 | 3.82                          | $2.74 \pm 0.53$                               |
|                                                           | $D^0 \rightarrow K^-\pi^+\pi^0$                              | $42.1_{-8.6}^{+9.0}$   | 6.8                       | 13.5                 | 12.94                         | $2.49 \pm 0.53$                               |
|                                                           | $D^0 \rightarrow K^-(3\pi)^+$                                | $44.6_{-10.2}^{+10.7}$ | 5.3                       | 19.0                 | 7.48                          | $3.25 \pm 0.78$                               |
| Averaged $\mathcal{B}(\bar{B}^0 \rightarrow D^0\pi^0)$    |                                                              |                        | 12.1                      |                      |                               | $2.74_{-0.32}^{+0.36}$                        |
| $\bar{B}^0 \rightarrow D^{*0}\pi^0$                       | $D^{*0} \rightarrow D^0\pi^0, D^0 \rightarrow K^-\pi^+$      | $6.8_{-2.8}^{+3.2}$    | 2.4                       | 15.3                 | 2.36                          | $1.95 \pm 0.91$                               |
|                                                           | $D^{*0} \rightarrow D^0\pi^0, D^0 \rightarrow K^-\pi^+\pi^0$ | $7.3_{-3.6}^{+4.0}$    | 2.8                       | 5.5                  | 8.01                          | $1.72 \pm 0.94$                               |
|                                                           | $D^{*0} \rightarrow D^0\pi^0, D^0 \rightarrow K^-(3\pi)^+$   | $8.0_{-3.7}^{+4.2}$    | 3.1                       | 8.1                  | 4.63                          | $2.21 \pm 1.15$                               |
|                                                           | $D^{*0} \rightarrow D^0\gamma, D^0 \rightarrow K^-\pi^+$     | $6.4_{-2.7}^{+3.0}$    | 3.4                       | 11.4                 | 1.46                          | $3.99 \pm 1.89$                               |
| Averaged $\mathcal{B}(\bar{B}^0 \rightarrow D^{*0}\pi^0)$ |                                                              |                        | 5.9                       |                      |                               | $2.20_{-0.52}^{+0.59}$                        |




FIG. 2. Distribution of fit input variables for  $\bar{B}^0 \rightarrow D^0 \pi^0$ . The results of the unbinned, extended maximum likelihood fit are shown as the full line. The dotted line represents the fitted continuum and the dashed line is the fit result for the sum of  $B\bar{B}$  and continuum background. To enhance the signal for display purposes, the fit results are projected into the  $M_B$ - $\Delta E$  signal region  $-0.05 < \Delta E < 0.05$  GeV,  $5.275 < M_B < 5.285$  GeV.

feed between different modes. The dominant systematic uncertainty comes from the PDF shapes. The systematic uncertainty on the shapes is derived by varying the PDF shapes within the statistical errors of the fit parametrization, as well as comparing the CLEO data in the  $\Delta E$  and  $M_B$  sideband regions to the PDF shapes and taking differences as systematic errors. Figures 2 and 3 show our results for  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  with the number of signal,  $B\bar{B}$ , and continuum background as free parameters of the fit. The fit result is projected into a signal region, defined in the  $M_B$ - $\Delta E$  plane as  $-0.05 < \Delta E < 0.05$  GeV,  $5.275 < M_B < 5.285$  GeV. The fit results describe the data well. The background in the sidebands is also well modeled by the fit.

The observation of  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$  completes the measurement of  $D^{(*)} \pi$  final states. This allows us to calculate the relative phase between the isospin 1/2 and 3/2 amplitudes in the  $D^{(*)} \pi$  system. The basic relation can be expressed in an amplitude triangle:  $\mathcal{A}(\bar{D}^0 \pi^+) = \mathcal{A}(D^- \pi^+) + \sqrt{2} \mathcal{A}(\bar{D}^0 \pi^0)$ , following the formulation in [3]. With the Particle Data Group values [10]  $\mathcal{B}(\bar{D}^0 \pi^+) = (53 \pm 5) \times 10^{-4}$ ,  $\mathcal{B}(D^- \pi^+) = (30 \pm 4) \times 10^{-4}$ ,  $\mathcal{B}(\bar{D}^{*0} \pi^+) = (46 \pm 4) \times 10^{-4}$ ,  $\mathcal{B}(D^{*-} \pi^+) = (27.6 \pm 2.1) \times 10^{-4}$ ,  $\tau(B^+)/\tau(B^0) = 1.073 \pm 0.027$ , and our measurement of  $\bar{B}^0 \rightarrow D^{(*)0} \pi^0$ , we determine the relative phase between the isospin amplitudes to be  $\cos \delta_{I,D} = 0.89 \pm 0.08$  for the  $D\pi$  final state and  $\cos \delta_{I,D^*} = 0.89 \pm 0.08$  for  $D^* \pi$ . The ratios of isospin amplitudes




FIG. 3. Distribution of fit input variables for  $\bar{B}^0 \rightarrow D^{*0} \pi^0$ . The results of the unbinned, extended maximum likelihood fit are shown as the full line. The dotted line represents the fitted continuum and the dashed line is the fit result for the sum of  $B\bar{B}$  and continuum background. To enhance the signal for display purposes, the fit results are projected into the  $M_B$ - $\Delta E$  signal region.

$A_{1/2}/A_{3/2}$  are  $0.70 \pm 0.11$  ( $D\pi$ ) and  $0.74 \pm 0.08$  ( $D^* \pi$ ). A similar calculation has been performed in [12] using our preliminary results [13] and preliminary results obtained by the Belle Collaboration [14].

Models of hadronic  $B$  decay [2] have successfully described experimental results using two phenomenological parameters,  $a_1$  and  $a_2$ , that characterize nonfactorizable contributions. Both are believed to be process dependent, but so far experimental data have been consistent with universal values for  $a_1$  and  $a_2$ . Recent work by Beneke *et al.* [15] has shown that  $a_1$  is only slightly process dependent. Based on our  $\bar{B}^0 \rightarrow D^0 \pi^0$  measurement, we derive a value  $a_2 = 0.57 \pm 0.06$ . Comparing our result to the  $a_2$  value from two-body  $B$  decays to charmonium,  $a_2 = 0.29$  [2], the process dependence of  $a_2$  is favored [12].

To summarize, we observed the color-suppressed decays  $\bar{B}^0 \rightarrow D^0 \pi^0$  and  $\bar{B}^0 \rightarrow D^{*0} \pi^0$ . The number of signal events in our data sample was obtained from an unbinned extended maximum likelihood fit in four variables. The measurements of the two branching fractions are  $\mathcal{B}(\bar{B}^0 \rightarrow D^0 \pi^0) = (2.74^{+0.36}_{-0.32} \pm 0.55) \times 10^{-4}$  and  $\mathcal{B}(\bar{B}^0 \rightarrow D^{*0} \pi^0) = (2.20^{+0.59}_{-0.52} \pm 0.79) \times 10^{-4}$ . The first error is statistical and the second systematic. The statistical significance of the  $D^0 \pi^0$  signal is  $12.1\sigma$  ( $5.9\sigma$  for  $D^{*0} \pi^0$ ).

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the National Science

Foundation, the U.S. Department of Energy, the Research Corporation, and the Texas Advanced Research Program.

- 
- [1] T. E. Browder, K. Honscheid, and D. Pedrini, *Annu. Rev. Nucl. Part. Sci.* **46**, 395 (1996).
- [2] M. Neubert and B. Stech, in *Heavy Flavors*, edited by A. J. Buras and M. Lindner (World Scientific, Singapore, 1998), 2nd ed.
- [3] J. L. Rosner, *Phys. Rev. D* **60**, 074029 (1999).
- [4] CLEO Collaboration, B. Nematy *et al.*, *Phys. Rev. D* **57**, 5363 (1998).
- [5] CLEO Collaboration, Y. Kubota *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **320**, 66 (1992).
- [6] T. Hill, *Nucl. Instrum. Methods Phys. Res., Sect. A* **418**, 32 (1998).
- [7] CLEO Collaboration, D. Cronin-Hennessy *et al.*, *Phys. Rev. Lett.* **85**, 515 (2000).
- [8] S. L. Wu, *Phys. Rep. C* **107**, 59 (1984).
- [9] A bifurcated Gaussian is a Gaussian with different widths for the left and right parts of the curve.
- [10] Particle Data Group, D. E. Groom *et al.*, *Eur. Phys. J. C* **15**, 1 (2000); 2001 partial update for edition 2002 (URL: <http://pdg.lbl.gov>).
- [11] CLEO Collaboration, J. P. Alexander *et al.*, *Phys. Rev. Lett.* **86**, 2737 (2001).
- [12] M. Neubert and A. Petrov, *Phys. Lett. B* **519**, 50 (2001).
- [13] E. von Toerne, in Proceedings of International Europhysics Conference on High Energy Physics, Budapest, 2001 (to be published).
- [14] R-S. Lu, hep-ex/0107048 [in Proceedings of International Europhysics Conference on High Energy Physics, Budapest (to be published)]; final results by the Belle Collaboration, K. Abe *et al.*, *Phys. Rev. Lett.* **88**, 052002 (2002).
- [15] M. Beneke *et al.*, *Nucl. Phys.* **B591**, 313 (2000).