



The Genetics of Self-Compatibility and Associated Floral Characters in *Tolpis* (Asteraceae) in the Canary Islands

Author(s): Fabiola Soto-Trejo, John K. Kelly, Jenny K. Archibald, Mark E. Mort, Arnaldo Santos-Guerra, and Daniel J. Crawford

Source: *International Journal of Plant Sciences*, Vol. 174, No. 2 (February 2013), pp. 171-178

Published by: [The University of Chicago Press](#)

Stable URL: <http://www.jstor.org/stable/10.1086/668788>

Accessed: 20/03/2014 14:35

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



The University of Chicago Press is collaborating with JSTOR to digitize, preserve and extend access to *International Journal of Plant Sciences*.

<http://www.jstor.org>

THE GENETICS OF SELF-COMPATIBILITY AND ASSOCIATED FLORAL CHARACTERS IN *TOLPIS* (ASTERACEAE) IN THE CANARY ISLANDS

Fabiola Soto-Trejo,* John K. Kelly,† Jenny K. Archibald,†‡ Mark E. Mort,†‡
Arnoldo Santos-Guerra,§ and Daniel J. Crawford¹†‡

*Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México, Avenida de Los Barrios s/n, 54090 Tlalnepantla, Estado de México; †Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, Kansas 66045, U.S.A.; ‡Biodiversity Institute, University of Kansas, Lawrence, Kansas 66045, U.S.A.; and §Unidad de Botánica, (ICIA) Retama 2, Puerto de la Cruz, Tenerife, Canary Islands, Spain

Members of the genus *Tolpis* (Asteraceae) endemic to the Canary Islands comprise a monophyletic group with the dispersal of one species to the Cape Verde Islands. Most species are self-incompatible or pseudo-self-compatible perennials. However, one species, *Tolpis coronopifolia*, consists of self-compatible annual plants with several floral features typical of the selfing syndrome. The evolution of self-compatibility and the selfing syndrome was studied by producing synthetic hybrids between self-incompatible/pseudo-self-compatible and self-compatible plants to determine the genetic architecture of breeding system and floral trait differences. There was a correlation between self-compatibility and selfing floral traits in the F₂ generation. Self-incompatibility in *Tolpis* appears to be controlled by a locus of major effect but with modifier loci affecting seed set. Segregation of floral traits indicates that they are controlled by multiple loci. The high molecular similarity between plants with the two breeding systems suggests that divergence of the self-compatible *T. coronopifolia* from self-incompatible/pseudo-self-compatible ancestors has been recent. The association between breeding system and floral features likely results from pleiotropy or close linkage and may have facilitated the rapid evolution of the morphologically distinct self-compatible *T. coronopifolia*.

Keywords: Asteraceae, Canary Islands, self-compatibility, *Tolpis*.

Online enhancements: appendix figures.

Introduction

Self-incompatibility (SI; also used to indicate self-incompatible) is the most common mechanism for enforcing outcrossing in flowering plants (Igic et al. 2008). However, the loss of SI to produce self-compatible (SC; also used to designate self-compatibility) plants is also quite common and has occurred repeatedly in angiosperms (Goodwillie 1999; Barrett 2002; Beck et al. 2006; Igic et al. 2008; Foxe et al. 2010). The two primary advantages commonly attributed to selfers over outcrossers is an inherent transmission advantage (Fisher 1941) and the ability to reproduce sexually without mates (Kaliz et al. 2004; Charlesworth 2006). Higher selfing may also decrease hybridization with closely related sympatric species (Antonovics 1968; Levin 1978, 2000 [p. 11]; Fishman and Wyatt 1999; Wendt et al. 2002; Grossenbacher and Whittall 2011). Potential downsides of selfing include inbreeding depression and reduced genetic variation, since selfing increases homozygosity and lowers the effective population size (Goodwillie et al. 2005; Yang and Hodges 2010).

Selfing facilitates colonization following dispersal because one individual can initiate a sexually reproducing population in the absence of mates and pollination vectors (Baker 1955;

Stebbins 1957; Pannell and Barrett 1998). Stebbins (1957) viewed this idea of such general significance that he called it Baker's law. Selfing ability may be especially critical in the persistence of plants on isolated oceanic islands because ancestral colonizers likely arrived by rare, long-distance dispersal events (Cowie and Holland 2006). However, just as Allard (1965) and others had argued for the importance of genetic variation in selfing species, Carlquist (1966) suggested that Baker's law may not be of general application to the insular situation because of the low genetic variation present in a colonizer originating from a selfing source population. He thus emphasized that the higher genetic diversity typical of outcrossed populations is crucial for the persistence and diversification of insular lineages. Reviews of the literature by Ehrendorfer (1979) and Barrett (1996) supported the Baker (1955) hypothesis because the majority of endemic flowering plants examined from islands are SC. Two well-studied island systems with a high frequency of SC species are the Galápagos (McMullen 1987, 1990) and New Zealand (Webb and Kelly 1993). More recent compilations for the Robinson Crusoe (Juan Fernandez) Islands likewise document the prevalence of SC species (Anderson et al. 2001; Bernardello et al. 2001). However, some of the most successful insular radiations in Asteraceae, the plant family with the highest number of oceanic island endemics (Crawford et al. 2009b, 2011), originated from SI ancestors in which there was likely some "leakiness" in the system. That is, the plants were pseudo-

¹ Author for correspondence; e-mail: dcrawfor@ku.edu.

Manuscript received March 2012; revised manuscript received August 2012.

self-compatible (PSC; also used to designate pseudo-self-compatibility; Levin 1996; reviewed in Crawford et al. 2009b, 2011). Because PSC colonizers originated from potentially outcrossing source populations, they would likely carry more genetic variation than a propagule from a highly selfing population, yet they could set some selfed seed (Levin 1996; Crawford et al. 2008).

True SC is achieved via irreversible breakdown of a functioning S-locus, resulting in the potential to produce high levels of selfed seed (Ilgic et al. 2008; Pettengill and Moeller 2011). In addition, a suite of floral characters often evolves in association with SC (Ornduff 1969; Slotte et al. 2012). Available evidence indicates that many of the SC species that have been studied on oceanic islands originated from SC colonizers (Carlquist 1974, chap. 9; McMullen 1987, 1990; Sun and Ganders 1988; Anderson et al. 2001; Bernardello et al. 2001). Despite the apparent prevalence of SC species, there are at least some reports of various levels of PSC in island endemics, such as the silversword alliance in Hawaii (Carr et al. 1986;ARRIER et al. 1999; Baldwin 2003; but see Bainbridge and Baldwin 2010) and *Scalesia* from the Galápagos Islands (Nielsen et al. 2000, 2003). A recent study of the Canarian clade of *Tolpis* (Asteraceae) has shown that, with one exception, species are

SI or PSC (i.e., SI-PSC). The one exception is the SC species *Tolpis coronopifolia* (Crawford et al. 2008), which has several components of the selfing syndrome, including much smaller capitula with fewer, smaller florets (fig. 1). Also, unlike other Canary Island endemic *Tolpis* species, *T. coronopifolia* is an annual (or occasionally biennial) rather than perennial. The SC condition is likely derived, as it is in most other flowering plants (Beck et al. 2006; Ilgic et al. 2008), and evolved in situ in the Canaries. We are unaware of the in situ evolution of SC and the selfing floral syndrome during the radiation of other island lineages in Asteraceae; one species in the silversword alliance is SC but does not exhibit the selfing syndrome (G. D. Carr, personal communication).

The genus *Tolpis* is a good system to address questions about the evolution of SC and the selfing syndrome in an insular lineage. While the contrasting floral features of closely related outcrossing and selfing species have long been recognized (Ornduff 1969), little is known about the dynamics of change in a variety of floral features associated with the breakdown of SI (but see Slotte et al. 2012). Because the SC *T. coronopifolia* can easily be hybridized with SI-PSC taxa of *Tolpis* (Crawford et al. 2009a), it is feasible to study the segregation of characters in F₂ interspecific hybrids. Molecular divergence between *T.*



Fig. 1 Capitula of *Tolpis* sp. nov. 1 (population Santos-Guerra 4; A), *Tolpis coronopifolia* (population Crawford et al. 1850; B), an F₁ hybrid (C), and two F₂ hybrids (D) showing the range of size variation in this generation.

coronopifolia and the other species in the Canary clade is minimal to nonexistent, using data sets normally effective in resolving relationships among congeneric species. These include inter-simple sequence repeat markers (Archibald et al. 2006), allozymes (Crawford et al. 2006), and nuclear internal transcribed spacer (M. E. Mort, unpublished data; R. K. Jansen, personal communication) and noncoding plastid DNA sequences (Mort et al. 2010), suggesting that it has diverged from its SI ancestors recently and rapidly.

Available data from reproductive biology, floral morphology, and molecular markers suggest that SC and several floral traits typically associated with selfing have evolved rapidly in *Tolpis* in the Canary Islands. The purpose of this study was to examine the pattern of inheritance of SC and floral morphology in hybrids between SC and SI-PSC plants of *Tolpis*. Of particular interest was testing the hypothesis that rapid evolution of an SC species of *Tolpis* may have been facilitated by the genetic architecture of the selfing suite of traits.

Material and Methods

Hybrids were made between an undescribed SI-PSC species of Canary *Tolpis* and the SC species *Tolpis coronopifolia*

(Crawford et al. 2008). The SI-PSC species belongs to the *Tolpis laciniata* complex and has been designated *Tolpis* sp. nov. 1 (Crawford et al. 2009a). The methods for determining the breeding systems of the parental species were described by Crawford et al. (2008). Briefly, capitula were enclosed in fine netting before anthesis, and pollen was distributed over the stigmatic surfaces of the same capitula on successive days until the capitula closed. Self seed set was determined by the proportion of plump, black fruits with embryos. Plants grown from wild-collected seed served as the parents for hybridization, including seeds from population Santos-Guerra 4 (on La Palma) for the SI-PSC parent and seeds from population Crawford et al. 1850 (on Tenerife) for the SC parent. These seeds and all subsequent progeny were reared in consistent environmental conditions in the University of Kansas greenhouses. Plants used to generate F₁ hybrids had proportion pollen viability exceeding 0.80. Crosses were effected by bagging capitula before anthesis and rubbing them together on successive days until fruit was set, which is indicated by the onset of senescence of the peduncle below the capitulum. The seedlings of F₁ hybrids could be detected by intermediate leaf morphology (fig. 2). When SI-PSC plants were the egg parent, nearly all progeny were hybrids, but with SC *T. coronopifolia* as egg parent, fewer than 15% of the progeny were




Fig. 2 Leaves of *Tolpis* sp. nov. 1 (population Santos-Guerra 4; A), *Tolpis coronopifolia* (population Crawford et al. 1850; B), and two F₁ hybrid individuals (C).

hybrids, and the rest were selfs. Proportion pollen viability of the F_1 hybrids used to generate the F_2 generation was higher than 0.90. Vouchers for parental and F_1 hybrids are deposited in the McGregor Herbarium (KANU) at the University of Kansas.

Nine F_2 families were generated by bagging F_1 plants before anthesis, collecting the resulting fruits, and germinating them. From the resulting plants, a total of 129 individuals were measured for the following floral characters: number of florets per capitulum, diameter of capitula, and floret, anther, and stigma lengths. For the first two characters, the average of two capitula per plant was used, and for the latter three characters, the average of five florets from each of two capitula was taken. Pollen viability for each plant was determined by examining at least 200 pollen grains stained with lactophenol aniline blue (Kearns and Inouye 1993; Crawford et al. 2008). Self seed set for F_2 individuals was determined by the same methods used to assess self seed set in the parental species. Large, plump, black to brown-black fruits containing viable embryos were easily distinguished from shriveled, light brown to tan fruits with no embryos (Crawford et al. 2008). Five tan, shrunken fruits from two plants of each of the nine F_2 families were dissected, and there was no evidence of embryos. The pollen:ovule ratio (average of two capitula) was determined for 126 plants, and proportion pollen viability and proportion self seed set (in 124 plants) were calculated by pooling from two or more capitula.

We used general linear models (regression and ANOVA) to determine whether measurements differed among F_1 families. The proportion variables, proportion self seed set and pollen viability, were arcsine square root transformed for these analyses. Calculations were done using Minitab (ver. 14).

Results

The F_2 generation segregated roughly into SI-PSC and SC classes in a 3:1 ratio, regardless of the specific criteria used to define the classes (fig. 3A). For example, if ≤ 0.10 seed set and ≥ 0.70 are used to define SI-PSC and SC classes, respectively, then the ratio is 3.16:1, whereas using ≤ 0.30 and ≥ 0.50 results in a 2.94:1 ratio. The segregation is bimodal, with most individuals having very low or relatively high self seed set. A substantial proportion of the variation in self seed set may be due to the availability of viable self pollen (fig. 3B). The regression of self seed set onto pollen viability is highly significant ($F_{1,122} = 24.3$, $P < 0.001$). However, the abundance of plants with high pollen viability yet low self seed set indicates that pollen viability does not fully account for differences in self seed set (fig. 3B). Pollen viability differed significantly among families within the F_2 generation ($F_{8,119} = 7.83$, $P < 0.001$), but self seed set did not ($F_{8,115} = 1.86$, $P = 0.07$). All five of the morphological traits differed significantly among families.

The summary statistics for measurements of floral traits in F_2 individuals are given in table 1, and their frequency distributions are provided in figures A1–A5 (available in the online edition of the *International Journal of Plant Sciences*). Each trait displays continuous variation, suggesting that multiple—and perhaps many—loci contribute to differences between the




Fig. 3 A, Histogram showing number of individuals at each proportion of self seed set in the F_2 generation. B, Scatter plot of proportion pollen viability versus proportion viable seed set for F_2 generation.

parental genotypes. As expected, mean floral dimensions (i.e., florets per capitulum, capitulum diameter, and floret, anther, and stigma lengths) are positively correlated (table 2). Pollen:ovule ratio is significantly correlated with floret, anther, and stigma length but not with self seed set (table 2). Interestingly, the floral dimensions exhibit significantly negative correlations with self seed set and pollen viability. The relationship between mean anther length and self seed set is illustrated in figure 4. Anther length shows no relationship with self seed set for anthers less than $28 \mu\text{m}$. However, self seed set drops to nearly 0 for plants with anthers longer than $28 \mu\text{m}$. Similar but less extreme patterns are observed for other floral dimensions.

Discussion

The roughly 3:1 ratio for SI-PSC versus SC individuals in the F_2 generations suggests that the breeding system in *Tolpis*

Table 1**Summary Statistics for Floral Traits Measured in *Tolpis* F₂ Hybrids**

Variable	N	Mean	SD
No. florets/capitulum	129	88.41	18.94
Capitulum diameter (cm)	129	1.5155	.2920
Floret length (cm)	129	.9030	.1554
Anther length (μm)	129	26.627	3.123
Stigma length (μm)	129	5.6885	.8926
Proportion pollen viability	129	.7608	.2902
Proportion self seed set	124	.2447	.3469
Pollen : ovule ratio	128	1904.5 : 1	499.7

Note. N, number of plants.

is controlled by a major locus, as it is in the sporophytic SI (SSI) system of other Asteraceae such as *Senecio squalidus* (Hiscock and Tabah 2003), and that SI (at least with regard to the alleles present in the parents of the hybrids) is dominant to SC. The match to a 3 : 1 ratio is even closer if one excludes those plants with low pollen viability. Although not always explicitly considered in studies of seed set, low pollen viability can clearly reduce the level of self seed set even in fully SC individuals (Gottlieb 1973). The complexity of the SSI system in Asteraceae has been documented though not fully elucidated. A major factor conferring complexity is the possible dominance relationships among S-alleles with SSI (Brauner and Gottlieb 1987; Levin 1996; Hiscock and Tabah 2003; Busch and Schoen 2008; Brennan et al. 2011). Dominance among alleles may differ in pollen and stigma, resulting in different compatibility results for reciprocal crosses (Hiscock and Tabah 2003). Another source of complexity in elucidating the genetics of SSI is the occurrence of modifier loci unlinked to the S-locus (Good-Avila and Stephensen 2002; Brennan et al. 2011). These loci may alter dominance relationships among S-alleles or increase compatible matings not interpretable in the context of S-alleles in an SSI system (Hiscock 2000; Brennan et al. 2003, 2011). One or more of these factors as well as others may have influenced results in this study. For example, among the F₂ plants in the SI-PSC category, only five had no viable seed, indicating that SI is leaky, with modifier loci a possible cause. Only two of the SC plants had 1.0 proportion self seed set, and in some instances, reduced self seed set of SC plants co-occurs with low pollen viability (fig. 3B). Resource limitation may also have been a cause of reduced seed set, although this is not likely because only vigorous hybrid plants were selected for study. While much remains to be determined about the genetics of

SI in *Tolpis*, present data indicate that a single major locus influences seed set, with alleles for SI dominant to those for SC but with other factors likely involved.

The loss of SI has been documented between sister or progenitor-derivative species (Ornduff 1966; Gottlieb 1973; Beck et al. 2006; Busch and Urban 2011; Slotte et al. 2012), between sister subspecies (Pettengill and Moeller 2011), among populations of a species (Busch 2005; Koelling et al. 2011), and among individuals within populations (Brauner and Gottlieb 1987; Tsukamoto et al. 1999; Mable et al. 2005). The breakdown of SI may be associated with one or more floral features collectively known as the selfing syndrome (Ornduff 1969), so that closely related SI-PSC and SC species differ by several of these floral characters (Ornduff 1966; Foxe et al. 2009; Slotte et al. 2012). Components of the selfing syndrome typically include fewer, smaller flowers and loss of floral rewards for pollinators, ostensibly because of reduced selection for features attracting pollinators (Ornduff 1969). Also, there may be selection for features promoting autonomous autogamy, such as anthers and stigmas in close proximity and synchrony of anther dehiscence and stigma receptivity (Ornduff 1969).

A major result of this study is that five floral traits that are part of the selfing syndrome are associated with viable self seed set (SC) in the F₂ individuals. This might be expected if plants with these traits have higher seed set because of more efficient autonomous autogamy. However, preliminary observations have not confirmed this hypothesis; selfed seed set in *Tolpis* does not differ whether capitula of a plant are unmanipulated or florets are manually pollinated with pollen from the same capitulum (D. J. Crawford, M. E. Mort, and A. Santos-Guerra, unpublished data). Thus, while more data are needed, there is no evidence at present that the selfing syndrome in *Tolpis* increases seed set by more efficient autonomous selfing. It may be that selection for pollinator attraction has precluded development of the selfing syndrome in some PSC plants, whereas outcrossed seed production is not an issue with SC plants. However, this does not explain the correlation of selfing morphological features and seed set in these F₂ plants, where the influence of selection on co-occurrence of traits has been removed. This correlation may instead be due to either genetic linkage or pleiotropy. The measured floral dimensions appear to be controlled by multiple loci (figs. A1–A5). This suggests that genetic linkage is unlikely, since it would mean that several loci (involved in floral morphology) would coincidentally have to be genetically proximate to each other and to the major locus responsible for SI/SC. Pleiotropy remains as a possibility, although we are aware of no precedents for an SI locus directly influencing morphology.

Table 2**Correlations (P Value) between Floral Traits in *Tolpis* F₂ Hybrids**

	Florets/capitulum	Capitulum diameter	Floret length	Anther length	Stigma length	Pollen : ovule ratio	Pollen viability
Capitulum diameter	.661 (.000)	...					
Floret length	.530 (.000)	.904 (.000)	...				
Anther length	.503 (.000)	.650 (.000)	.697 (.000)	...			
Stigma length	.229 (.009)	.393 (.000)	.405 (.000)	.542 (.000)	...		
Pollen : ovule ratio	.091 (.305)	.133 (.135)	.191 (.031)	.252 (.004)	.192 (.030)	...	
Pollen viability	-.287 (.001)	-.342 (.000)	-.359 (.000)	-.352 (.000)	-.379 (.000)	-.135 (.128)	...
Self seed set	-.224 (.012)	-.292 (.001)	-.376 (.000)	-.443 (.000)	-.257 (.015)	-.108 (.230)	.391 (.000)




Fig. 4 Proportion self seed set versus anther length in F_2 generation.

Pollen : ovule ratios have been shown to be good indicators of breeding system, with selfers having a lower pollen : ovule ratio than outcrossers (Cruden 1977). Differences are seen between pollen : ovule ratios in the SC *Tolpis coronopifolia* (400–900 : 1) and all of the SI-PSC species (2,500–3,800 : 1) of Canarian *Tolpis* (Crawford et al. 2008), though the pollen : ovule ratios for *T. coronopifolia* are not as low as the means given by Cruden (1977) for obligate or facultative autogamy. Pollen : ovule ratios in the F_2 individuals measured here are correlated with the lengths of florets, anthers, and stigmas (table 2). This may simply be due to the number of pollen grains being constrained by general floret size and especially by the length of anthers, that is, the associations being caused by functional constraints. Causation is not shown here, but reduction of floral size may be the mechanism behind lower pollen : ovule ratios in the SC *T. coronopifolia*; pollen : ovule ratios were not found to be correlated directly with self seed set in these F_2 individuals (table 2).

The low divergence among species of Canarian *Tolpis* for several molecular markers (Archibald et al. 2006; Crawford et al. 2006; Mort et al. 2010) suggests that SC and the selfing syndrome evolved recently and rapidly in *T. coronopifolia*. This is in agreement with other studies in which sequence data indicate that SI-SC species pairs have diverged recently, for example, with estimates of 20,000–50,000 yr for two species of *Capsella* (Brassicaceae; Foxe et al. 2009; Guo et al. 2009). Within members of Asteraceae, molecular, geological, and biogeographical data support recent divergence between SI and SC species of *Lasthenia* (Ornduff 1966; Chan et al. 2001) and *Stephanomeria* (Gottlieb 1973; Lee et al. 2002). The correlation between the floral traits and SC would aid selection on the entire suite of features and facilitate rapid divergence between *T. coronopifolia* and the SI-PSC species of *Tolpis* in the Canarian clade. Similarly, the correlation among floral morphological traits may speed divergence of this suite of characteristics. The latter correlation may be partly due to developmental constraints dictating that smaller flowers will lead to smaller floral parts. The shorter generation time for the annual *T. coronopifolia* could also accelerate divergence compared with perennials.

The factors driving evolution of SC in *T. coronopifolia* on the Canary Islands remain obscure, but there are several possible

reasons for loss of SI-PSC and evolution of the floral selfing syndrome. Leaky SI (PSC) is pervasive in species of *Tolpis* in the Canaries (Crawford et al. 2008, 2010). Evidence indicates that PSC can facilitate increased selfing with weakening of SI and the further increase of PSC to eventually produce nonfunctional S-alleles (reviewed in Levin 1996; Busch and Schoen 2008). Levin (1996, p. 326) stated after review of the literature, “I am unaware of any instance in which S_f [self-fertility] alleles increased in a species that was not already pseudo-self-fertile.” PSC varies quantitatively, and selection can raise or lower self seed set over generations (Levin 1996). Factors such as strength of S-alleles and modifier loci are likely responsible for the variation on which selection can act. By contrast, once an S-allele becomes nonfunctional, it is highly unlikely, if not impossible, that reversal to the SI-PSC condition will occur (Igc et al. 2008).

One factor that could increase SC is the inherent 50% transmission advantage of selfing over strictly outcrossing plants because selfing plants contribute two sets of gametes to each selfed offspring and can contribute to the next generation by both selfing and siring offspring with other plants (Fisher 1941; Jain 1976). Another factor that could select for increased selfing, both PSC and SC, is the limitation of compatible mates (Busch and Schoen 2008). This could include small populations with very low S-allele diversity and single colonizers founding new populations following dispersal. *Tolpis coronopifolia* occurs on the island of Tenerife in primarily open habitats of the xerophytic scrub zone (Jarvis 1980). This species is an annual and thus has only one season of reproductive opportunity, which might favor the evolution of SC. Unlike several SI-PSC perennial *Tolpis* species that occur on more than one island in the Canarian archipelago, *T. coronopifolia* has not dispersed from Tenerife and become established on any of the four other high islands. However, in the Canaries, frequent colonization events may be likely even within single islands. Disturbances such as landslides and volcanic activity have occurred throughout the history of the Canaries, including recent historical times (Carracedo and Day 2002, chap. 6). Human disturbance, such as development for tourism and road building, are also constantly creating open habitats such as those favored by *T. coronopifolia* (Caujapé-Castells et al. 2010). Species of *Tolpis* are found along roadsides over altitudinal gradients and vegetation types on the island of Tenerife (Arévalo et al. 2005).

Field studies have not detected natural hybrids between *T. coronopifolia* and other endemic species of *Tolpis* in the Canaries (A. Santos-Guerra, unpublished data). Given that experimental hybrids are readily synthesized in cultivation, one or more factors presumably restrict interspecific gene flow in the field. *Tolpis coronopifolia* is known to occur with other species of *Tolpis* on occasion, but it more often occurs in open habitats where other species are much less common (Jarvis 1980). The highly selfing breeding system of *T. coronopifolia* may also reduce gene flow with congeneric species, as has been suggested for other plants (Antonovics 1968; Fishman and Wyatt 1999; Wendt et al. 2002; Grossenbacher and Whittall 2011). Reduced pollen viability of some F_2 plants (fig. 3B) suggests some Dobzhansky-Muller incompatibility.

tibilities between the parental species (Fishman and Willis 2001), although their role restricting the occurrence of advanced generation or backcross hybrids remains to be elucidated.

In summary, present evidence indicates that *T. coronopifolia* evolved recently from SI or, more precisely, PSC ancestors on Tenerife in the Canary Islands. The correlation of SC with several floral traits likely facilitated the rapid origin of the *T. coronopifolia* phenotype. The most feasible hypothesis is that selection for selfing in the colonization of open areas drove the origin and evolution of *T. coronopifolia*.

Acknowledgments

This research was supported by the Department of Ecology and Evolutionary Biology, the Biodiversity Institute, and Graduate Research Funds at the University of Kansas. A Kansas National Science Foundation Experimental Program to Stimulate Competitive Research (EPSCoR) Ecological Genomics postdoctoral award was given to J. K. Archibald and D. J. Crawford. We thank Katie Sadler for her assistance in greenhouse cultivation of plants, Craig Freeman for advice on micromorphological studies, Clare Prohaska for assistance with greenhouse work, and Gilbert Ortiz for production of the figures.

Literature Cited

- Allard RW 1965 Genetic systems associated with colonizing ability in predominately self-pollinated species. Pages 49–76 in HG Baker, GL Stebbins, eds. The genetics of colonizing species. Academic Press, New York.
- Anderson GJ, G Bernardello, TF Stuessy, DJ Crawford 2001 Breeding system and pollination of selected plants endemic to Juan Fernández islands. *Am J Bot* 88:220–233.
- Antonovics J 1968 Evolution in closely adjacent plant populations. V. Evolution of self-fertility. *Heredity* 23:219–228.
- Archibald JK, DJ Crawford, A Santos-Guerra, ME Mort 2006 The utility of automated analysis of inter-simple sequence repeat (ISSR) loci for resolving relationships in the Canary Island species of *Tolpis* (Asteraceae). *Am J Bot* 93:1154–1162.
- Arévalo J, RD Delgado, R Otto, A Naranjo, M Salas, JM Fernández-Palacios 2005 Distribution of alien vs. native plant species in roadside communities along an altitudinal gradient in Tenerife and Gran Canaria (Canary Islands). *Perspect Plant Ecol Evol Syst* 7: 185–202.
- Bainbridge SJ, BG Baldwin 2010 Self-incompatibility, pollen limitation, and endangerment in the Hawaiian silversword alliance (Compositae) on Kaua'i. Botany Conference, Providence, RI.
- Baker HG 1955 Self-compatibility and establishment after “long-distance” dispersal. *Evolution* 9:347–349.
- Baldwin BG 2003 Natural history of the continental tarweeds and the Hawaiian silversword alliance (Asteraceae-Madiinae). Pages 1–16 in S Carlquist, BG Baldwin, GD Carr, eds. Tarweeds and silverswords: evolution of the Madiinae (Asteraceae). Missouri Botanical Garden Press, St. Louis.
- Barrett SCH 1996 The reproductive biology and genetics of island plants. *Philos Trans R Soc B* 351:725–733.
- 2002 The evolution of plant sexual diversity. *Nat Rev Genet* 3:274–284.
- Barrier M, BG Baldwin, RH Robichaux, MD Purugganan 1999 Interspecific hybrid ancestry of a plant adaptive radiation: allopolyploidy of the Hawaiian silversword alliance (Asteraceae) inferred from floral homeotic gene duplications. *Mol Phylogenet Evol* 16: 1105–1113.
- Beck JB, IA Al-Shehbaz, BA Schaal 2006 *Leavenworthia* (Brassicaceae) revisited: testing classic systematic and mating system hypotheses. *Syst Bot* 31:151–159.
- Bernardello G, GJ Anderson, TF Stuessy, DJ Crawford 2001 A survey of floral traits, breeding systems, floral visitors, and pollination systems of the angiosperms of the Juan Fernandez Islands (Chile). *Bot Rev (Lancaster)* 67:255–308.
- Brauner S, LD Gottlieb 1987 A self-compatible plant of *Stephanomeria exigua* subsp. *coronaria* (Asteraceae) and its relevance to the origin of its self-pollinating derivative *S. malheurensis*. *Syst Bot* 12: 29–304.
- Brennan AC, SA Harris, SJ Hiscock 2003 The population genetics of sporophytic self-incompatibility in *Senecio squalidus* L. (Asteraceae): avoidance of mating constraints imposed by low S-allele numbers. *Philos Trans R Soc B* 358:1047–1050.
- Brennan AC, DA Tabah, SA Harris, SJ Hiscock 2011 Sporophytic self-incompatibility in *Senecio squalidus* (Asteraceae): S allele dominance interactions and modifiers of cross-compatibility and selfing rates. *Heredity* 106:113–123.
- Busch JW 2005 Inbreeding depression in self-incompatible and self-compatible populations of *Leavenworthia alabamica*. *Heredity* 94: 159–165.
- Busch JW, DJ Schoen 2008 The evolution of self-incompatibility when mates are limiting. *Trends Plant Sci* 13:128–136.
- Busch JW, L Urban 2011 Insights from 50 years of studying the evolution of self-compatibility in *Leavenworthia* (Brassicaceae). *Evol Biol* 38:15–27.
- Carlquist S 1966 The biota of long-distance dispersal. IV. Genetic systems in the floras of oceanic islands. *Evolution* 20:433–455.
- 1974 Island biology. Columbia University Press, New York.
- Carr GD, EA Powell, DW Kyhos 1986 Self-incompatibility in the Hawaiian Madiinae (Compositae): an exception to Baker's rule. *Evolution* 40:430–434.
- Carracedo JC, S Day 2002 Canary Islands. *Classic Geology in Europe* 4. Terra, Hertfordshire.
- Caujapé-Castells J, A Tye, DJ Crawford, A Santos-Guerra, A Sakai, K Beaver, W Lobin, et al 2010 Conservation of oceanic island floras: present and future global challenges. *Perspect Plant Ecol Evol Syst* 12:107–129.
- Chan R, BG Baldwin, R Ornduff 2001 Goldfields revisited: a molecular phylogenetic perspective on the evolution of *Lasthenia* (Compositae: Heliantheae sensu lato). *Int J Plant Sci* 162:1347–1360.
- Charlesworth D 2006 Evolution of plant breeding systems. *Curr Biol* 16:R726–R735.
- Cowie RH, BS Holland 2006 Dispersal is fundamental to biogeography and the evolution of biodiversity on oceanic islands. *J Biogeogr* 33:193–198.
- Crawford DJ, GJ Anderson, G Bernardello 2011 The reproductive biology of island plants. Pages 11–36 in D Bramwell, J Caujapé-Castells, eds. The biology of island floras. Cambridge University Press, Cambridge.
- Crawford DJ, JK Archibald, JK Kelly, ME Mort, A Santos-Guerra 2010 Mixed mating in the “obligately outcrossing” *Tolpis* (Asteraceae) of the Canary Islands. *Plant Species Biol* 25:114–119.
- Crawford DJ, JK Archibald, ME Mort, A Santos-Guerra 2009a Pollen fertility of synthetic intra- and interspecific F₁ hybrids in Canary Island *Tolpis* (Asteraceae). *Plant Syst Evol* 279:103–113.
- Crawford DJ, JK Archibald, A Santos-Guerra, ME Mort 2006 Al-

- lozyme diversity within and divergence among species of *Tolpis* (Asteraceae-Lactuceae) in the Canary Islands: systematic, evolutionary, and biogeographical implications. *Am J Bot* 93:656–664.
- Crawford DJ, JK Archibald, D Stormier, ME Mort, JK Kelly, A Santos-Guerra 2008 A test of Baker's law: the radiation of *Tolpis* (Asteraceae) in the Canary Islands. *Int J Plant Sci* 169:782–791.
- Crawford DJ, TK Lowrey, GJ Anderson, G Bernardello, A Santos-Guerra, TF Stuessy 2009b Genetic diversity in the colonizing ancestors of Asteraceae endemic to oceanic islands: Baker's law and polyploidy. Pages 139–151 in VA Funk, A Susanna, TF Stuessy, R Bayer, eds. Systematics, evolution, and biogeography of the Compositae. International Association of Plant Taxonomy, Vienna.
- Cruden RW 1977 Pollen-ovule ratios: a conservative indicator of breeding systems in flowering plants. *Evolution* 31:32–46.
- Ehrendorfer E 1979 Reproductive biology of island plants. Pages 293–306 in D Bramwell, ed. Plants and islands. Academic Press, New York.
- Fisher RA 1941 Average excess and average effect on a gene substitution. *Ann Eugen* 11:53–63.
- Fishman L, JH Willis 2001 Evidence for Dobzhansky-Muller incompatibilities contributing to the sterility of hybrids between *Mimulus guttatus* and *M. nasutus*. *Evolution* 55:1932–1942.
- Fishman L, RA Wyatt 1999 Pollinator-mediated competition, reproductive character displacement, and the evolution of selfing in *Arenaria* (Caryophyllaceae). *Evolution* 53:1723–1733.
- Foxe JP, T Slotte, EA Stahl, B Neuffer, H Hurka, SI Wright. 2009 Recent speciation associated with the evolution of selfing in *Capsella*. *Proc Natl Acad Sci USA* 106:5241–5245.
- Foxe JP, M Stift, A Tedder, A Haudry, SI Wright, BK Mable 2010 Reconstructing origins of loss of self-incompatibility and selfing in North American *Arabidopsis lyrata*: a population genetic context. *Evolution* 64:3495–3510.
- Good-Aliva SV, AG Stephensen 2002 The inheritance of modifiers conferring self-fertility in the partially self-incompatible perennial, *Campanula rapunculoides* L. (Campanulaceae). *Evolution* 56:263–272.
- Goodwillie C 1999 Multiple origins of self-compatibility in *Linanthus* section *Leptosiphon* (Polemoniaceae): phylogenetic evidence from internal-transcribed-spacer sequence data. *Evolution* 53:1387–1395.
- Goodwillie C, S Kalisz, CG Eckert 2005 The evolutionary enigma of mixed mating systems in plants: occurrence, theoretical explanations, and empirical evidence. *Annu Rev Ecol Syst* 36:47–79.
- Gottlieb LD 1973 Genetic differentiation, sympatric speciation, and origin of a diploid species of *Stephanomeria*. *Am J Bot* 60:545–553.
- Grossenbacher DL, JB Whittall 2011 Increased floral divergence in sympatric monkeyflowers. *Evolution* 65:2712–2718.
- Guo YL, JS Bechsgaard, T Slotte, B Neuffer, M Lascoux, D Weigel, MH Schierup 2009 Recent speciation of *Capsella rubella* from *Capsella grandiflora*, associated with loss of self-incompatibility and an extreme bottleneck. *Proc Natl Acad Sci USA* 106:5246–5251.
- Hiscock SJ 2000 Genetic control of self-incompatibility in *Senecio squalidus* (Asteraceae): a successful colonizing species. *Heredity* 85:10–19.
- Hiscock SJ, DA Tabah 2003 The different mechanisms of sporophytic self-incompatibility. *Philos Trans R Soc B* 358:1037–1045.
- Igic B, R Lande, JR Kohn 2008 Loss of self-incompatibility and its evolutionary consequences. *Int J Plant Sci* 169:93–104.
- Jain SK 1976 Evolution of inbreeding in plants. *Annu Rev Ecol Syst* 7:469–495.
- Jarvis CE 1980 Systematic studies in the genus *Tolpis* Adanson. PhD diss. University of Reading.
- Kalisz S, DW Volger, KM Hanley 2004 Context-dependent autonomous self-fertilization yields reproductive assurance and mixed mating. *Nature* 430:884–887.
- Kearns CA, DW Inouye 1993 Techniques for pollination biologists. University Press of Colorado, Niwot.
- Koelling V, JL Hamraick, R Mauricio 2011 Genetic diversity and structure in two species of *Leavenworthia* with self-incompatible and self-compatible populations. *Heredity* 106:310–318.
- Lee J, BG Baldwin, LD Gottlieb 2002 Phylogeny of *Stephanomeria* and related genera (Compositae-Lactuceae) based on analysis of 18S-26S nuclear rDNA ITS and ETS sequences. *Am J Bot* 89:160–168.
- Levin DA 1978 The origin of isolating mechanisms in flowering plants. *Evol Biol* 11:85–317.
- 1996 The evolutionary significance of pseudo-self-fertility. *Am Nat* 148:321–332.
- 2000 The origin, expansion, and demise of plant species. Oxford University Press, New York.
- Mable BK, AV Robertson, S Dart, C Di Bernardo, L Witham 2005 Breakdown of self-incompatibility in the perennial *Arabidopsis lyrata* (Brassicaceae) and its genetic consequences. *Evolution* 59:1437–1448.
- McMullen CK 1987 Breeding systems of selected Galápagos Islands angiosperms. *Am J Bot* 74:1694–1705.
- 1990 Reproductive biology of Galápagos Islands angiosperms. Pages 35–45 in JE Lawesson, O Hamann, G Rogers, G Reck, H Ochoa, eds. Botanical research and management in the Galápagos. Missouri Botanical Garden Press, St. Louis.
- Mort ME, DJ Crawford, JK Archibald, TR O'Leary, A Santos-Guerra 2010 Plant DNA barcoding: a test using Macaronesian taxa of *Tolpis*. *Taxon* 59:581–587.
- Nielsen LR, M Philipp, H Adersen, HR Siegismund 2000 Breeding system of *Scalesia divisa* Andersson, an endemic Asteraceae from the Galápagos Islands. *Det Norske Videnskaps-Akademi I. Matematisk-Naturvidenskabelige Klasse, Shriftr, Ny Serie* 39:127–138.
- Nielsen LR, HS Siegismund, M Philipp 2003 Partial self-incompatibility in the polyploid endemic species *Scalesia affinis* (Asteraceae) from the Galápagos: remnants of a self-incompatibility system? *Bot J Linn Soc* 142:93–101.
- Ornduff R 1966 A biosystematic study of the goldfield genus *Lasthenia*. *Univ Calif Publ Bot* 40:1–92.
- 1969 Reproductive biology in relation to systematics. *Taxon* 18:121–133.
- Pannell JR, SCH Barrett 1998 Baker's law revisited: reproductive assurance and the metapopulation. *Evolution* 52:657–668.
- Pettengill JB, DA Moeller 2011 Tempo and mode of mating system evolution between insipient *Clarkia* species. *Evolution* 66:1210–1225.
- Slotte T, KM Hazzouri, D Stern, P. Andolfatto, SI Wright 2012 Genetic architecture and adaptive significance of the selfing syndrome in *Capsella*. *Evolution* 66:1360–1374.
- Stebbins GL 1957 Self-fertilization and population variability in higher plants. *Am Nat* 91:337–354.
- Sun M, FR Ganders 1988 Mixed mating systems in Hawaiian *Bidens* (Asteraceae). *Evolution* 42:516–527.
- Tsukamoto T, T Ando, H Kokubun, H Watanabe, M Masada, X Zhu, E Marchesi, T Kao 1999 Breakdown of self-incompatibility in a natural population of *Petunia axillaris* (Solanaceae) in Uruguay containing both self-incompatible and self-compatible plants. *Sex Plant Reprod* 12:6–13.
- Webb CJ, D Kelly 1993 The reproductive biology of the New Zealand flora. *Trends Ecol Evol* 8:442–447.
- Wendt TM, BF Canela, DE Klein, RI Rios 2002 Selfing facilitates reproductive isolation among three sympatric species of *Pitcairnia* (Bromeliaceae). *Plant Syst Evol* 232:201–212.
- Yang JY, SA Hodges 2010 Early inbreeding depression selects for high outcrossing rates in *Aquilegia formosa* and *Aquilegia pubescens*. *Int J Plant Sci* 171:860–871.