

Full Length Research Paper

Yaoundé-like virus in resident wild bird, Ghana

Williams, Richard A. J.^{1,2*}, Vázquez, Ana³, Asante, Ivy⁴, Bonney, Kofi⁴, Odoom, Shirley⁴, Pupilampu, Naiki⁴, Ampofo, William⁴, Sánchez-Seco, María Paz³, Tenorio, Antonio³ and Peterson, A. Townsend¹

¹University of Kansas Biodiversity Institute, 1345 Jayhawk Blvd, Lawrence, Kansas, USA.

²Departamento de Zoología y Antropología Física, Complutense Universidad, Madrid 28040, Spain.

³Centro Nacional de Microbiología, Institute of Health Carlos III, Majadahonda, Spain.

⁴Noguchi Memorial Institute for Medical Research, Accra, Ghana.

Accepted 19 December, 2011

Tissue and swab samples from 551 wild birds collected in Ghana (October-November 2007) were assayed for alphaviruses, flaviviruses, and influenza A viruses using polymerase chain (PCR) techniques. One pool sample tested positive for *Flavivirus* RNA; further testing revealed that the amplified sequence was Yaoundé virus (YAOV), or closely related to it. YAOV is an apparently rare *Flavivirus* closely related to medically important human pathogens Japanese Encephalitis virus and West Nile virus. It is known only from West Africa. This is the first detection from Ghana, and only the second detection from a bird. Samples were negative for alphaviruses and Influenza A virus.

Key words: Birds, Flavivirus, Ghana, West Nile virus, Yaoundé virus.

INTRODUCTION

Recent range expansion of bird-borne viruses, particularly highly pathogenic avian influenza H5N1 (HPAI-H5N1), and West Nile virus (WN), have caused significant mortality in humans, wildlife and livestock (Daszak et al., 2000). Both viruses, and related taxa, are associated with wild birds, and spill over from wild bird reservoirs to infect humans (Webster et al., 1992; Komar, 2003). Although, these bird-borne pathogen shifts have attracted considerable research and public health attention, the details of bird-virus host associations in many cases remain poorly known.

Host-virus dynamics may vary according to viral and avian species, for instance WN amplifies principally among passerine songbirds (Ezenwa, 2006; Komar et al., 2003), whereas JE amplifies principally among Ciconiiformes (herons and allies; Buescher et al., 1959; Endy et al., 2002; Jamgaonkar, 2003), and so forth. Certainly experimental infection of avian hosts

demonstrates that both susceptibility to infection and the level and duration of viremia vary by species for WN (Komar et al., 2003) and St. Louis Encephalitis virus (McLean et al., 1985). Similarly, Anseriformes (ducks and geese) and Charadriiformes (shorebirds, gulls, and terns) are considered the primary reservoirs for avian influenza (AI) strains, notwithstanding the detection of more broadly across class Aves (Hesterberg et al., 2009).

No data are available on host range, prevalence, or habitat type for Flaviviruses or Influenza A viruses in wild birds in Ghana. Indeed, very little information is available for West Africa. Several Flaviviruses, notably WN, are present in West Africa (MacKenzie, 2002), and likely present in Ghana. A recent human survey found high WN seroprevalences, 30% (Wang, 2009), suggesting transmission in amplifying (avian) hosts. Investigation of AI strains in wild birds in sub-Saharan Africa has thus far concentrated on water birds, and several AI strains have been detected in West Africa (Gaidet et al., 2007), including HPAI-H5N1 (Enserink, 2006), which has been detected throughout West Africa, including Ghana (Williams et al., 2008).

Although, AI strains and Flaviviruses are clearly

*Corresponding author. E-mail: ricw.ku@gmail.com. Tel: +34 91 854 0360. Fax: +1 785 864 5335.

versatile as regards their range of possible hosts, some viruses appear to demonstrate some degree of non-random host association. These associations nevertheless remain untested and uncharacterized both over most virus distributions, and across most of host diversity.

As a result, in October and November 2007, we conducted broad-scale virus surveillance among land birds, in two distinct habitat types, savannah-woodland and rain forest, to assess the distribution of avian Influenza A viruses and Flaviviruses, using PCR techniques. In addition, we tested all samples for the presence of another bird-associated group, Alphaviruses. To the best of our knowledge, this survey is the first broad-scale study of these virus groups in Ghanaian land birds.

MATERIALS AND METHODS

We sampled and tested 551 free ranging birds and 12 backyard chickens for *Alphavirus*, *Flavivirus*, and influenza A virus infections using PCR techniques. Sampling was conducted in Ghana, at Ankasa Conservation Area ($N = 305$ individuals, 70 species), Western Region, and Gbele Resource Reserve ($N = 259$, 73 species), Upper West Region, in October–November 2007. Birds were obtained by mist netting and selective harvesting with shotguns; all birds were apparently healthy at the time of collection. Buccal and cloacal swabs were collected for up to 30 individuals per species, and mixed tissue samples (brain, gonad, intestine, kidney and lung) for up to 10 individuals per species. Samples were frozen immediately in liquid nitrogen, and voucher specimens were prepared for all samples and deposited at the University of Kansas Biodiversity Institute. 95.5% of individuals sampled were resident, 2.9% were Palearctic migrants, and the remaining 1.6% were species with populations that include internal African migrants.

Viral RNA was extracted from homogenized tissue pools using RNEasy minikit (QIAGEN, Valencia, CA, USA), and from swab pools prepared in VTM (containing 2.5% veal infusion broth, 0.5% BSA, 100 $\mu\text{g}/\text{ml}$ gentamicin sulfate and 2 $\mu\text{g}/\text{ml}$ Fungizone) using QIAamp Viral RNA Mini Kit (QIAGEN). Tissues were homogenized using TissueRuptor (QIAGEN). Molecular RNA detection techniques were used to detect genomic material from alphaviruses (Sánchez-Seco et al., 2001), flaviviruses (Pierre et al., 1994; Sánchez-Seco et al., 2005) and influenza viruses (CDC, 2009; Spackman et al., 2002). A flavivirus was detected from one pool of tissues from seven birds using the nested RT-PCR technique, although other tests resulted negative. The 100 nt amplification product was detected by electrophoresis, and purified using QIAquick PCR Purification Kit (QIAGEN). Sequencing reactions were performed with ABI Prism BigDye Terminator Cycle Sequencing v.3.1 Ready Reaction (Applied Biosystems, Foster City, CA, USA), and analyzed using an ABI PRISM model 3730 automated sequencer (Applied Biosystems). Assembly of consensus sequences and translation into amino acid sequences was performed with EditSeq (DNASTAR Inc., Madison, WI, USA). Phylogenetic analyses were developed relative to 13 *Flavivirus* sequences; all aligned using ClustalX (<http://www.clustal.org/>). Phylogenetic trees were constructed by Bayesian analysis, using MrBayes v. 3.1.2 (<http://mrbayes.csit.fsu.edu/>), with 10,000,000 cycles for the Markov chain Monte Carlo algorithm, and specifying TBEV (GenBank Acc No: NC_001672) as outgroup. Bayesian posterior probabilities were calculated from the consensus of 10,000 trees.

RESULTS AND DISCUSSION

One tissue pool of seven resident wild birds captured among the two sites was positive for *Flavivirus* NS5 sequence. The positive pool contained one *Alethe diademata*, two individuals of *Andropadus virens*, one *Oriolus brachyrhynchus*, and one *Tauraco macrorhynchus* from Ankasa, and one each of *Fringilla bicalcaratus* and *Petronia dentata* from Gbele. Attempts to determine viral host identity by testing samples from seven individuals from the *Flavivirus* positive pool individually, using the generic RT-nested PCR (Sánchez-Seco, 2005), were negative, presumably hampered by multiple failures of the cold chain for the diagnostic samples. Triturated tissue (and associated swab storage buffer) from individuals in the positive pool were inoculated into C6/36 (*Aedes albopictus* cells) and Vero (African green monkey cells) monolayer cells; no cytopathic effects were detected after three blind passages. Supernatant samples drawn from all passages of cell cultures were negative using the generic RT-nested PCR (Sánchez-Seco, 2005). Further attempts to amplify a more informative sequence using nine generic RT-nested PCR protocols (Kuno et al., 1998, and ISCI designed primers) were unsuccessful.

The putative YAOV sequence (GenBank accession number HQ290163) showed 92% sequence identity with a published YAOV strain (EU074036), with 8 nt differences (Figure 1), none producing amino acid replacements. Phylogenetic analyses (data not shown) confirmed the close relation of our sequence and YAOV (93% posterior probability). The virus we detected is thus YAOV, or one that is quite closely related.

Yaoundé virus (YAOV) is a mosquito-borne *Flavivirus* within the *Japanese encephalitis virus* (JEV) serocomplex which includes human pathogens such as JEV and *West Nile virus* (WNV; Kuno et al., 1998). YAOV has been isolated from one bird (*Bycanistes sharpii*), 2 mammal species (*Praomys* sp. and *Cavia porcellus*), and 10 mosquito species from four sub-Saharan African nations (Cameroon, Central African Republic, Congo, and Senegal; Adam and Digoutte, 2005). The virus was first isolated from a *Culex nebulosus* pool collected from degraded semi-deciduous forest near Nkolbisson, Cameroon, in 1968 (Digoutte, 1992). Although, YAOV has not been detected in humans (Adam and Digoutte, 2005), many other JEV group taxa have been.

To our knowledge, an active JEV group infection has not previously been detected directly in Ghana (that is, through the detection of viral RNA, viral culture, etc.). However, WNV presence has been inferred from a human serologic study that found an IgG seroprevalence of 27.9% in adults, and IgM seroprevalence of 2.4% in children <6 years old (Wang et al., 2009), the latter result indicates recent infection, presumably within Ghana. Given the close serologic relationships among JEV serocomplex viruses, it is possible that Yaoundé virus

Strain	0'	50'
YAOV: EU074036	AAAACGTGAG AAGAAGCCAG GGGAAATTCGG AAAGGCAAAA GGAAGCCGAG	
Ave-Ghana: HQ290163	-----A-----	-----G-
JEV: M18370	---A-A--- ---T- -A-G--T-- ---A--T--- ---A-G-	
WNV: DQ211652	---GA-A--- --A--A--C- -A--G----- ---C--G ---A---	
	51'	100'
YAOV: EU074036	CCATCTGGTA CATGTGGCTC GGAGCCCGAT TCTTGAGT TGAAGCCCTT	
Ave-Ghana: HQ290163	----T-----	---G-----
JEV: M18370	----T----T -----T -----A-G- ATC-A----- -----TT-G	
WNV: DQ211652	----T----T -----T--C- -TC----- C--G--T--G	

Figure 1. Nucleotide level comparison of sequenced fragment and *Yaoundé virus* (YAOV) sequences available on GenBank (catalog numbers provided). Comparisons with published sequences of known flaviviruses were performed by searches with the FASTA program in the EMBL database (www.ebi.ac.uk/embl) to identify the detected agent and study the level of homology. Here we show a nucleotide level comparison of sequenced fragment and *Yaoundé virus* (YAOV) available on GenBank (catalog numbers provided). Sequences from *Japanese Encephalitis virus* (JEV; 22 nt differences) and *West Nile virus* lineage 1 (WNV1; 21 nt differences), taxa closely related to YAOV are included to illustrate the close relationship between our sequence and YAOV. Dash indicates coincident nucleotide representation.

could be the source of WNV-reactive antibodies.

In conclusion, we detected one Flavivirus in a survey of asymptomatic wild birds in Ghana. The sequence showed a close relationship with YAOV. No alphaviruses or Influenza A viruses were recovered, suggesting that circulation of these virus groups is low, or zero, at these sites at the time of sampling.

ACKNOWLEDGEMENTS

We thank field companions including E. Bonaccorso, M. Robbins, and M. Thompson; laboratory companions, including A. Negrodo and N. Reyes; Dr. F. Pozo and I. Casas for introduction to ISCIII for their collaboration; and L. Benitez for introduction to ISCIII. This study was supported by the US National Biological Information Infrastructure, the GAINS program of the Wildlife Conservation Society, and the Centers for Disease Control and Prevention, grants FIS PI07/1308 of the Red de Investigacion de Centros de Enfermedades Tropicales RD06/0021, and the agreement signed between the Institute of Health Carlos III and the Spanish Ministry of Health and Social Policy for the surveillance of imported viral hemorrhagic fevers. Influenza work was supported by grant GR09/0040 (MPY-1440/09) ISCIII. RW is supported by grant CGL2010-15734/BOS, Ministerio de Ciencia e Innovación, Spain.

REFERENCES

Adam F, Digoutte JP (2005). Virus d'Afrique (base de données).

- <http://www.pasteur.fr/recherche/banques/CRORA>.
- Buescher EL, Scherer WF, McClure HE, Meyer JT, Rosenberg MZ, Yoshi UM, Okada Y (1959). Ecological studies of Japanese Encephalitis in Japan. IV. Avian infection. *Am. J. Trop. Med. Hyg.*, 8: 678-688.
- Centers for Disease Control and Prevention (CDC) (2009). CDC protocol of real-time RTPCR for influenza A(H1N1); Revision 2. www.who.int/csr/resources/publications/swineflu/realtimeptcr/en/ind ex.html.
- Daszak P, Cunningham AA, Hyatt AD (2000). Wildlife ecology - Emerging infectious diseases of wildlife - Threats to biodiversity and human health. *Science*, 287: 443-449.
- Digoutte JP (1992). Rapport sur le fonctionnement technique de l'Institut Pasteur de Dakar. Année 1991, Institut Pasteur de Dakar, Dakar.
- Endy TP, Nisalak A (2002). Japanese Encephalitis virus: ecology and epidemiology. In: Mackenzie JS, Barrett ADT, Deubel V (eds) *Japanese Encephalitis and West Nile Viruses*. Springer-Verlag, Berlin, pp. 11-48.
- Enserink M (2006). Avian influenza. H5N1 moves into Africa, European Union, deepening global crisis. *Science*, 311: 932.
- Ezenwa VO, Godsey MS, King RJ, Guptill SC (2006). Avian diversity and West Nile virus: testing associations between biodiversity and infectious disease risk. *Proc. Roy. Soc., B* 273:109-117.
- Gaidet N, Dodman T, Caron A, Balança G, Desvaux S, Goutard F, Cattoli G, Lamarque F, Hagemeijer W, Monicat F (2007). Avian influenza viruses in water birds, Africa. *Emerg. Infect. Dis.*, 13: 626-629.
- Hesterberg U, Harris K, Stroud D, Guberti V, Busani L, Pittman M, Piazza V, Cook A, Brown I (2009). Avian influenza surveillance in wild birds in the European Union in 2006. *Influenza Other Respir. Viruses*, 3: 1-14.
- Jamgaonkar AV, Yergolkar PN, Geevarghese G, Joshi GD, Joshi MV, Mishra AC (2003). Serological evidence for Japanese Encephalitis virus and West Nile virus infections in water frequenting and terrestrial wild birds in Kolar District, Karnataka State, India. A retrospective study. *Acta Virol.*, 47: 185-188.
- Komar N (2003). West Nile virus: epidemiology and ecology in North America. *Adv. Virus Res.*, 61: 185-234.
- Komar N, Langevin S, Hinten S, Nemeth N, Edwards E, Hettler D, Davis B, Bowen R, Bunning M (2003). Experimental infection of North American birds with the New York 1999 strain of West Nile virus.

- Emerg. Infect. Dis., 9: 311-322.
- Kuno G, Chang GJ, Tsuchiya KR, Karabatsos N, Cropp CB (1998). Phylogeny of the Genus *Flavivirus*. *J. Virol.*, 72: 73-83.
- Mackenzie JS, Barrett ADT, Deubel V (2002). The Japanese Encephalitis serological group of flaviviruses: a brief introduction to the group. In: Mackenzie JS, Barrett ADT, Deubel V (eds) *Japanese Encephalitis and West Nile Viruses*. Springer-Verlag, Berlin, pp. 1-10.
- McLean RG, Francly DB, Campos EG (1985). Experimental studies of St. Louis Encephalitis virus in vertebrates. *J. Wildlife Dis.*, 21: 85-93.
- nested-RT-PCR followed by sequencing for detection and identification of members of the *Alphavirus* genus. *J. Virol. Methods*, 95: 153-161.
- Pierre V, Drouet MT, Deubel (1994). Identification of mosquito-borne *Flavivirus*.
- Sánchez-Seco MP, Rosario D, Domingo C, Hernández L, Valdés K, Guzmán MG, Tenorio A (2005). Generic RT-nested-PCR for detection of flaviviruses using degenerated primers and internal control followed by sequencing for specific identification. *J. Virol. Methods*, 126: 101-109.
- Sánchez-Seco MP, Rosario D, Quiroz E, Guzmán G, Tenorio A (2001). A generic sequences using universal primers and reverse transcriptase-polymerase chain-reaction. *Res. Virol.*, 145: 93-104.
- Spackman E, Senne DA, Myers TJ, Bulaga LL, Garber LP, Perdue ML, Lohman K, Daum LT, Suarez DL (2002). Development of a real-time reverse transcriptase PCR assay for type A influenza virus and the avian H5 and H7 hemagglutinin subtypes. *J. Clin. Microbiol.*, 40: 3256-3256.
- Wang WJ, Sarkodie F, Danso K, Addo-Yobo E, Owusu-Ofori S, Allain JP, Li CY (2009). Seroprevalence of *West Nile virus* in Ghana. *Viral Immunol.*, 22: 17-22.
- Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y (1992). Evolution and ecology of influenza-A viruses. *Microbiol. Rev.*, 56: 152-179.
- Williams RAJ, Fasina FO, Peterson AT (2008). Predictable ecology and geography of avian influenza (H5N1) transmission in Nigeria and West Africa. *Trans. Roy. Soc. Trop. Med. Hyg.*, 102: 471-479.