

Measurements of the Mass, Total Width, and Two-Photon Partial Width of the η_c Meson

G. Brandenburg,¹ A. Ershov,¹ Y. S. Gao,¹ D. Y.-J. Kim,¹ R. Wilson,¹ T. E. Browder,² Y. Li,² J. L. Rodriguez,² H. Yamamoto,² T. Bergfeld,³ B. I. Eisenstein,³ J. Ernst,³ G. E. Gladding,³ G. D. Gollin,³ R. M. Hans,³ E. Johnson,³ I. Karliner,³ M. A. Marsh,³ M. Palmer,³ C. Plager,³ C. Sedlack,³ M. Selen,³ J. J. Thaler,³ J. Williams,³ K. W. Edwards,⁴ R. Janicek,⁵ P. M. Patel,⁵ A. J. Sadoff,⁶ R. Ammar,⁷ A. Bean,⁷ D. Besson,⁷ R. Davis,⁷ N. Kwak,⁷ X. Zhao,⁷ S. Anderson,⁸ V. V. Frolov,⁸ Y. Kubota,⁸ S. J. Lee,⁸ R. Mahapatra,⁸ J. J. O'Neill,⁸ R. Poling,⁸ T. Riehle,⁸ A. Smith,⁸ C. J. Stepaniak,⁸ J. Urheim,⁸ S. Ahmed,⁹ M. S. Alam,⁹ S. B. Athar,⁹ L. Jian,⁹ L. Ling,⁹ M. Saleem,⁹ S. Timm,⁹ F. Wappler,⁹ A. Anastassov,¹⁰ J. E. Duboscq,¹⁰ E. Eckhart,¹⁰ K. K. Gan,¹⁰ C. Gwon,¹⁰ T. Hart,¹⁰ K. Honscheid,¹⁰ D. Hufnagel,¹⁰ H. Kagan,¹⁰ R. Kass,¹⁰ T. K. Pedlar,¹⁰ H. Schwarthoff,¹⁰ J. B. Thayer,¹⁰ E. von Toerne,¹⁰ M. M. Zoeller,¹⁰ S. J. Richichi,¹¹ H. Severini,¹¹ P. Skubic,¹¹ A. Undrus,¹¹ S. Chen,¹² J. Fast,¹² J. W. Hinson,¹² J. Lee,¹² D. H. Miller,¹² E. I. Shibata,¹² I. P. J. Shipsey,¹² V. Pavlunin,¹² D. Cronin-Hennessy,¹³ A. L. Lyon,¹³ E. H. Thorndike,¹³ C. P. Jessop,¹⁴ H. Marsiske,¹⁴ M. L. Perl,¹⁴ V. Savinov,¹⁴ D. Ugolini,¹⁴ X. Zhou,¹⁴ T. E. Coan,¹⁵ V. Fadeyev,¹⁵ Y. Maravin,¹⁵ I. Narsky,¹⁵ R. Stroynowski,¹⁵ J. Ye,¹⁵ T. Wlodek,¹⁵ M. Artuso,¹⁶ R. Ayad,¹⁶ C. Boulahouache,¹⁶ K. Bukin,¹⁶ E. Dambasuren,¹⁶ S. Karamov,¹⁶ G. Majumder,¹⁶ G. C. Moneti,¹⁶ R. Mountain,¹⁶ S. Schuh,¹⁶ T. Skwarnicki,¹⁶ S. Stone,¹⁶ G. Viehhauser,¹⁶ J. C. Wang,¹⁶ A. Wolf,¹⁶ J. Wu,¹⁶ S. Kopp,¹⁷ A. H. Mahmood,¹⁸ S. E. Csorna,¹⁹ I. Danko,¹⁹ K. W. McLean,¹⁹ Sz. Márka,¹⁹ Z. Xu,¹⁹ R. Godang,²⁰ K. Kinoshita,^{20,*} I. C. Lai,²⁰ S. Schrenk,²⁰ G. Bonvicini,²¹ D. Cinabro,²¹ S. McGee,²¹ L. P. Perera,²¹ G. J. Zhou,²¹ E. Lipeles,²² S. P. Pappas,²² M. Schmidler,²² A. Shapiro,²² W. M. Sun,²² A. J. Weinstein,²² F. Würthwein,^{22,†} D. E. Jaffe,²³ G. Masek,²³ H. P. Paar,²³ E. M. Potter,²³ S. Prell,²³ V. Sharma,²³ D. M. Asner,²⁴ A. Eppich,²⁴ T. S. Hill,²⁴ R. J. Morrison,²⁴ R. A. Briere,²⁵ T. Ferguson,²⁵ H. Vogel,²⁵ B. H. Behrens,²⁶ W. T. Ford,²⁶ A. Gritsan,²⁶ J. Roy,²⁶ J. G. Smith,²⁶ J. P. Alexander,²⁷ R. Baker,²⁷ C. Bebek,²⁷ B. E. Berger,²⁷ K. Berkelman,²⁷ F. Blanc,²⁷ V. Boisvert,²⁷ D. G. Cassel,²⁷ M. Dickson,²⁷ P. S. Drell,²⁷ K. M. Ecklund,²⁷ R. Ehrlich,²⁷ A. D. Foland,²⁷ P. Gaidarev,²⁷ R. S. Galik,²⁷ L. Gibbons,²⁷ B. Gittelman,²⁷ S. W. Gray,²⁷ D. L. Hartill,²⁷ B. K. Heltsley,²⁷ P. I. Hopman,²⁷ C. D. Jones,²⁷ D. L. Kreinick,²⁷ M. Lohner,²⁷ A. Magerkurth,²⁷ T. O. Meyer,²⁷ N. B. Mistry,²⁷ E. Nordberg,²⁷ J. R. Patterson,²⁷ D. Peterson,²⁷ D. Riley,²⁷ J. G. Thayer,²⁷ P. G. Thies,²⁷ D. Urner,²⁷ B. Valant-Spaight,²⁷ A. Warburton,²⁷ P. Avery,²⁸ C. Prescott,²⁸ A. I. Rubiera,²⁸ J. Yelton,²⁸ and J. Zheng²⁸

(CLEO Collaboration)

¹Harvard University, Cambridge, Massachusetts 02138

²University of Hawaii at Manoa, Honolulu, Hawaii 96822

³University of Illinois, Urbana-Champaign, Illinois 61801

⁴Carleton University, Ottawa, Ontario, Canada K1S 5B6
and the Institute of Particle Physics, Canada

⁵McGill University, Montréal, Québec, Canada H3A 2T8
and the Institute of Particle Physics, Canada

⁶Ithaca College, Ithaca, New York 14850

⁷University of Kansas, Lawrence, Kansas 66045

⁸University of Minnesota, Minneapolis, Minnesota 55455

⁹State University of New York at Albany, Albany, New York 12222

¹⁰The Ohio State University, Columbus, Ohio 43210

¹¹University of Oklahoma, Norman, Oklahoma 73019

¹²Purdue University, West Lafayette, Indiana 47907

¹³University of Rochester, Rochester, New York 14627

¹⁴Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309

¹⁵Southern Methodist University, Dallas, Texas 75275

¹⁶Syracuse University, Syracuse, New York 13244

¹⁷University of Texas, Austin, Texas 78712

¹⁸University of Texas-Pan American, Edinburg, Texas 78539

¹⁹Vanderbilt University, Nashville, Tennessee 37235

²⁰Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

²¹Wayne State University, Detroit, Michigan 48202

²²California Institute of Technology, Pasadena, California 91125

²³University of California, San Diego, La Jolla, California 92093

²⁴University of California, Santa Barbara, California 93106

²⁵*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213*

²⁶*University of Colorado, Boulder, Colorado 80309-0390*

²⁷*Cornell University, Ithaca, New York 14853*

²⁸*University of Florida, Gainesville, Florida 32611*

(Received 20 June 2000)

Using 13.4 fb^{-1} of data collected with the CLEO detector at the Cornell Electron Storage Ring, we have observed 300 events for the two-photon production of ground-state pseudoscalar charmonium in the decay $\eta_c \rightarrow K_S^0 K^\mp \pi^\pm$. We have measured the η_c mass to be $[2980.4 \pm 2.3 \text{ (stat)} \pm 0.6 \text{ (syst)}] \text{ MeV}$ and its full width as $[27.0 \pm 5.8 \text{ (stat)} \pm 1.4 \text{ (syst)}] \text{ MeV}$. We have determined the two-photon partial width of the η_c meson to be $[7.6 \pm 0.8 \text{ (stat)} \pm 0.4 \text{ (syst)} \pm 2.3 \text{ (br)}] \text{ keV}$, with the last uncertainty associated with the decay branching fraction.

PACS numbers: 14.40.Gx, 13.20.Gd

In this Letter, we report a study of two-photon production of the ground-state pseudoscalar charmonium, i.e., $\gamma\gamma \rightarrow \eta_c$. The two spacelike photons are radiated by e^+ and e^- beams, each at an energy of approximately 5.3 GeV. The charmonium spectrum is an ideal testing ground for quantum chromodynamics (QCD) calculations, and producing C-even charmonium states through $\gamma\gamma$ fusion provides a clean environment for this purpose.

The two-photon partial width of the η_c meson can be expressed in next-to-leading order (NLO) perturbative QCD (PQCD), in terms of the e^+e^- partial width of the J/ψ meson, as [1]

$$\frac{\Gamma_{\gamma\gamma}^{\eta_c}}{\Gamma_{ee}^{\psi}} = \frac{4}{3} (1 + 1.96\alpha_s/\pi) \times \frac{|\Psi_{\eta_c}(0)|^2}{|\Psi_{\psi}(0)|^2}. \quad (1)$$

Using the world average value [2] of Γ_{ee}^{ψ} , a value of the strong coupling constant α_s evaluated at the charm mass scale [1] of (0.28 ± 0.02) , and the assumption that the two 1S wave functions, Ψ , are the same at the origin, this relationship predicts $\Gamma_{\gamma\gamma}^{\eta_c} = (8.2 \pm 0.6) \text{ keV}$.

The total width of the η_c meson can be assumed to be dominated by its two-gluon component, i.e., $\Gamma_{\text{tot}}^{\eta_c} \approx \Gamma_{gg}^{\eta_c}$. The ratio of the rates for $\eta_c \rightarrow gg$ and $\eta_c \rightarrow \gamma\gamma$ is an especially clean prediction of PQCD because the dependencies of these rates on the wave functions and non-perturbative factors are identical in the numerator and denominator. The ratio depends only on the coupling constants and has been calculated in NLO [1],

$$\frac{\Gamma_{\text{tot}}^{\eta_c}}{\Gamma_{\gamma\gamma}^{\eta_c}} \approx \frac{9\alpha_s^2}{8\alpha^2} \times \frac{(1 + 4.8\alpha_s/\pi)}{(1 - 3.4\alpha_s/\pi)}. \quad (2)$$

Using the value of $\Gamma_{\gamma\gamma}^{\eta_c}$ estimated in NLO gives $\Gamma_{\text{tot}}^{\eta_c}$ as $(28 \pm 6) \text{ MeV}$; using instead the world average value [2] of $\Gamma_{\gamma\gamma}^{\eta_c}$, one obtains an estimate of $\Gamma_{\text{tot}}^{\eta_c}$ as $(26 \pm 6) \text{ MeV}$. A calculation with fully relativistic decay amplitudes and a sophisticated QCD potential model [3] predicts $\Gamma_{\text{tot}}^{\eta_c} \approx 23 \text{ MeV}$. The current world average [2] of $\Gamma_{\text{tot}}^{\eta_c} = 13.2_{-3.2}^{+3.8} \text{ MeV}$ disagrees with these theoretical expectations. A precise measurement of the full width and two-photon partial width of the η_c is important for the verification of these PQCD calculations and approximations.

In the two-photon process $e^+e^- \rightarrow e^+e^-\gamma\gamma \rightarrow e^+e^-\eta_c$, the photon propagators dictate that the cross

section naturally peaks at low momentum transfer, so the photons are almost real (“on shell”). The incident leptons are scattered at very low angles and continue traveling down the beam pipe undetected. Such “untagged” events typically have low net transverse momentum and low visible energy. The production of the η_c meson in this untagged two-photon process was searched for in the $K_S^0 K^\mp \pi^\pm$ decay mode.

The data used in this study correspond to an integrated luminosity of 13.4 fb^{-1} and were collected with two configurations (CLEO II [4] and CLEO II.V [5]) of the CLEO detector at the Cornell Electron Storage Ring (CESR). Approximately one-third of the data were taken with the CLEO II configuration. The detector components most useful for this study were the concentric tracking devices for charged particles, operating in a 1.5 T superconducting solenoid. For CLEO II, this tracking system consisted of a 6-layer straw tube chamber, a 10-layer precision drift chamber, and a 51-layer main drift chamber. The main drift chamber also provided measurements of the specific ionization loss, dE/dx , used for particle identification. For CLEO II.V, the straw tube chamber was replaced by a 3-layer, double-sided silicon vertex detector and the gas in the main drift chamber was changed from a 50:50 argon-ethane mixture to a 60:40 helium-propane mixture. These changes gave rise to significant improvements in the momentum and dE/dx resolutions for charged tracks. Photons were detected using the high-resolution electromagnetic calorimeter consisting of 7800 CsI crystals. The Monte Carlo simulation of the CLEO detector response was based upon GEANT [6]. Simulated events were processed in the same manner as the data to determine the $\eta_c \rightarrow K_S^0 K^\mp \pi^\pm$ detection efficiency and the $K_S^0 K^\mp \pi^\pm$ mass resolution at the η_c mass.

The K_S^0 vertex was reconstructed from its decay to $\pi^+\pi^-$ and was required to be displaced from the e^+e^- interaction point; the amount of this displacement varied with detector configuration but was $\approx 1.5 \text{ mm}$. The K_S^0 candidate was also required to be within 4 standard deviations (σ) of the known K_S^0 mass [2]; here σ was determined on an event-by-event basis from the momenta measurements. Furthermore, the K_S^0 momentum vector was required to point back to the interaction point. Of the two remaining charged tracks, the K^\mp and π^\pm candidates,

the one with lower momentum was typically uniquely identified using the particle's specific ionization (dE/dx). This fixed the identity of the only remaining unidentified track, because the presence of the K_S^0 dictated that exactly one of these two be a kaon to conserve strangeness in the event.

The background from processes other than two-photon production was suppressed by requiring that the η_c candidate have net transverse momentum less than 0.6 GeV/ c and that visible energy in the event be less than 6 GeV. Also, because the final state had no expected energy deposits in the calorimeter from neutral particles, the total calorimeter energy in the event not matched to charged tracks was required to be less than 0.6 GeV.

For the mass measurement only, we restricted ourselves to those events in which the K^\mp and π^\pm daughters of the η_c candidate traversed all layers of the tracking volume. The K_S^0 daughter pions were not required to satisfy the same criterion, in that the kinematic fitting of the K_S^0 decay corrected for any possible momentum mismeasurement of the daughter tracks. We did not make any such requirements while measuring $\Gamma_{\text{tot}}^{\eta_c}$ and $\Gamma_{\gamma\gamma}^{\eta_c}$, because these quantities are relatively insensitive to precise measurements of the track momenta; the distribution of candidate invariant masses for the determination of these two quantities is shown in Fig. 1.

We fitted the background with a power law function ($A \cdot W_{\gamma\gamma}^n$, with $W_{\gamma\gamma}$ the $K_S^0 K^\mp \pi^\pm$ invariant mass and A a multiplicative constant) and the signal with a spin-0 relativistic Breit-Wigner function (describing the natural line shape) convolved with a double Gaussian function (to take into account the detector resolution). The parameters for this double Gaussian were obtained from a Monte Carlo




FIG. 1. Results of the simultaneous fit to η_c candidates in (a) CLEO II and (b) CLEO II.V for width and yield measurements, with a combined $\chi^2/\text{d.o.f.} = 226/243$.

sample that had the η_c intrinsic width set to zero. We performed a simultaneous, binned, maximum-likelihood fit to the invariant mass distributions for the CLEO II and CLEO II.V datasets, constraining the physical variables M_{η_c} , $\Gamma_{\text{tot}}^{\eta_c}$, and $\Gamma_{\gamma\gamma}^{\eta_c}$ in the two datasets to be the same. The constraint on $\Gamma_{\gamma\gamma}^{\eta_c}$ was accomplished by requiring the ratio of the fitted yields to be the same as the ratio of the integrated luminosities of the two data sets times the efficiencies as determined from our simulations. The invariant mass resolution was approximately 9 MeV in CLEO II and 7 MeV in CLEO II.V. The bin width for fitting was chosen as approximately the average of these two resolutions.

As noted above, two separate sets of fits were performed, one for the measurements of $\Gamma_{\text{tot}}^{\eta_c}$ and $\Gamma_{\gamma\gamma}^{\eta_c}$ and another for the determination of the η_c mass. The full width and yield were obtained from the distributions shown in Fig. 1, with the total observed yield being $N_{\text{obs}} = 300 \pm 32$. The fit to the width gives $\Gamma_{\text{tot}}^{\eta_c} = (27.0 \pm 5.8)$ MeV, with the uncertainty being only statistical. The two-photon partial width was determined by first correcting for the detector efficiency, ϵ , and then dividing by the number of events expected, N_1 , for a two-photon partial width of 1 keV:

$$\Gamma_{\gamma\gamma}^{\eta_c} = N_{\text{obs}} / (\epsilon \cdot N_1). \quad (3)$$

The quantity N_1 was determined using

$$N_1 = \mathcal{L} \cdot \mathcal{B}_{\eta_c} \cdot \mathcal{B}_{K_S} \cdot \sigma_{e^+e^- \rightarrow e^+e^-\eta_c}; \quad (4)$$

\mathcal{L} is the integrated luminosity. The cross section for $e^+e^- \rightarrow e^+e^-\eta_c$ was obtained from Monte Carlo simulation, using the formalism of Budnev *et al.* [7] and setting $\Gamma_{\gamma\gamma}^{\eta_c} = 1$ keV; this choice of this value has no effect on the extracted result. Also, $\mathcal{B}_{\eta_c} \equiv \mathcal{B}(\eta_c \rightarrow K_S^0 K^\mp \pi^\pm)$ and $\mathcal{B}_{K_S} \equiv \mathcal{B}(K_S^0 \rightarrow \pi^+ \pi^-)$, with the world average values [2] used. This procedure gives $\Gamma_{\gamma\gamma}^{\eta_c} = (7.6 \pm 0.8)$ keV, with this uncertainty coming from statistics only.

The mass was obtained from fits to the more restrictive set of events, as described above, with a total signal size of 195 ± 24 , yielding $M_{\eta_c} = (2980.4 \pm 2.3)$ MeV, the uncertainty being statistical only.

Possible sources of systematic uncertainty for the measured mass, full width, and two-photon partial width were studied. The results are summarized in Table I, in which the individual uncertainties are added in quadrature to obtain the total systematic uncertainty.

The mass calibration of our detector was checked by measuring the masses of the well known K_S^0 , ϕ , D^+ , and J/ψ mesons using decay modes involving only charged tracks. The measurements were found to be in good agreement with their respective world averages when we limited ourselves to events in which the charged tracks traversed all layers of the tracking volume. The fitted mass including events outside this more restrictive detector volume is consistent with the value we quote, but with substantially larger systematic uncertainties. Our particle identification procedure did not introduce any significant systematic bias to the mass measurement. The measured mass was also

TABLE I. Systematic uncertainties in the three measurements. The overall value is obtained by adding the individual contributions in quadrature.

Source of systematic uncertainty	M_{η_c} (MeV)	$\Gamma_{\text{tot}}^{\eta_c}$ (MeV)	$\Gamma_{\gamma\gamma}^{\eta_c}$ (keV)
Mass calibration	0.6	<0.1	<0.1
Particle identification	<0.1	1.3	0.1
Signal shape	<0.1	0.3	0.1
Detector resolution	<0.1	0.3	<0.1
Trigger	0.2
Tracking	0.2
Resonant substructure	<0.1	<0.1	0.2
Luminosity	0.1
K_S^0 selection	<0.1	<0.1	0.1
Event selection	<0.1	<0.1	<0.1
Total	0.6	1.4	0.4

insensitive to variations in the signal shape used to fit the η_c resonance.

The systematic uncertainties in the measurement of the full width were dominated by effects due to the mass resolution, the particle identification procedure, and the signal shape used in the fit. Our ability to predict the actual mass resolution was tested by studying the reconstructed D^+ mesons in Monte Carlo simulation and data; the agreement was found to be better than 0.1 MeV.

The particle identification procedure was unable to distinguish between a charged kaon and a charged pion if the track momentum was above 0.8 GeV/ c , for which the expected ionization losses are nearly equal for the two species. This led to a broadening of the reconstructed resonance and was taken into account by the wider Gaussian of the double Gaussian resolution function. This effect was limited to less than 5% of the events. We estimated the possible uncertainty due to this misassignment of particle species by completely removing the fraction of events having two possible η_c candidates and assigning the corresponding change in the measured width as the systematic uncertainty.

The accuracy of the fitting method in extracting the Breit-Wigner width of the resonance was checked by extracting the η_c widths from sets of simulation events generated with different intrinsic widths. We varied the parameters of the signal shape used to fit the η_c resonance within their uncertainties, derived from a comparison of the fit to D^+ meson decays in Monte Carlo simulation and data, to estimate the effect on the measured width. The measured width from the more restrictive set of events used for mass measurement was within 0.2 MeV of the corresponding measurement using the full sample, and no significant correlation was found between the measured mass and full width.

There were several sources of uncertainty for the estimation of the efficiency, which in turn affected the measurement of $\Gamma_{\gamma\gamma}^{\eta_c}$. These were dominated by the uncertainties in the tracking and trigger efficiencies. The effect of a possible presence of resonant substructure

($K^*\bar{K}$) in η_c decay was studied and was found to give an insignificant variation in the detection efficiency. We estimated the uncertainty in the measured partial width from this effect by considering the possibility that all the η_c mesons decay through $K^*\bar{K}$. Our initial investigation showed roughly a third of the $\eta_c \rightarrow K_S^0 K^\mp \pi^\pm$ events proceed via $K^*(1430)$ with no evidence for any $K^*(892)$; a detailed analysis of this substructure is beyond the scope of this Letter.

The selection requirements on total visible energy and unmatched energy clusters were shown by simulation to be essentially 100% efficient for our signal process and free of systematic bias. Possible bias from the transverse momentum requirement was investigated by changing the nature of the form factor in the simulation and shown to also be negligible.

In our analysis, we investigated the possible effects of interference between the $K_S^0 K^\mp \pi^\pm$ resonant and nonresonant final states. From the distribution of net transverse momentum for events in the sidebands of the signal line shape, we estimated that one-third of the background events were not of the type $\gamma\gamma \rightarrow K_S^0 K^\mp \pi^\pm$; these included events with at least one missing particle (π_0 , γ) as well as events of the type $e^+e^- \rightarrow$ hadrons, $e^+e^- \rightarrow \tau^+\tau^-$, and $\gamma\gamma \rightarrow \tau^+\tau^-$. Such events cannot have interference with our signal $K_S^0 K^\mp \pi^\pm$ events. Study of the helicity angle distributions in the η_c rest frame indicated that the sideband events predominantly have $J = 2$ (or higher) while our signal has $J = 0$. Because of the preferential production of states with natural parity in two-photon untagged processes, a majority of the remaining background events were expected to have the natural spin parity (0^+) compared to the unnatural spin parity (0^-) of the signal events. The acceptance of our detector is symmetric in polar angle and uniform in azimuth, making the interference between these states of opposite parity vanish. We did not include any possible effects due to interference on the measured mass, full width, and two-photon partial width. Further, the signal shape showed no distortions and the goodness of fit to the hypothesis that ignored interference was very good, as shown in Fig. 1.

In summary, we have measured the mass, full width, and two-photon partial width of the η_c produced in two-photon collisions. The mass measurement of $[2980.4 \pm 2.3 \text{ (stat)} \pm 0.6 \text{ (syst)}]$ MeV compares well with the world average [2] of (2979.8 ± 2.1) MeV. The measured total width of $[27.0 \pm 5.8 \text{ (stat)} \pm 1.4 \text{ (syst)}]$ MeV disagrees with the world average [2] of $13.2_{-3.2}^{+3.8}$ MeV, which consists of measurements with large relative uncertainties (40%–100%). Our measured width is consistent with theoretical expectations [1,3]. The measured two-photon partial width of the η_c meson of $[7.6 \pm 0.8 \text{ (stat)} \pm 0.4 \text{ (syst)}]$ keV agrees well with the world average [2] of $7.5_{-1.4}^{+1.6}$ keV and theoretically expected values and is a significant improvement in terms of experimental precision. We use the world average [2] of

the $\eta_c \rightarrow K\bar{K}\pi$ branching fraction of $(5.5 \pm 1.7)\%$. The uncertainty in $\Gamma_{\gamma\gamma}^{\eta_c}$ due to the uncertainty in this branching fraction is ± 2.3 keV and is stated separately from the other contributions. From the ratio of our measured full width and two-photon partial width, we have extracted α_s at the charm mass scale to be 0.285 ± 0.025 , for which we have added our sources of uncertainty in quadrature. We have used the NLO calculation in Eq. (2) to estimate α_s , thus making the result dependent on renormalization scheme and scale; we have not included such theoretical uncertainties in our quoted value.

Our measurements of $\Gamma_{\text{tot}}^{\eta_c}$ and $\Gamma_{\gamma\gamma}^{\eta_c}$ show that PQCD calculations are able to reliably predict the ratios of the decay rates of a heavy quarkonium system, where nonperturbative effects cancel.

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the National Science Foundation, the U.S. Department of Energy, the Research Corporation, the Natural Sciences and Engineering Research Council of Canada, the A.P. Sloan Foun-

ation, the Swiss National Science Foundation, the Texas Advanced Research Program, and the Alexander von Humboldt Stiftung.

*Permanent address: University of Cincinnati, Cincinnati, OH 45221.

†Permanent address: Massachusetts Institute of Technology, Cambridge, MA 02139.

- [1] W. Kwong *et al.*, Phys. Rev. D **37**, 3210 (1988).
- [2] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. C **3**, 1 (1998).
- [3] S. N. Gupta, J. M. Johnson, and W. W. Repko, Phys. Rev. D **54**, 2075 (1996).
- [4] CLEO Collaboration, Y. Kubota *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **320**, 66 (1992).
- [5] T. Hill, Nucl. Instrum. Methods Phys. Res., Sect. A **418**, 32 (1998).
- [6] R. Brun *et al.*, GEANT 3.15, CERN Report No. DD/EE/84-1, 1987.
- [7] V. M. Budnev *et al.*, Phys. Rep. C **15**, 181 (1975).