

## First Observation of the Decay $\tau^- \rightarrow K^- \eta \nu_\tau$

J. Bartelt,<sup>1</sup> S. E. Csorna,<sup>1</sup> V. Jain,<sup>1</sup> S. Marka,<sup>1</sup> A. Freyberger,<sup>2</sup> D. Gibaut,<sup>2</sup> K. Kinoshita,<sup>2</sup> P. Pomianowski,<sup>2</sup> S. Schrenk,<sup>2</sup> D. Cinabro,<sup>3</sup> B. Barish,<sup>4</sup> M. Chadha,<sup>4</sup> S. Chan,<sup>4</sup> G. Eigen,<sup>4</sup> J. S. Miller,<sup>4</sup> C. O'Grady,<sup>4</sup> M. Schmidtler,<sup>4</sup> J. Urheim,<sup>4</sup> A. J. Weinstein,<sup>4</sup> F. Würthwein,<sup>4</sup> D. M. Asner,<sup>5</sup> M. Athanas,<sup>5</sup> D. W. Bliss,<sup>5</sup> W. S. Brower,<sup>5</sup> G. Masek,<sup>5</sup> H. P. Paar,<sup>5</sup> J. Gronberg,<sup>6</sup> C. M. Korte,<sup>6</sup> R. Kutschke,<sup>6</sup> S. Menary,<sup>6</sup> R. J. Morrison,<sup>6</sup> S. Nakanishi,<sup>6</sup> H. N. Nelson,<sup>6</sup> T. K. Nelson,<sup>6</sup> C. Qiao,<sup>6</sup> J. D. Richman,<sup>6</sup> D. Roberts,<sup>6</sup> A. Ryd,<sup>6</sup> H. Tajima,<sup>6</sup> M. S. Witherell,<sup>6</sup> R. Balest,<sup>7</sup> K. Cho,<sup>7</sup> W. T. Ford,<sup>7</sup> M. Lohner,<sup>7</sup> H. Park,<sup>7</sup> P. Rankin,<sup>7</sup> J. Roy,<sup>7</sup> J. G. Smith,<sup>7</sup> J. P. Alexander,<sup>8</sup> C. Bebek,<sup>8</sup> B. E. Berger,<sup>8</sup> K. Berkelman,<sup>8</sup> K. Bloom,<sup>8</sup> D. G. Cassel,<sup>8</sup> H. A. Cho,<sup>8</sup> D. M. Coffman,<sup>8</sup> D. S. Crowcroft,<sup>8</sup> M. Dickson,<sup>8</sup> P. S. Drell,<sup>8</sup> D. J. Dumas,<sup>8</sup> R. Ehrlich,<sup>8</sup> R. Elia,<sup>8</sup> P. Gaidarev,<sup>8</sup> R. S. Galik,<sup>8</sup> B. Gittelman,<sup>8</sup> S. W. Gray,<sup>8</sup> D. L. Hartill,<sup>8</sup> B. K. Heltsley,<sup>8</sup> C. D. Jones,<sup>8</sup> S. L. Jones,<sup>8</sup> J. Kandaswamy,<sup>8</sup> N. Katayama,<sup>8</sup> P. C. Kim,<sup>8</sup> D. L. Kreinick,<sup>8</sup> T. Lee,<sup>8</sup> Y. Liu,<sup>8</sup> G. S. Ludwig,<sup>8</sup> J. Masui,<sup>8</sup> J. Mevissen,<sup>8</sup> N. B. Mistry,<sup>8</sup> C. R. Ng,<sup>8</sup> E. Nordberg,<sup>8</sup> J. R. Patterson,<sup>8</sup> D. Peterson,<sup>8</sup> D. Riley,<sup>8</sup> A. Soffer,<sup>8</sup> C. Ward,<sup>8</sup> P. Avery,<sup>9</sup> C. Prescott,<sup>9</sup> S. Yang,<sup>9</sup> J. Yelton,<sup>9</sup> G. Brandenburg,<sup>10</sup> R. A. Briere,<sup>10</sup> T. Liu,<sup>10</sup> M. Saulnier,<sup>10</sup> R. Wilson,<sup>10</sup> H. Yamamoto,<sup>10</sup> T. E. Browder,<sup>11</sup> F. Li,<sup>11</sup> J. L. Rodriguez,<sup>11</sup> T. Bergfeld,<sup>12</sup> B. I. Eisenstein,<sup>12</sup> J. Ernst,<sup>12</sup> G. E. Gladding,<sup>12</sup> G. D. Gollin,<sup>12</sup> M. Palmer,<sup>12</sup> M. Selen,<sup>12</sup> J. J. Thaler,<sup>12</sup> K. W. Edwards,<sup>13</sup> K. W. McLean,<sup>13</sup> M. Ogg,<sup>13</sup> A. Bellerive,<sup>14</sup> D. I. Britton,<sup>14</sup> R. Janicek,<sup>14</sup> D. B. MacFarlane,<sup>14</sup> P. M. Patel,<sup>14</sup> B. Spaan,<sup>14</sup> A. J. Sadoff,<sup>15</sup> R. Ammar,<sup>16</sup> P. Baringer,<sup>16</sup> A. Bean,<sup>16</sup> D. Besson,<sup>16</sup> D. Coppage,<sup>16</sup> N. Coptly,<sup>16</sup> R. Davis,<sup>16</sup> N. Hancock,<sup>16</sup> S. Kotov,<sup>16</sup> I. Kravchenko,<sup>16</sup> N. Kwak,<sup>16</sup> S. Anderson,<sup>17</sup> Y. Kubota,<sup>17</sup> M. Lattery,<sup>17</sup> J. K. Nelson,<sup>17</sup> S. Patton,<sup>17</sup> R. Poling,<sup>17</sup> T. Riehle,<sup>17</sup> V. Savinov,<sup>17</sup> M. S. Alam,<sup>18</sup> I. J. Kim,<sup>18</sup> Z. Ling,<sup>18</sup> A. H. Mahmood,<sup>18</sup> J. J. O'Neill,<sup>18</sup> H. Severini,<sup>18</sup> C. R. Sun,<sup>18</sup> S. Timm,<sup>18</sup> F. Wappler,<sup>18</sup> J. E. Duboscq,<sup>19</sup> R. Fulton,<sup>19</sup> D. Fujino,<sup>19</sup> K. K. Gan,<sup>19</sup> K. Honscheid,<sup>19</sup> H. Kagan,<sup>19</sup> R. Kass,<sup>19</sup> J. Lee,<sup>19</sup> M. Sung,<sup>19</sup> A. Undrus,<sup>19,\*</sup> C. White,<sup>19</sup> R. Wanke,<sup>19</sup> A. Wolf,<sup>19</sup> M. M. Zoeller,<sup>19</sup> X. Fu,<sup>20</sup> B. Nemati,<sup>20</sup> S. J. Richichi,<sup>20</sup> W. R. Ross,<sup>20</sup> P. Skubic,<sup>20</sup> M. Wood,<sup>20</sup> M. Bishai,<sup>21</sup> J. Fast,<sup>21</sup> E. Gerndt,<sup>21</sup> J. W. Hinson,<sup>21</sup> T. Miao,<sup>21</sup> D. H. Miller,<sup>21</sup> M. Modesitt,<sup>21</sup> E. I. Shibata,<sup>21</sup> I. P. J. Shipsey,<sup>21</sup> P. N. Wang,<sup>21</sup> M. Yurko,<sup>21</sup> L. Gibbons,<sup>22</sup> S. D. Johnson,<sup>22</sup> Y. Kwon,<sup>22</sup> S. Roberts,<sup>22</sup> E. H. Thorndike,<sup>22</sup> C. P. Jessop,<sup>23</sup> K. Lingel,<sup>23</sup> H. Marsiske,<sup>23</sup> M. L. Perl,<sup>23</sup> S. F. Schaffner,<sup>23</sup> R. Wang,<sup>23</sup> T. E. Coan,<sup>24</sup> J. Dominick,<sup>24</sup> V. Fadeyev,<sup>24</sup> I. Korolkov,<sup>24</sup> M. Lambrecht,<sup>24</sup> S. Sanghera,<sup>24</sup> V. Shelkov,<sup>24</sup> R. Stroynowski,<sup>24</sup> I. Volobouev,<sup>24</sup> G. Wei,<sup>24</sup> M. Artuso,<sup>25</sup> A. Efimov,<sup>25</sup> M. Gao,<sup>25</sup> M. Goldberg,<sup>25</sup> R. Greene,<sup>25</sup> D. He,<sup>25</sup> N. Horwitz,<sup>25</sup> S. Kopp,<sup>25</sup> G. C. Moneti,<sup>25</sup> R. Mountain,<sup>25</sup> Y. Mukhin,<sup>25</sup> S. Playfer,<sup>25</sup> T. Skwarnicki,<sup>25</sup> S. Stone,<sup>25</sup> and X. Xing<sup>25</sup>

(CLEO Collaboration)

<sup>1</sup>Vanderbilt University, Nashville, Tennessee 37235

<sup>2</sup>Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

<sup>3</sup>Wayne State University, Detroit, Michigan 48202

<sup>4</sup>California Institute of Technology, Pasadena, California 91125

<sup>5</sup>University of California, San Diego, La Jolla, California 92093

<sup>6</sup>University of California, Santa Barbara, California 93106

<sup>7</sup>University of Colorado, Boulder, Colorado 80309-0390

<sup>8</sup>Cornell University, Ithaca, New York 14853

<sup>9</sup>University of Florida, Gainesville, Florida 32611

<sup>10</sup>Harvard University, Cambridge, Massachusetts 02138

<sup>11</sup>University of Hawaii at Manoa, Honolulu, Hawaii 96822

<sup>12</sup>University of Illinois, Champaign-Urbana, Illinois 61801

<sup>13</sup>Carleton University, Ottawa, Ontario, Canada K1S 5B6  
and the Institute of Particle Physics, Montréal, Québec, Canada

<sup>14</sup>McGill University, Montréal, Québec, Canada H3A 2T8  
and the Institute of Particle Physics, Montréal, Québec, Canada

<sup>15</sup>Ithaca College, Ithaca, New York 14850

<sup>16</sup>University of Kansas, Lawrence, Kansas 66045

<sup>17</sup>University of Minnesota, Minneapolis, Minnesota 55455

<sup>18</sup>State University of New York at Albany, Albany, New York 12222

<sup>19</sup>The Ohio State University, Columbus, Ohio 43210

<sup>20</sup>University of Oklahoma, Norman, Oklahoma 73019

<sup>21</sup>Purdue University, West Lafayette, Indiana 47907

<sup>22</sup>University of Rochester, Rochester, New York 14627

<sup>23</sup>Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309

<sup>24</sup>Southern Methodist University, Dallas, Texas 75275

<sup>25</sup>Syracuse University, Syracuse, New York 13244

(Received 12 January 1996)

The decay  $\tau^- \rightarrow K^- \eta \nu_\tau$  has been observed with the CLEO II detector. The  $\eta$  meson is reconstructed using two decay channels,  $\eta \rightarrow \gamma\gamma$  and  $\pi^+ \pi^- \pi^0$ . The measured branching fraction is  $B(\tau^- \rightarrow K^- \eta \nu_\tau) = (2.6 \pm 0.5 \pm 0.4) \times 10^{-4}$ , somewhat higher than theoretical estimates. An improved upper limit for the second-class-current decay  $\tau^- \rightarrow \pi^- \eta \nu_\tau$  is set,  $B(\tau^- \rightarrow \pi^- \eta \nu_\tau) < 1.4 \times 10^{-4}$  at 95% C.L., consistent with theoretical expectations. [S0031-9007(96)00324-9]

PACS numbers: 13.35.Dx, 14.40.Aq, 14.60.Fg

The decays of the  $\tau$  lepton involving  $\eta$  mesons are of particular interest since they are suppressed in comparison with all-pion decays. The decay  $\tau^- \rightarrow \pi^- \eta \nu_\tau$  [1] violates  $G$ -parity conservation and proceeds via a second-class current. The standard model predicts a branching fraction of  $(1.2-1.5) \times 10^{-5}$  [2,3] for this decay mode. Because of SU(3) symmetry breaking, there is no  $G$ -parity constraint on the analogous Cabibbo-suppressed decay,  $\tau^- \rightarrow K^- \eta \nu_\tau$ , so this branching fraction is expected to be larger by an order of magnitude,  $(1.2-1.6) \times 10^{-4}$  [2,4]. Previously, CLEO II measured [5] the branching fraction of the first  $\tau$  decay involving  $\eta$  mesons,  $B(\tau^- \rightarrow \pi^- \eta \pi^0 \nu_\tau) = (1.7 \pm 0.2 \pm 0.2) \times 10^{-3}$ , and set the following upper limits,  $B(\tau^- \rightarrow \pi^- \eta \nu_\tau) < 3.4 \times 10^{-4}$  and  $B(\tau^- \rightarrow K^- \eta \nu_\tau) < 4.7 \times 10^{-4}$  at 95% C.L. Using a 4 times larger data sample we report in this Letter the first measurement of the decay  $\tau^- \rightarrow K^- \eta \nu_\tau$  and an improved upper limit for  $\tau^- \rightarrow \pi^- \eta \nu_\tau$ .

The data used in this analysis were collected with the CLEO II detector from  $e^+e^-$  collisions at the Cornell Electron-positron Storage Ring (CESR) at a center-of-mass energy  $E_{\text{cm}} \sim 10.6$  GeV. The total integrated luminosity of the sample is  $3.5 \text{ fb}^{-1}$ , corresponding to the production of  $3.2 \times 10^6$   $\tau^+\tau^-$  events. The CLEO II detector has been described in detail elsewhere [6]. Charged particles are identified using the drift chamber ( $dE/dx$ ) and time-of-flight (TOF) scintillation counters. For hadrons, the TOF system provides  $K/\pi$  separation of  $>2\sigma$  (standard deviation) for particle momenta below 1.07 GeV/ $c$ . The  $dE/dx$  measurement provides  $K/\pi$  separation of  $>2\sigma$  for particle momenta below 0.75 GeV/ $c$ ; for the relativistic rise region above 2.0 GeV/ $c$ , it provides  $\sim 1.8\sigma$   $K/\pi$  separation.

We reconstruct the  $\eta$  meson using the  $\gamma\gamma$  and  $\pi^+ \pi^- \pi^0$  decay channels. The  $\tau^+ \tau^-$  candidate events must contain two or four charged tracks and have zero net charge. To reject beam-gas events, the distance of the closest approach of each charged track to the IP must be within 0.5 cm transverse to the beam and 5 cm along the beam direction. We divide each event into two hemispheres (tag vs signal) using the plane perpendicular to the thrust axis, calculated using both charged tracks and photons. The photons are defined as energy clusters in the calorimeter with at least 60 MeV in the barrel,  $|\cos\theta| < 0.80$ , and 100 MeV in the end cap,  $0.80 < |\cos\theta| < 0.95$ . The opening angle

between the total momentum vectors of the two  $\tau$  decay products must be greater than  $120^\circ$ . The tag hemisphere must contain only one charged particle (plus any number of photons), and in the events with  $\eta \rightarrow \gamma\gamma$ , its momentum ( $p_{\text{tag}}$ ) must be greater than 1.0 GeV/ $c$ . The hadronic background is suppressed by the requirements that the total invariant mass of the particles in the tag and signal hemispheres be less than 1.2 and 1.7 GeV, respectively. Backgrounds from two-photon production, Bhabha scattering, and hadronic events are suppressed by the requirements on the total visible energy,  $0.25 < E_{\text{tot}}/E_{\text{cm}} < 0.85$ , and on the measured net transverse momentum of the event,  $p_\perp > 0.5$  GeV/ $c$ . All charged particles and photons are included in the calculation of the kinematic variables. For the decay  $\tau^- \rightarrow \pi^- \eta \nu_\tau$ , we reconstruct the  $\eta$  meson using the  $\gamma\gamma$  decay channel only and select the events with a lepton tag in order to further suppress the hadronic background. An electron candidate must have an energy deposition in the calorimeter consistent with its measured momentum. A muon candidate must penetrate more than three absorption lengths of iron.

The kaon and pion candidates in the signal hemisphere are restricted to be in the central part of the detector,  $|\cos\theta| < 0.81$ . Particle identification is based on a confidence level ratio which is constructed from the confidence levels for  $\pi$  and  $K$  hypotheses,  $C_\pi$  and  $C_K$ . The confidence level ratio for  $K$  is

$$R_K = \frac{C_K}{C_\pi + C_K},$$

and similarly for  $\pi$  ( $R_\pi = 1 - R_K$ ). The confidence level is computed from the  $\chi^2$  probability for a particle hypothesis using a combination of the TOF and  $dE/dx$  information. We use a sample of kaons from  $D^{*+} \rightarrow D^0 \pi^+ \rightarrow K^- \pi^+ \pi^+$  and pions from  $K_S$  decays in hadronic events to study the optimum method of combining the information as a function of momentum. For the momentum ranges 0.2 to 1.0 and 1.3 to 1.7 GeV/ $c$ , we require the track to have  $dE/dx$  information; we also use the TOF information if available. In the momentum region 1.0 to 1.3 GeV/ $c$ , where  $\pi$  and  $K$  have similar  $dE/dx$  energy losses, we use TOF information only. For high momentum tracks, 1.7 to 4.5 GeV/ $c$ , we use  $dE/dx$  information only. A  $K$  candidate is then defined as a particle with  $R_K > 0.5$ ; otherwise the particle is considered a pion. Above 1 GeV/ $c$  where

80% of the kaons from the decay  $\tau^- \rightarrow K^- \eta \nu_\tau$  populate, the  $R_K$  requirement gives an identification efficiency of  $\sim 80\%$  and a misidentification rate ( $\pi$  as  $K$ ) of  $\sim 20\%$ .

The  $\eta$  mesons are reconstructed using barrel photons in the signal hemisphere. If there are more than two photons with energy above 100 MeV, including the end cap, the event is rejected. For the  $\pi^+ \pi^- \pi^0$  decay channel, the invariant mass of the photon pair is constrained to the  $\pi^0$  mass. For the  $\gamma\gamma$  channel, the requirements on the photons are more stringent in order to suppress the background. Each photon must have an energy above 150 MeV and a lateral profile of energy deposition consistent with that expected of a photon. In addition, we do not use showers that are likely fragments of nearby showers. A signal photon may not combine with any other photon to form a  $\pi^0$  ( $|M_{\gamma\gamma} - M_{\pi^0}| < 20$  MeV).

For the  $\gamma\gamma$  channel, the invariant mass spectrum of the two photons accompanying the kaon candidate is shown in Fig. 1(a). The mass spectrum is expressed in standard deviations from the nominal  $\eta$  mass,  $S_{\gamma\gamma} = (M_{\gamma\gamma} - M_\eta)/\sigma_{\gamma\gamma}$ , with the mass resolution  $\sigma_{\gamma\gamma}$  calculated from the measured angle and energy resolution of each photon. An  $\eta$  signal is observed. The curve is a fit [7] to the data using a Gaussian with a low mass tail (to account for shower leakage) over a polynomial background. An  $\eta$  signal is also observed in the  $\pi^+ \pi^- \pi^0$  channel as




FIG. 1. The invariant mass spectrum of the  $\eta$  candidates. Each  $\eta$  candidate is accompanied by a kaon candidate in (a) and (b) and by a pion candidate in (c). The  $\eta$  candidates in (c) are selected with a lepton tag. Each curve shows a fit to the mass spectrum (see text).

shown in Fig. 1(b). For the decay  $\tau^- \rightarrow \pi^- \eta \nu_\tau$ , no  $\eta$  signal is evident as shown in Fig. 1(c). All fits have the  $\eta$  width constrained to the Monte Carlo expectation. For the  $\pi\eta$  analysis, the  $\eta$  mass is also constrained to the Monte Carlo prediction since there is no  $\eta$  signal in the data. The event yield extracted from the fit is summarized in Table I.

As a verification of the  $K\eta$  signal in the  $\eta \rightarrow \gamma\gamma$  decay channel, we show in Fig. 2 the  $R_K$  distribution of the charged track in the signal hemisphere accompanied by an  $\eta$  candidate (sideband subtracted). There is an enhancement at  $R_K = 0$  and 1.0 as expected. The distribution is well described by the Monte Carlo simulation. The enhancement at high  $R_K$  is dominated by the  $K\eta$  decay while at low  $R_K$  it is saturated by the hadronic background.

The detection efficiencies for the candidate events are calculated using a Monte Carlo simulation. The KORALB program is used to generate  $\tau^+ \tau^-$  pairs according to the standard electroweak theory, including  $\alpha^3$  radiative corrections [8]. The decay  $\tau^- \rightarrow K^- \eta \nu_\tau$  [ $\pi^- \eta \nu_\tau$ ] is modeled with the  $K^*(1410)$  [ $a_0(980)$ ] resonance assuming a  $V-A$  weak interaction. The detector response is simulated using the GEANT program [9]. Detector activity not attributable to the  $e^+ e^-$  interaction is modeled by embedding random trigger events obtained during data taking into the generated events. The identification and misidentification efficiencies are calibrated as a function of momentum by comparing the efficiencies measured from the  $D^*$  and  $K_S$  data sample with the Lund Monte Carlo expectations [10].

The hadronic background is estimated from the data, using events in which the invariant mass in one hemisphere is above the  $\tau$  mass [11]. The background to  $\tau^- \rightarrow K^- \eta \nu_\tau$  from two-photon interactions is estimated using the fact that the two-photon events are produced at low  $p_\perp$ . From the paucity of events at  $p_\perp < 0.1$  GeV/ $c$ , we set an upper limit [12] of 2% contamination at 95% C.L.

We use the measured branching fraction [5] for  $\tau^- \rightarrow \pi^- \eta \pi^0 \nu_\tau$  to estimate its contribution to the  $\tau^- \rightarrow K^- \eta \nu_\tau$  sample. The measured branching fractions for  $\tau^- \rightarrow K^- \eta \nu_\tau$  and  $\pi^- \eta \nu_\tau$  presented in Table I are used to estimate the cross-feed backgrounds between the two decay modes. The background from  $\tau^- \rightarrow K^- \pi^0 \eta \nu_\tau$  is neglected since it is expected to be  $\sim 0.1\%$  [2] of the signal. The signals, backgrounds, and detection efficiencies are summarized in Table I. The efficiencies

TABLE I. Summary of signals, backgrounds, detection efficiencies, and branching fractions. All errors are statistical.

	$K^- \gamma\gamma$	$K^- \pi^+ \pi^- \pi^0$	$\pi^- \gamma\gamma$
$\eta$ signal	$61 \pm 11$	$24 \pm 7$	$0_{-0}^{+5.3}$
$q\bar{q}$	$8.2 \pm 3.8$	$5.9 \pm 3.1$	$2.7 \pm 1.9$
$\pi^- \pi^0 \eta$	$3.2 \pm 0.8$	$3.8 \pm 1.0$	$3.9 \pm 0.9$
Cross-feed eff (%)	$1.3 \pm 0.1$	$0.8 \pm 0.1$	$0.8 \pm 0.1$
Eff (%)	$7.6 \pm 0.1$	$3.9 \pm 0.1$	$3.5 \pm 0.1$
$B$ ( $10^{-4}$ )	$2.6 \pm 0.6$	$2.5 \pm 1.3$	$0_{-0}^{+0.62}$




FIG. 2. The observed  $R_K$  spectrum of the charged particle in the hemisphere containing an  $\eta$  meson after sideband subtraction. The histogram shows the Monte Carlo expectation which is a sum of the predictions for  $\tau^- \rightarrow K^- \eta \nu_\tau$  (dashed),  $\tau^- \rightarrow \pi^- \pi^0 \eta \nu_\tau$  (shaded), and  $e^+ e^- \rightarrow q \bar{q}$  (dotted).

exclude the decay branching fractions of  $\eta \rightarrow \gamma\gamma$  and  $\pi^+ \pi^- \pi^0$  [13].

The observed  $K\eta$  mass spectrum for  $\eta \rightarrow \gamma\gamma$  is compared with the Monte Carlo expectation in Fig. 3, without the 1.7 GeV/c mass cut. Also shown is the prediction in which the  $K^*(1410)$  resonance is replaced by a spectral function determined simply from the  $K$ - $\eta$  phase space. The data are consistent with either model.

There are several sources of systematic errors as shown in Table II. These include the uncertainties in the luminosity,  $\tau^+ \tau^-$  cross section, branching fractions of  $\eta \rightarrow \gamma\gamma$  and  $\pi^+ \pi^- \pi^0$  [13], identification efficiency, background subtraction, fitting procedure, acceptance calculation, decay modeling, as well as the uncertainty due to limited Monte Carlo statistics. The systematic error in the identification efficiency has two components: (i) the




FIG. 3. The  $K\eta$  invariant mass spectrum after the subtraction of all backgrounds. The histograms show the Monte Carlo expectations for the  $K^*(1410)$  (solid) and phase space (dashed) models assuming a  $V$ - $A$  weak interaction. The number of events above the  $\tau$  mass is consistent with zero.

statistical uncertainty due to the limited  $D^*$  and  $K_S$  samples, (ii) the potential difference in the efficiencies between  $\tau$  and hadronic events. This is estimated from the dependence of the efficiency in the hadronic events on the number of useful  $dE/dx$  hits and the isolation of the TOF hits from the other charged tracks. There is also a similar systematic error in the misidentification efficiency. This is included in the systematic error of the background subtraction. Also included are the statistical errors in the background estimate (Table II) and the uncertainties in the branching fractions of the decay modes. The systematic error due to the fitting procedure was estimated by comparing the results for various fit ranges and for different orders of polynomial background. We estimated the systematic error in the acceptance calculation using the well-measured decay channels with the similar topology as the reactions under study,  $\tau^- \rightarrow h^- \pi^0 \nu_\tau$  and  $h^- h^- h^+ \pi^0 \nu_\tau$ , where  $h$  can either be a  $\pi$  or  $K$ . By comparing the measured branching fractions with the world averages [14], we assign a systematic error of 5% for the  $K\gamma\gamma$  mode, 7% for the  $K\pi^+ \pi^- \pi^0$  mode, and 11% for the  $\pi\gamma\gamma$  mode. The systematic error in the decay modeling is estimated by comparing the detection efficiencies for the resonance and phase space ( $S$  wave) models.

The branching fraction for  $\tau^- \rightarrow K^- \eta \nu_\tau$  is extracted after correcting for the backgrounds and detection efficiencies. The results are  $(2.6 \pm 0.6 \pm 0.4) \times 10^{-4}$  and  $(2.5 \pm 1.3 \pm 0.7) \times 10^{-4}$  for the  $\gamma\gamma$  and  $\pi^+ \pi^- \pi^0$  channels, where the first error is statistical and the second systematic. Combining these two results yields

$$B(\tau^- \rightarrow K^- \eta \nu_\tau) = (2.6 \pm 0.5 \pm 0.4) \times 10^{-4}.$$

The upper limit on  $\tau^- \rightarrow \pi^- \eta \nu_\tau$  is obtained without correcting for the backgrounds,  $B(\tau^- \rightarrow \pi^- \eta \nu_\tau) < 1.2 \times 10^{-4}$  at 95% C.L. Loosening the limit by one unit of the total systematic error yields

$$B(\tau^- \rightarrow \pi^- \eta \nu_\tau) < 1.4 \times 10^{-4} \text{ at } 95\% \text{ C.L.}$$

In summary, we have measured, for the first time, the branching fraction of  $\tau^- \rightarrow K^- \eta \nu_\tau$  and set a more stringent limit on  $B(\tau^- \rightarrow \pi^- \eta \nu_\tau)$ . The measured branching

TABLE II. Summary of systematic errors (%).

	$K^- \gamma\gamma$	$K^- \pi^- \pi^+ \pi^0$	$\pi^- \gamma\gamma$
$N_{\tau\tau}$	1.4	1.4	1.4
$\eta B$	1.3	2.5	1.3
Identification	2	2	2
Background	9	24	...
Fit	5	8	...
Acceptance	5	7	11
Decay model	6	5	5
MC statistics	2	3	2
Total	13	27	13

fraction is somewhat higher than the theoretical predictions [24] while the upper limit on the second-class-current decay is consistent with theoretical expectations [2,3].

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the National Science Foundation, the U.S. Department of Energy, the Heisenberg Foundation, the Alexander von Humboldt Stiftung, the Natural Sciences and Engineering Research Council of Canada, and the A. P. Sloan Foundation.

---

\*Permanent address: BINP, RU-630090 Novosibirsk, Russia.

[1] In this Letter, charge conjugate states are implied.

[2] A. Pich, Phys. Lett. B **196**, 561 (1987).

[3] S. Tisserant and T.N. Truong, Phys. Lett. **115B**, 264 (1982); H. Neufeld and H. Rupertsberger, Z. Phys. C **68**, 91 (1995).

[4] G.J. Aubrecht II, N. Chahroui, and K. Slanec, Phys. Rev. D **24**, 1318 (1981).

[5] M. Artuso *et al.*, Phys. Rev. Lett. **69**, 3278 (1992).

[6] Y. Kubota *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **320**, 66 (1992).

[7] The peak and width of the mass resolution curve are well modeled by the Monte Carlo simulation, verified using  $\eta$ 's in hadronic events.

[8] S. Jadach and Z. Was, Comput. Phys. Commun. **36**, 191 (1985); **64**, 267 (1991); S. Jadach, J. H. Kuhn, and Z. Was, *ibid.* **64**, 275 (1991).

[9] R. Brun *et al.*, CERN Report No. CERN-DD/EE/84-1, 1987 (unpublished).

[10] T. Sjostrand and M. Bengtsson, Comput. Phys. Commun. **43**, 367 (1987).

[11] We have verified the hadronic background calculation by measuring the branching fraction for  $\tau^- \rightarrow K^- \eta \nu_\tau$  with a lepton tag and obtained a consistent result.

[12] The background is estimated using a simple Monte Carlo simulation based on the equivalent photon approximation [V.M. Budnev *et al.*, Phys. Rep. C **15**, 181 (1975)].

[13] L. Montanet *et al.*, Phys. Rev. D **50**, 1173 (1994).

[14] K. K. Gan, in Proceedings of the International Europhysics Conference on High Energy Physics, Brussels, 1995 (to be published).