

First Measurement of the Form Factors in the Decays $D^0 \rightarrow \rho^- e^+ \nu_e$ and $D^+ \rightarrow \rho^0 e^+ \nu_e$

S. Dobbs,¹ Z. Metreveli,¹ K. K. Seth,¹ A. Tomaradze,¹ T. Xiao,¹ L. Martin,² A. Powell,² G. Wilkinson,² H. Mendez,³ D. M. Asner,⁴ G. Tatishvili,⁴ J. Y. Ge,⁵ G. S. Huang,^{5,*} D. H. Miller,⁵ V. Pavlunin,^{5,†} I. P. J. Shipsey,⁵ B. Xin,⁵ G. S. Adams,⁶ D. Hu,⁶ B. Moziak,⁶ J. Napolitano,⁶ K. M. Ecklund,⁷ J. Insler,⁸ H. Muramatsu,⁸ C. S. Park,⁸ L. J. Pearson,⁸ E. H. Thorndike,⁸ S. Ricciardi,⁹ C. Thomas,^{2,9} M. Artuso,¹⁰ S. Blusk,¹⁰ R. Mountain,¹⁰ T. Skwarnicki,¹⁰ S. Stone,¹⁰ L. M. Zhang,¹⁰ G. Bonvicini,¹¹ D. Cinabro,¹¹ A. Lincoln,¹¹ M. J. Smith,¹¹ P. Zhou,¹¹ J. Zhu,¹¹ P. Naik,¹² J. Rademacker,¹² K. W. Edwards,¹³ K. Randrianarivony,¹³ R. A. Briere,¹⁴ H. Vogel,¹⁴ P. U. E. Onyisi,¹⁵ J. L. Rosner,¹⁵ J. P. Alexander,¹⁶ D. G. Cassel,¹⁶ S. Das,¹⁶ R. Ehrlich,¹⁶ L. Gibbons,¹⁶ S. W. Gray,¹⁶ D. L. Hartill,¹⁶ B. K. Heltsley,¹⁶ D. L. Kreinick,¹⁶ V. E. Kuznetsov,¹⁶ J. R. Patterson,¹⁶ D. Peterson,¹⁶ D. Riley,¹⁶ A. Ryd,¹⁶ A. J. Sadoff,¹⁶ X. Shi,¹⁶ W. M. Sun,¹⁶ J. Yelton,¹⁷ P. Rubin,¹⁸ N. Lowrey,¹⁹ S. Mehrabyan,¹⁹ M. Selen,¹⁹ J. Wiss,¹⁹ J. Libby,²⁰ M. Kornicer,²¹ R. E. Mitchell,²¹ C. M. Tarbert,²¹ D. Besson,²² T. K. Pedlar,²³ D. Cronin-Hennessy,²⁴ and J. Hietala²⁴

(CLEO Collaboration)

¹Northwestern University, Evanston, Illinois 60208, USA

²University of Oxford, Oxford OX1 3RH, United Kingdom

³University of Puerto Rico, Mayaguez, Puerto Rico 00681, USA

⁴Pacific Northwest National Laboratory, Richland, Washington 99352, USA

⁵Purdue University, West Lafayette, Indiana 47907, USA

⁶Rensselaer Polytechnic Institute, Troy, New York 12180, USA

⁷Rice University, Houston, Texas 77005, USA

⁸University of Rochester, Rochester, New York 14627, USA

⁹STFC Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, United Kingdom

¹⁰Syracuse University, Syracuse, New York 13244, USA

¹¹Wayne State University, Detroit, Michigan 48202, USA

¹²University of Bristol, Bristol BS8 1TL, United Kingdom

¹³Carleton University, Ottawa, Ontario K1S 5B6, Canada

¹⁴Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

¹⁵University of Chicago, Chicago, Illinois 60637, USA

¹⁶Cornell University, Ithaca, New York 14853, USA

¹⁷University of Florida, Gainesville, Florida 32611, USA

¹⁸George Mason University, Fairfax, Virginia 22030, USA

¹⁹University of Illinois, Urbana-Champaign, Illinois 61801, USA

²⁰Indian Institute of Technology Madras, Chennai, Tamil Nadu 600036, India

²¹Indiana University, Bloomington, Indiana 47405, USA

²²University of Kansas, Lawrence, Kansas 66045, USA

²³Luther College, Decorah, Iowa 52101, USA

²⁴University of Minnesota, Minneapolis, Minnesota 55455, USA

(Received 13 December 2011; published 26 March 2013)

The beauty to up quark coupling constant $|V_{ub}|$ can be extracted from $B \rightarrow \rho e^+ \nu_e$ combined with the form factors for $D \rightarrow K^* e^+ \nu_e$ and $B \rightarrow V \ell^+ \ell^-$ and $D \rightarrow \rho e^+ \nu_e$. Using the entire CLEO-c $\psi(3770) \rightarrow D\bar{D}$ event sample, corresponding to an integrated luminosity of 818 pb^{-1} and approximately 5.4×10^6 $D\bar{D}$ events, we measure the form factors for the decays $D^0 \rightarrow \rho^- e^+ \nu_e$ and $D^+ \rightarrow \rho^0 e^+ \nu_e$ for the first time and the branching fractions with improved precision. A four-dimensional unbinned maximum likelihood fit determines the form factor ratios to be $V(0)/A_1(0) = 1.48 \pm 0.15 \pm 0.05$ and $A_2(0)/A_1(0) = 0.83 \pm 0.11 \pm 0.04$. Assuming Cabibbo-Kobayashi-Maskawa unitarity, the known D meson lifetimes, and our measured branching fractions we obtain the form factor normalizations $A_1(0)$, $A_2(0)$, and $V(0)$. We also present a measurement of the branching fraction for $D^+ \rightarrow \omega e^+ \nu_e$ with improved precision.

DOI: [10.1103/PhysRevLett.110.131802](https://doi.org/10.1103/PhysRevLett.110.131802)

PACS numbers: 13.20.Fc

A precise determination of the beauty to up quark coupling $|V_{ub}|$ is central to testing the quark mixing sector of the standard model. Quark couplings can be determined from the transition rates of semileptonic decays if strong

interaction effects binding quarks into hadrons parametrized by form factors are known. Exploiting one of the proposed double-ratio techniques [1–3], $D \rightarrow \rho e^+ \nu_e$ form factors, when combined with those of $D \rightarrow K^* e^+ \nu_e$ and

$B \rightarrow V\ell^+\ell^-$, can be used to extract $|V_{ub}|$ from $B \rightarrow \rho e^+\nu_e$. In this Letter, we present the first measurement of the form factors in $D \rightarrow \rho e^+\nu_e$.

The transition rate for $D \rightarrow \rho e^+\nu_e$ decays depends on the charm to down quark coupling $|V_{cd}|$, which is precisely known from unitarity of the Cabibbo-Kobayashi-Maskawa matrix [4], and in the limit of negligible lepton mass, three dominant form factors: two axial and one vector, A_1 , A_2 ,

and V , respectively, which are functions of q^2 , the invariant mass of the lepton-neutrino system. They are not amenable to unquenched lattice quantum chromodynamics calculations due to the large total decay width of the ρ meson, but model predictions exist [5–7].

The differential decay rate of $D \rightarrow \rho e^+\nu_e$ can be expressed in terms of three helicity amplitudes ($H_+(q^2)$, $H_-(q^2)$, and $H_0(q^2)$) [8]:

$$\begin{aligned} \frac{d\Gamma}{dq^2 d\cos\theta_\pi d\cos\theta_e d\chi dm_{\pi\pi}} &= \frac{3}{8(4\pi)^4} G_F^2 |V_{cd}|^2 \frac{p_\rho q^2}{M_D^2} \mathcal{B}(\rho \rightarrow \pi\pi) |\mathcal{B}\mathcal{W}(m_{\pi\pi})|^2 [(1 + \cos\theta_e)^2 \sin^2\theta_\pi |H_+(q^2, m_{\pi\pi})|^2 \\ &+ (1 - \cos\theta_e)^2 \sin^2\theta_\pi |H_-(q^2, m_{\pi\pi})|^2 + 4\sin^2\theta_e \cos^2\theta_\pi |H_0(q^2, m_{\pi\pi})|^2 \\ &+ 4\sin\theta_e(1 + \cos\theta_e) \sin\theta_\pi \cos\theta_\pi \cos\chi H_+(q^2, m_{\pi\pi}) H_0(q^2, m_{\pi\pi}) \\ &- 4\sin\theta_e(1 - \cos\theta_e) \sin\theta_\pi \cos\theta_\pi \cos\chi H_-(q^2, m_{\pi\pi}) H_0(q^2, m_{\pi\pi}) \\ &- 2\sin^2\theta_e \sin^2\theta_\pi \cos 2\chi H_+(q^2, m_{\pi\pi}) H_-(q^2, m_{\pi\pi})], \end{aligned} \quad (1)$$

where G_F is the Fermi constant, p_ρ is the momentum of the ρ in the D rest frame, $\mathcal{B}(\rho \rightarrow \pi\pi)$ is a branching fraction, θ_π is the angle between the π and the D direction in the ρ rest frame, θ_e is the angle between the e^+ and the D direction in the $e^+\nu_e$ rest frame, χ is the acoplanarity angle between the $\pi^+\pi^-$ and $e^+\nu_e$ decay planes, $m_{\pi\pi}$ is the invariant mass of the two pions, and $\mathcal{B}\mathcal{W}(m_{\pi\pi})$ is the Breit-Wigner function that describes the ρ line shape. Following Ref. [9], we use the relativistic form

$$\mathcal{B}\mathcal{W}(m_{\pi\pi}) = \frac{\sqrt{m_0\Gamma_0}(p/p_0)}{m_0^2 - m_{\pi\pi}^2 - im_0\Gamma(m_{\pi\pi})} \frac{B(p)}{B(p_0)}, \quad (2)$$

where m_0 and Γ_0 are the mass and width of the ρ meson [4], p is the momentum of the pion in the $\pi\pi$ rest frame, p_0 is equal to p when $m_{\pi\pi} = m_0$, and $B(p)$ is a Blatt-Weisskopf form factor given by $B(p) = 1/(1 + R^2 p^2)^{1/2}$, with $R = 3 \text{ GeV}^{-1}$, and $\Gamma(m_{\pi\pi}) = (p/p_0)^3 (m_0/m_{\pi\pi}) \times \Gamma_0 [B(p)/B(p_0)]^2$. The interference term between a possible s -wave $\pi\pi$ component and the ρ amplitude has not been included in Eq. (1). Its absence is treated as a source of systematic uncertainty on the measurement.

The helicity amplitudes are related to the form factors

$$H_\pm(q^2) = MA_1(q^2) \mp 2 \frac{M_D p_\rho}{M} V(q^2), \quad (3)$$

$$\begin{aligned} H_0(q^2) &= \frac{1}{2m_{\pi\pi}\sqrt{q^2}} \left[(M_D^2 - m_{\pi\pi}^2 - q^2) MA_1(q^2) \right. \\ &\quad \left. - 4 \frac{M_D^2 p_\rho^2}{M} A_2(q^2) \right], \end{aligned} \quad (4)$$

where M_D is the mass of the D meson and $M = M_D + m_{\pi\pi}$. Because $A_1(q^2)$ is common to all three helicity amplitudes, it is natural to define two form factor ratios as

$$r_V = \frac{V(0)}{A_1(0)} \quad \text{and} \quad r_2 = \frac{A_2(0)}{A_1(0)}. \quad (5)$$

We assume a simple pole form [10] for $A_1(q^2)$, $A_2(q^2)$, and $V(q^2)$, where the pole mass is $M_{D^*(1-)} = 2.01 \text{ GeV}/c^2$ and $M_{D^*(1+)} = 2.42 \text{ GeV}/c^2$ [4] for the vector and axial form factors, respectively. We have also explored a double-pole parametrization [7].

We report herein the first measurement of the form factor ratios and absolute form factor normalization in $D \rightarrow \rho e^+\nu_e$, and improved branching fraction measurements for these decays and $D^+ \rightarrow \omega e^+\nu_e$. (Throughout this Letter charge-conjugate modes are implied.) These decays were studied previously using a smaller CLEO-c data sample [11]. The data sample used here consists of an integrated luminosity of 818 pb^{-1} at the $\psi(3770)$ resonance and includes about 3.0×10^6 $D^0\bar{D}^0$ and 2.4×10^6 D^+D^- events. The CLEO-c detector is described in detail elsewhere [12].

The analysis technique was employed in previous CLEO-c studies [11,13]. The presence of two D mesons in a $D\bar{D}$ event allows a tag sample to be defined in which a \bar{D} is reconstructed in a hadronic decay mode. A subsample is then formed in which a positron and a set of hadrons, as a signature of a semileptonic decay, are required in addition to the tag. The semileptonic decay branching fraction \mathcal{B}_{SL} is given by

$$\mathcal{B}_{\text{SL}} = \frac{N_{\text{tag,SL}}}{N_{\text{tag}}} \frac{\epsilon_{\text{tag}}}{\epsilon_{\text{tag,SL}}} = \frac{N_{\text{tag,SL}}/\epsilon}{N_{\text{tag}}}, \quad (6)$$

where N_{tag} and ϵ_{tag} are the yield and reconstruction efficiency, respectively, for the hadronic tag, $N_{\text{tag,SL}}$ and $\epsilon_{\text{tag,SL}}$ are those for the combined semileptonic decay and hadronic tag, and $\epsilon = \epsilon_{\text{tag,SL}}/\epsilon_{\text{tag}}$ is the effective signal efficiency.

Candidate events are selected by reconstructing a \bar{D}^0 or D^- tag in the following hadronic final states: $K^+\pi^-$, $K^+\pi^-\pi^0$, and $K^+\pi^-\pi^-\pi^+$ for neutral tags, and $K_S^0\pi^-$, $K^+\pi^-\pi^-$, $K_S^0\pi^-\pi^0$, $K^+\pi^-\pi^-\pi^0$, $K_S^0\pi^-\pi^-\pi^+$, and $K^-K^+\pi^-$ for charged tags. Tagged events are selected based on two variables: $\Delta E \equiv E_D - E_{\text{beam}}$, the difference between the energy of the D^- tag candidate E_D and the beam energy E_{beam} , and the beam-constrained mass $M_{\text{bc}} \equiv (E_{\text{beam}}^2/c^4 - |\mathbf{p}_D|^2/c^2)^{1/2}$, where \mathbf{p}_D is the measured momentum of the D^- candidate. Selection criteria for tracks, π^0 , and K_S^0 candidates used in the reconstruction of tags are described in Ref. [14]. If multiple candidates are present in the same tag mode, one candidate per tag charge with the smallest $|\Delta E|$ is chosen. The yield of each tag mode is obtained from fits to the M_{bc} distributions [14]. The data sample comprises 661232 ± 879 and 481927 ± 810 reconstructed neutral and charged tags, respectively.

After a tag is identified, we search for an e^+ and a ρ^- ($\pi^-\pi^0$ mode), ρ^0 ($\pi^+\pi^-$ mode), or ω ($\pi^+\pi^-\pi^0$ mode) recoiling against the tag following Ref. [14]. A $\rho \rightarrow \pi\pi$ candidate satisfies $|m_{\pi\pi} - m_0| < 150 \text{ MeV}/c^2$. The combined tag and semileptonic candidates must account for all tracks in the event. Semileptonic decays are identified with $U \equiv E_{\text{miss}} - c|\mathbf{p}_{\text{miss}}|$, where E_{miss} and \mathbf{p}_{miss} are the missing energy and momentum of the D^+ meson. If the decay products have been correctly identified, U is expected to be zero, since only a neutrino is undetected. The resolution in U is improved by constraining the magnitude and direction of the D^+ momentum to be $p_{D^+} = (E_{\text{beam}}^2/c^2 - c^2m_D^2)^{1/2}$, and

$\hat{\mathbf{p}}_{D^+} = -\hat{\mathbf{p}}_{D^-}$ [11], respectively. Due to the finite resolution of the detector, the distribution in U is approximately Gaussian, with resolution $\sim 17 \text{ MeV}$ for $D^0 \rightarrow \rho^- e^+ \nu_e$ and $D^+ \rightarrow \omega e^+ \nu_e$ and $\sim 8 \text{ MeV}$ for $D^+ \rightarrow \rho^0 e^+ \nu_e$. To remove multiple candidates in each semileptonic mode, which arise due to multiple possible assignments of the particles on the signal side, one combination is chosen per tag mode per tag charge based on the proximity of the invariant masses of the ρ^0 , ρ^+ , or ω candidates to their expected masses.

The U and invariant mass distributions for $D^0 \rightarrow \rho^- e^+ \nu_e$, $D^+ \rightarrow \rho^0 e^+ \nu_e$, and $D^+ \rightarrow \omega e^+ \nu_e$ with all tag modes combined are shown in Fig. 1. The yield for each of the three modes is determined from a binned likelihood fit to the U distribution where the signal is described by a modified Crystal Ball function with two power-law tails [15] which account for initial- and final-state radiation and mismeasured tracks. The signal parameters are fixed with a GEANT-based Monte Carlo (MC) simulation [16] in fits to the data. The background functions are determined by a MC simulation that incorporates all available data on D meson decays, which we refer to as ‘‘generic MC’’ simulation. For $D^0 \rightarrow \rho^- e^+ \nu_e$, the backgrounds arise mostly from $D^0 \rightarrow K^{*-} e^+ \nu_e$, peaking at positive U and modeled with a Gaussian function, and events with misidentified tags, which are accounted for in the fit by a fourth order polynomial. The background to $D^+ \rightarrow \rho^0 e^+ \nu_e$ has its largest contribution from $D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e$, $\bar{K}^{*0} \rightarrow K^- \pi^+$, with the peak at higher U due to charged kaons




FIG. 1 (color online). Fits to the U and hadron invariant mass distributions in data (filled circles with error bars) for (a) and (d) $D^0 \rightarrow \rho^- e^+ \nu_e$, $\rho^- \rightarrow \pi^-\pi^0$; (b) and (e) $D^+ \rightarrow \rho^0 e^+ \nu_e$, $\rho^0 \rightarrow \pi^+\pi^-$; and (c) and (f) $D^+ \rightarrow \omega e^+ \nu_e$, $\omega \rightarrow \pi^+\pi^-\pi^0$. The solid line represents the fit of the sum of the signal function and background function to the data. The dashed line indicates the background contribution. The arrows indicate a $\pm 48 \text{ MeV}$ region around the K_S^0 mass, which has been removed for display.

misidentified as charged pions, and the peak at $U \sim 0$ from either decay-in-flight kaons or interactions with detector material. We categorize the background components according to their shape in U and parametrize the overall background shape using combinations of polynomials and Gaussian functions. The background shape parameters are fixed in fits to the data, while the background normalizations are allowed to float. The signal shapes for the invariant mass distributions of the hadronic system are modeled with a Breit-Wigner function, and the background shapes are modeled with generic MC simulation. The peaking background for $D^+ \rightarrow \rho^0 e^+ \nu_e$ that arises from $D^+ \rightarrow \omega e^+ \nu_e$, $\omega \rightarrow \pi^+ \pi^-$, is subtracted. Due to the tag, backgrounds from the non- $D\bar{D}$ processes $e^+ e^- \rightarrow q\bar{q}$, where q is a u, d , or s quark, $e^+ e^- \rightarrow \tau^+ \tau^-$, and $e^+ e^- \rightarrow \psi(2S)\gamma$, are negligible [13]. The signal yields $N_{\text{tag,SL}}$ are given in Table I.

The second row of Fig. 1 shows the $m_{\pi^-\pi^0}$, $m_{\pi^+\pi^-}$, and $m_{\pi^+\pi^-\pi^0}$ distributions with $|U| < 60$ MeV for the three signal modes. The peaking background at $m_{\pi^-\pi^0} \sim 0.49$ GeV/ c^2 arises from $D^0 \rightarrow K^- e^+ \nu$ with $K^- \rightarrow \pi^-\pi^0$. The small background peak at $m_{\pi^+\pi^-} \sim 0.78$ GeV/ c^2 is due to $D^+ \rightarrow \omega e^+ \nu_e$ with $\omega \rightarrow \pi^+\pi^-$ [17].

The absolute branching fractions in Table I are obtained using Eq. (6). The signal efficiencies ϵ are determined by MC simulation and have been weighted by the tag yields in the data.

The systematic uncertainties for the branching fractions of $D^0 \rightarrow \rho^- e^+ \nu_e$ and $D^+ \rightarrow \rho^0 e^+ \nu_e$ are dominated by uncertainties in the line shape of the ρ (5.0%), and the nonresonant background (−1.5% for $D^0 \rightarrow \rho^- e^+ \nu_e$ and −8.4% for $D^+ \rightarrow \rho^0 e^+ \nu_e$). The uncertainty due to the line shape of the ρ is estimated by (1) requiring $|U| < 60$ MeV and fitting the $m_{\pi\pi}$ distribution, and (2) varying the selection criterion $|m_{\pi\pi} - m_0| < 150$ MeV. The uncertainty due to the nonresonant background is obtained by performing a form factor fit, with an additional interfering nonresonant $D \rightarrow \pi\pi e^+ \nu_e$ (s -wave) component modeled following Ref. [18], then integrating over the kinematic variables to recalculate the branching fractions. The unknown form factors in $D^+ \rightarrow \omega e^+ \nu_e$ are the dominant uncertainty in its branching fraction (3.0%). The remaining systematic uncertainties include the track and π^0 finding efficiencies, positron and charged hadron identification,

the number of tags, selecting tags with the smallest $|\Delta E|$, the no-additional-track requirement, the shape of the signal and background functions, and the MC final-state radiation and form factor modeling. These estimates are added in quadrature to obtain the total systematic uncertainties on the branching fractions: $^{+5.7}_{-5.9}\%$, $^{+5.5}_{-10.0}\%$, 4.1%, for $D^0 \rightarrow \rho^- e^+ \nu_e$, $D^+ \rightarrow \rho^0 e^+ \nu_e$, and $D^+ \rightarrow \omega e^+ \nu_e$, respectively.

A form factor analysis is performed for $D \rightarrow \rho e^+ \nu_e$. We calculate the energy and momentum of the neutrino using $E_\nu = E_{\text{miss}}$ and $|\mathbf{p}_\nu| = E_{\text{miss}}$, because E_{miss} is better measured than $|\mathbf{p}_{\text{miss}}|$. Without ambiguity, the four kinematic variables (q^2 , $\cos\theta_\pi$, $\cos\theta_e$, χ) are measured with resolutions of (0.021 GeV $^2/c^4$, 0.020, 0.048, 0.024) for $D^0 \rightarrow \rho^- e^+ \nu_e$, and (0.013 GeV $^2/c^4$, 0.013, 0.037, 0.019) for $D^+ \rightarrow \rho^0 e^+ \nu_e$.

A four-dimensional maximum likelihood fit in a manner similar to Ref. [19] is performed in the space of q^2 , $\cos\theta_\pi$, $\cos\theta_e$, and χ . The technique makes possible a multidimensional fit to variables modified by experimental acceptance and resolution taking into account correlations among the variables. The signal probability density function for the likelihood function is estimated at each data point using signal MC events by sampling the MC distribution at the reconstructed level in a search volume around the data point, then weighting by the ratio of the decay distribution for the trial values of r_V and r_2 to that of the generated distribution. The search volumes are one tenth the full kinematic range of each of the four dimensions. Large MC samples are generated to ensure that each search volume has sufficient statistics. The background probability density function is modeled using events from the generic MC simulation. Due to the low statistics of the background in the generic MC simulation, we reduce the four-dimensional space to lower dimensional subspaces. Because of the correlation between q^2 and $\cos\theta_e$, the two subspaces are chosen to be (q^2 , $\cos\theta_e$) and ($\cos\theta_\pi$, χ). The background normalization is fixed in the fits to the values measured in the determination of the branching fractions.

Using the above method, a simultaneous fit is made to the isospin-conjugate modes $D^0 \rightarrow \rho^- e^+ \nu_e$ and $D^+ \rightarrow \rho^0 e^+ \nu_e$. We find $r_V = 1.48 \pm 0.15$ and $r_2 = 0.83 \pm 0.11$, with a correlation coefficient $\rho_{V2} = -0.18$. The confidence level of the fit is determined to be 5.0% by comparing the negative log-likelihood from the data to the distribution from MC pseudoexperiments. Figure 2 shows

TABLE I. Signal efficiencies, yields, and branching fractions (\mathcal{B}_{SL}) for $D^0 \rightarrow \rho^- e^+ \nu_e$, $D^+ \rightarrow \rho^0 e^+ \nu_e$, and $D^+ \rightarrow \omega e^+ \nu_e$, from this work, our previous (prev) measurements [11], and three model predictions: QCD sum rules (QCD SR) [5], ISGW2 [6] and FK [7]. All \mathcal{B}_{SL} are in units of 10^{-3} . The uncertainties for ϵ and $N_{\text{tag,SL}}$ are statistical, while the uncertainties for branching fractions are statistical and systematic in that order. The efficiencies include the ρ and ω decay branching fractions from the Particle Data Group (PDG) [4].

Decay Mode	ϵ (%)	$N_{\text{tag,SL}}$	\mathcal{B}_{SL}	\mathcal{B}_{SL} (prev)	\mathcal{B}_{SL} (QCD SR)	\mathcal{B}_{SL} (ISGW2)	\mathcal{B}_{SL} (FK)
$D^0 \rightarrow \rho^- e^+ \nu_e$	26.03 ± 0.02	304.6 ± 20.9	$1.77 \pm 0.12 \pm 0.10$	$1.94 \pm 0.39 \pm 0.13$	0.5 ± 0.1	1.0	2.0
$D^+ \rightarrow \rho^0 e^+ \nu_e$	42.84 ± 0.03	447.4 ± 24.5	$2.17 \pm 0.12^{+0.12}_{-0.22}$	$2.1 \pm 0.4 \pm 0.1$...	1.3	2.5
$D^+ \rightarrow \omega e^+ \nu_e$	14.67 ± 0.03	128.5 ± 12.6	$1.82 \pm 0.18 \pm 0.07$	$1.6^{+0.7}_{-0.6} \pm 0.1$...	1.3	2.5




FIG. 2. Projections of the combined ρ^- and ρ^0 data (points with statistical error bars) and the fit (solid histogram) onto q^2 , $\cos\theta_e$, $\cos\theta_\pi$, and χ . The dashed lines show the sum of the background distributions.

the q^2 , $\cos\theta_e$, $\cos\theta_\pi$, and χ projections for the combined ρ^- and ρ^0 data and the fit. We also make fits to the two modes separately. The results are consistent. We note that the difference between the data and the fit projection for $\cos\theta_\pi$ might be due to s -wave interference.

We have considered the following sources of systematic uncertainty in the form factor measurement. Our estimates of their magnitude are given in parentheses for r_V and r_2 , respectively. The uncertainty associated with background modeling (0.01, 0.02) is estimated by changing the normalization of the three largest background components by a factor of 2 in each semileptonic mode. The uncertainty due to imperfect knowledge of the ρ line shape (0.01, 0.02) is estimated by modifying the ρ line shape by increasing and decreasing the population of signal MC events below and above the nominal ρ mass [4] by 20%. The uncertainty due to a nonresonant background (0.01, 0.02) is obtained by repeating the fit with an additional interfering nonresonant $D \rightarrow \pi\pi e^+ \nu_e$ component (s -wave) following Ref. [18]. The procedure for extracting the form factor parameters is tested using the generic MC sample, from which events are drawn randomly to form mock data samples, each equivalent in size to the data sample. When backgrounds are absent, the measured form factor ratios are consistent with the input values. In the presence of background, a small statistically significant shift is observed. Its magnitude is taken as the uncertainty due to possible bias in the form factor fitter (0.03, 0.02). The

uncertainty associated with the unknown q^2 dependence of the form factors (0.03, 0.02) is estimated by introducing a second pole [7].

Adding all sources of systematic uncertainty in quadrature, the final result is $r_V = 1.48 \pm 0.15 \pm 0.05$ and $r_2 = 0.83 \pm 0.11 \pm 0.04$. Using $|V_{cd}| = 0.2252 \pm 0.0007$ obtained using Cabibbo-Kobayashi-Maskawa unitarity constraints [4] and the lifetimes $\tau_{D^0} = (410.1 \pm 1.5) \times 10^{-15}$ s and $\tau_{D^+} = (1040 \pm 7) \times 10^{-15}$ s [4], we combine our form factor ratio and branching fraction results to obtain $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$, $A_2(0) = 0.47 \pm 0.06 \pm 0.04$, and $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$.

Our branching fraction results are compared to previous measurements [11], with which they are consistent, and theoretical predictions in Table I. The results are consistent with isospin invariance: $\frac{\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e)}{2\Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e)} = 1.03 \pm 0.09^{+0.08}_{-0.02}$. Isospin symmetry is not expected to be exact due to $\rho^0 - \omega$ interference [17]. Theoretical predictions from QCD sum rules [5], the Isgur-Scora-Grinstein-Wise updated (ISGW2) model [6], and a model [Fajfer-Kamenik (FK)] that combines heavy-quark symmetry and properties of the chiral Lagrangian [7], are also listed in Table I. The branching fractions for QCD sum rules and ISGW2 are obtained by combining the partial rates in Refs. [5,6] with $|V_{cd}|$ and τ_D from the PDG [4]. Our branching fraction results are more consistent with the FK predictions than ISGW2.

QCD sum rules predict $A_1(0) = 0.5 \pm 0.2$, $A_2(0) = 0.4 \pm 0.1$, and $V(0) = 1.0 \pm 0.2$. The FK model predicts

$A_1(0) = 0.61$, $A_2(0) = 0.31$, and $V(0) = 1.05$. These values are compatible with our form factor measurements. No other experimental form factor results on these decays exist. Our values of r_V and r_2 are very similar to the current PDG average of $D^+ \rightarrow \bar{K}^{*0} e^+ \nu$ form factor ratios $r_V = 1.62 \pm 0.08$ and $r_2 = 0.83 \pm 0.05$ [4].

In summary, we have made the first measurement of the form factor ratios and absolute form factor normalization in $D \rightarrow \rho e^+ \nu_e$, and improved branching fraction measurements for these decays and $D^+ \rightarrow \omega e^+ \nu_e$. Our branching fractions are consistent with our previous measurements but with improved precision. The form factor measurement in $D \rightarrow \rho e^+ \nu_e$ is the first in a semileptonic Cabibbo-suppressed pseudoscalar-to-vector transition and will aid the determination of $|V_{ub}|$ via the double-ratio technique.

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the A. P. Sloan Foundation, the National Science Foundation, the U.S. Department of Energy, the Natural Sciences and Engineering Research Council of Canada, and the U.K. Science and Technology Facilities Council.

*Present address: University of Science and Technology of China, Hefei 230026, People's Republic of China.

†Present address: University of California, Santa Barbara, CA 93106, USA.

- [1] N. Isgur and M. B. Wise, *Phys. Rev. D* **42**, 2388 (1990).
- [2] B. Grinstein and D. Pirjol, *Phys. Rev. D* **70**, 114005 (2004).
- [3] B. Grinstein, *Phys. Rev. Lett.* **71**, 3067 (1993).
- [4] K. Nakamura *et al.* (Particle Data Group), *J. Phys. G* **37**, 075021 (2010).
- [5] P. Ball, *Phys. Rev. D* **48**, 3190 (1993).
- [6] D. Scora and N. Isgur, *Phys. Rev. D* **52**, 2783 (1995).
- [7] S. Fajfer and J. Kamenik, *Phys. Rev. D* **72**, 034029 (2005).
- [8] J. G. Körner and G. A. Schuler, *Z. Phys. C* **46**, 93 (1990); F. J. Gilman and R. L. Singleton, Jr., *Phys. Rev. D* **41**, 142 (1990).
- [9] J. M. Link *et al.* (FOCUS Collaboration), *Phys. Lett. B* **621**, 72 (2005).
- [10] J. D. Richman and P. Burchat, *Rev. Mod. Phys.* **67**, 893 (1995) and references therein.
- [11] T. E. Coan *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **95**, 181802 (2005); G. S. Huang *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **95**, 181801 (2005).
- [12] Y. Kubota *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **320**, 66 (1992); D. Peterson *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **478**, 142 (2002); M. Artuso *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **554**, 147 (2005).
- [13] D. Besson *et al.* (CLEO Collaboration), *Phys. Rev. D* **80**, 032005 (2009); J. Y. Ge *et al.* (CLEO Collaboration), *Phys. Rev. D* **79**, 052010 (2009); R. E. Mitchell *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **102**, 081801 (2009); J. Yelton *et al.* (CLEO Collaboration), *Phys. Rev. D* **84**, 032001 (2011).
- [14] S. Dobbs *et al.* (CLEO Collaboration), *Phys. Rev. D* **76**, 112001 (2007).
- [15] J. Gaiser, Ph.D. thesis, Stanford University Report No. SLAC-255, 1982; T. Skwarnicki, Ph.D thesis, Jagiellonian University in Krakow DESY Report No. F31-86-02, 1986.
- [16] R. Brun *et al.*, GEANT 3.21, CERN Program Library Long Writup W5013 (unpublished).
- [17] $\rho^0 - \omega$ interference is not included in the ρ^0 line shape but is accounted for both by fitting the U distribution to estimate and subtract the $D^+ \rightarrow \omega e^+ \nu_e$ with $\omega \rightarrow \pi^+ \pi^-$ contribution and by large variations in the ρ^0 (and ρ^+) line shape in the determination of the branching fractions and form factors.
- [18] J. M. Link *et al.* (FOCUS Collaboration), *Phys. Lett. B* **535**, 43 (2002); J. M. Link *et al.* (FOCUS Collaboration), *Phys. Lett. B* **544**, 89 (2002); J. M. Link *et al.* (FOCUS Collaboration), *Phys. Lett. B* **633**, 183 (2006).
- [19] D. M. Schmidt, R. J. Morrison, and M. S. Witherell, *Nucl. Instrum. Methods Phys. Res., Sect. A* **328**, 547 (1993).