

SOME SPACES RELATED TO TOPOLOGICAL INEQUALITIES PROVEN BY THE ERDÖS-RADO THEOREM

WILLIAM G. FLEISSNER

ABSTRACT. The Erdős-Rado theorem is very useful in proving cardinal inequalities in topology. It has been suggested that certain of these inequalities might be strengthened. We note that trees constructed by Jensen and Gregory using various extra axioms of set theory yield several counterexamples to these suggestions; for example, a space X , $|X| = \omega_2$, $c(X) = \omega_1$, $\chi(X) = \omega$, answering a question of Hajnal and Juhász. We consider the apparently similar relation between $|X|$, $e(X)$, and $d\Psi(X)$ of Ginsburg and Woods. Using combinatorial consequences of $V = L$, we construct G_δ tree families, and establish that, assuming $V = L$, an infinite cardinal κ is weakly compact iff $d\Psi(X) < \kappa$, $e_\alpha(X) < \kappa$ imply $|X| < \kappa$.

We consider products of countable chain condition spaces, and show that, using Cohen forcing that 2^ω can be anything allowed by König's theorem and there are spaces X, Y , $c(X) = c(Y) = \omega$, $c(X \times Y) = 2^\omega$. A variation is a space W with the property $c(W^n) = \omega_{n-1}$.

The Erdős-Rado theorem, $(2^\kappa)^+ \rightarrow (\kappa^+)_\alpha^2$, $\alpha < \kappa$, is very useful in establishing inequalities of cardinal functions, as amply illustrated in I. Juhász book. It has been suggested that certain of these inequalities might be sharpened. For example, $|X| \leq 2^{\chi(X)c(X)}$ gives a bound for $|X|$ in terms of the single variable $\max(\chi(X), c(X))$. Can a sharper bound be given as a function of two variables, $\chi(X)$ and $c(X)$? The product of k copies of the two point discrete space demonstrates that $\chi(X)$ must be in the exponent, but no example requiring $c(X)$ to be in the exponent is known. The starting point of this research was the conjecture that $|X| \leq c(X)^{\chi(X)}$.

In A-E below, we list five inequalities proven by the Erdős-Rado theorem. A, B, and C appear in [Ju], D in [GW₁] and E in [GW₂]. We also list how certain spaces, whose existence is known to be consistent with the usual axioms of set theory, refute conjectured improvements. B is a first countable ω_2 -Souslin line, L is the order compactification of B , S is the disjoint union of ω copies of L , and A is the antichain in an ω_2 , ω G_δ tree family.

- A. $|X| \geq 2^{\chi(X)c(X)}$ for T_2 spaces X : our space B refutes $|X| \leq c(X)^{\chi(X)}$.
- B. $|X| \leq 2^{\psi(X)s(X)}$ for T_2 spaces X : our space B refutes $|X| \leq s(X)^{\psi(X)}$.
- C. $c_\omega(X) \leq 2^{c(X)}$ for compact T_2 spaces X ; our space L refutes $c_\omega(X) \leq c(X)^\omega$.

Received by the editors March 12, 1977 and, in revised form, December 16, 1977.

AMS (MOS) subject classifications (1970). Primary 54A25, 04A20.

Key words and phrases. Cardinal functions in topology, trees, $V = L$, Erdős-Rado theorem, cellularity, character, spread, G_δ diagonal, products of ccc spaces, Cohen forcing.

© American Mathematical Society 1978

D. $c(\beta X - X) \leq 2^{K(X)c(X)}$ for completely regular spaces X ; our space S refutes $c(\beta X - X) \leq c(X)^{K(X)}$.

E. $|X| \leq 2^{d\psi(X)e(X)}$ for T_1 spaces X ; our space A refutes $|X| \leq e(X)^{d\psi(X)}$.

The plan of this paper is as follows. In §1 we introduce the notation and terminology used throughout this paper. We define convenient trees and use them to construct the spaces B, L and S . Next we define G_δ tree families and use them to define the space A . In §3, we construct convenient trees from Souslin trees and construct G tree families assuming $V = L$. Then we discuss a sup = max problem related to inequality E and contrast it with the analogous problems A–D. In §5, we discuss the cellularity of the product of two ccc spaces. The Erdős-Rado theorem gives an upper bound of c and we show that this upper bound can be attained no matter what the value of c .

1. We denote the cardinality of X by $|X|$. We use $[\alpha]^\beta$ to denote the set of subsets of α of cardinality β : $[\alpha]^{<\omega}$ is the set of finite subsets of α . The partition relation $\alpha \rightarrow (\beta)_\gamma^n$ means that whenever $P: [\alpha]^n \rightarrow \gamma$ there is $\delta \in \gamma$ and $H \in [\alpha]^\beta$ with $P''[H]^n = \{\delta\}$.

For a topological space X , with topology $\sigma(X)$, we define

cellularity	$c(X) = \omega \cdot \sup\{ \mathcal{U} : \mathcal{U} \subset \sigma(X), \mathcal{U} \text{ is disjoint}\};$
G_δ cellularity	$c_\omega(X) = \omega \cdot \sup\{ \mathcal{U} : U \in \mathcal{U} \text{ is } G_\delta, \mathcal{U} \text{ is disjoint}\};$
character	$\chi(X) = \sup\{\min\{ \mathcal{U}_p : \mathcal{U}_p \text{ is a nghd basis at } p\}: p \in X\};$
pseudocharacter	$\psi(X) = \sup\{\min\{ \mathcal{U}_p : \bigcap \mathcal{U}_p = \{p\}\}: p \in X\};$
spread	$s(X) = \omega \cdot \sup\{ D : D \subset X, D \text{ is discrete}\};$
extent	$e(X) = \omega \cdot \sup\{ D : D \subset X, D \text{ is closed, discrete}\};$
G_δ degree	$d\psi(X) = \min\{ \mathcal{U} : \mathcal{U} \subset \sigma(X^2), \bigcap \mathcal{U} = \text{diag } X\};$
K coweight	$K(X) = \min\{ \mathcal{K} : K \in \mathcal{K} \text{ is cpt, } \forall H \subset X,$

$$H \text{ cpt } \exists K \in \mathcal{K}, H \subset K\}.$$

We define (the phrase) $c_\alpha(X) < \kappa$ to mean that there is no set of cardinality κ of disjoint open sets of X . We similarly define $e_\alpha(X) < \kappa$ and $s_\alpha(X) < \kappa$. For a completely regular space X , βX is the Stone-Cech compactification. X is a ccc space iff $c(X) = \omega$.

A tree is a partially ordered set T , such that for every $x \in T$, the set $\hat{x} = \{y \in T: y < x\}$ is well ordered. Set $\bar{x} = \{x\} \cup \hat{x}$. The order $o(x)$ of x is the order type of \hat{x} . The α th level of T is $T_\alpha = \{x \in T: o(x) = \alpha\}$. The restriction of T to height α is

$$T|\alpha = \{x \in T: o(x) < \alpha\}.$$

A branch of T is a maximal linearly ordered subset of T ; an α -branch is a branch of order type α . We say an α -branch b of $T|\alpha$ continues if there is $x \in T_\alpha, \hat{x} = b$. An antichain of T is a set of incomparable elements of T ; we

abbreviate maximal antichain by mac. A λ -Souslin tree is a tree with elements of order α for $\alpha < \lambda$, but no branches or antichains of cardinality λ .

2. Problem 77/A of [EH] is whether $c(X) = \omega_1$ implies $|X| = \omega_1$ for first countable linearly ordered spaces X , assuming the Continuum Hypothesis. This is a special case of the question of whether inequality A can be sharpened. We begin this section by reviewing Jensen's solution to this problem.

Let us call a tree T λ, μ convenient if

- (i) every antichain of T has cardinality less than λ ;
- (ii) every element of T has an infinite number of immediate successors;
- (iii) there are at least λ branches of T of cofinality μ .

Let T be a c^+, ω convenient tree; let $B = \{b_\alpha : \alpha < c^+\}$ be a family of branches of cofinality ω ; let $\{x_\alpha^n : n < \omega\}$ be $<_T$ increasing and cofinal in b_α .

We obtain our spaces by "squashing" T to a line [Kp₁], [Ru]. Let $<_0$ be a linear order on T . Because of (ii) we may choose $<_0$ so that for all $x \in T$, the set of immediate successors of x has neither a $<_0$ greatest nor a $<_0$ least element. We define a linear order $<_1$ on $\text{Br}(T)$, the set of all branches of T , by

$$b <_1 b' \text{ iff } \exists \alpha \ b|\alpha = b'|\alpha \text{ and } b(\alpha) <_0 b'(\alpha).$$

We give $\text{Br}(T)$ the topology induced by the order $<_1$. Let $B (= \{b_\alpha : \alpha < c^+\})$ have the subspace topology. Let L be the order completion of $\text{Br}(T)$ with endpoints. By the Heine-Borel theorem, L is compact. Then $L \times \omega$, (i.e. countably many disjoint copies of L) is σ -compact.

For $x \in T$, define $(x) = \{b \in \text{Br}(T) : x \in b\}$. The point of (ii) and our choice of $<_0$ is that (x) is open in $\text{Br}(T)$, and in L , $(x) \subset \text{Interior}(\text{closure}(x))$. The point of (i) is that $c(\text{Br}(T)) < c^+$. The point of (iii) is that $\{(x_\alpha^n) : n < \omega\}$ is neighborhood base for b_α .

Let us determine the relevant cardinal functions for B, L , and S . The families $\{(x_\alpha^n) : n < \omega\}$ demonstrate that $\chi(X) = \omega = \psi(X)$. A nonempty subset of B contains some (x) , so by (i) $c(B) < c^+$. Because B is a linearly ordered topological space, $s(B) = c(B)$. Because B is dense in L , $c(L) = c(B)$, so $c(L) < \omega_2$. Each b_α is a G_δ in L , so $c_\omega(L) \geq c^+$. Let $S = \omega \times L$. Every compact subset of S is contained in some $m \times L$, $m < \omega$, so $K(S) = \omega$. Next $c(S) = c(L) \cdot \omega < c^+$. We define open subsets U_α , $\alpha < c^+$, of S : $U_\alpha = \{\{n\} \times (x_\alpha^n) : n < \omega\}$. Then $\{(\text{int}_{\beta S} \text{cl}_{\beta S} U_\alpha) - S : \alpha < c^+\}$ is a disjoint family of nonempty open subsets of $\beta S - S$.

The existence of a space such as A of E does not seem (at least to the author) to follow from the existence of a c^+, ω convenient tree. The results of §4 also tend to indicate that E is fundamentally different from A-D. So we define G_δ tree families. For λ, μ regular, $\mu < \lambda$, a family $\mathfrak{T} = \{T^\alpha : \alpha \leq \mu\}$ is a λ, μ G_δ tree family iff

- (i) $\forall \alpha \leq \mu \ T^\alpha \subset {}^\alpha \lambda$;
- (ii) $\forall \alpha < \mu \ T^\alpha$ is a λ Souslin tree, T^μ is not;
- (iii) $\forall \beta < \alpha < \mu \ \forall \sigma, \tau \in T^\alpha \ \sigma <_{T^\alpha} \tau$ implies $\sigma|\beta <_{T^\beta} \tau|\beta$;

(iv) $x, y \in T^\mu X <_{T^\mu} y$ iff $\forall \alpha < \mu \ x|\alpha <_{T^\alpha} y|\alpha$.

We will construct the space A from a c^+ , ω G_δ tree family \mathcal{T} . Let A be an antichain of T^ω of cardinality c^+ . The n th basic open neighborhood of a is $B_n(a) = \{a' \in A : a'|n <_n a|n\}$. We note $e(A) \leq s(A) \leq c$, because c^+ is regular and each T^n is a c^+ Souslin tree. A has a G_δ diagonal because for all $a \in A$

$$\bigcap_{n < \omega} \{a' \in A : a'|n \leq a|n \text{ or } a|n \leq a'|n\} = \{a\}.$$

3. In the previous section we constructed spaces from convenient trees and G_δ tree families. In this section we construct convenient trees and G_δ tree families.

Jensen [J] and Gregory [G] constructed ω_2 -Souslin trees from combinatorial principles. These trees are ω_2 , ω convenient trees as well. Not every ω_2 -Souslin tree is an ω_2 , ω convenient tree; using CH and \diamond_{ω_1} , ω_2 one can construct an ω_2 -Souslin tree “pruning” only at levels of cofinality ω_1 . However, we can construct a convenient tree from a Souslin tree.

THEOREM. *If there is a λ -Souslin tree S , then there is a λ , ω convenient tree T .*

PROOF. As in the previous section, we may “squash” S to an order complete line L_S . Because L_S is order complete and dense, whenever $a < b$ there is c , $a < c \leq b$, such that (a, c) has cofinality ω .

Using this fact we will construct a new tree T from L_S . Elements of T will be intervals (a, b) ; the order will be given by $(a, b) < (a', b')$ iff $(a, b) \supset (a', b')$. By induction on $\alpha < \lambda$, we construct T_α , the α th level of T to satisfy the following conditions

0. $T_0 = L_S$;
1. $\cup T_\alpha$ is dense in L_S ;
2. every element of $T_{\alpha+1}$ is a proper subset of an element of T_α and has cofinality ω ;
3. every element of T_α contains infinitely many elements of $T_{\alpha+1}$;
4. if $(a, c) \in T_\alpha$ and has cofinality ω , then for some a' $(a', c) \in T_{\alpha+1}$;
5. for δ a limit ordinal T_δ is the set of intersections $\cap \{t_\beta : \beta < \delta\}$, where $t_\beta \in T_\beta$.

We verify that T satisfies the three conditions to be a convenient tree. Because S was a λ -Souslin tree, T is, too, so T satisfies (i). Condition 3 insures that T satisfies (ii). By 4 and 5, if $(a, c) \in T$ and has cofinality ω , then the set of elements of T with c as endpoint determine a branch of cofinality ω . By 2, there are λ such branches.

We now construct a λ , μ G_δ tree family assuming $V = L$. There are two cases: λ is strongly inaccessible and $\lambda = \kappa^+$. If λ is strongly inaccessible, we use $ES\lambda$: There are $E \subset \lambda$, $S = \{S_\alpha : \alpha \in E\}$ satisfying

- (i) E is stationary in λ , $\forall \alpha < \lambda$ there is K_α closed unbounded in α , $K_\alpha \cap E = \emptyset$, and $\beta \in E$ implies $cf\beta = \mu$,

(ii) $\forall X \subset \lambda, \{\alpha: S_\alpha = X \cap \alpha\}$ is stationary.

If $\lambda = \kappa^+$, we use \square_κ^* , which is $ES\kappa^+$ plus the existence of $\{C_\beta: \beta < \lambda, \beta$ a limit ordinal $\}$ satisfying

(iii) C_β is closed and unbounded in β ,

(iv) if γ is a limit point of C_β then $\gamma \notin E$ and $C_\gamma = C_\beta \cap \gamma$. Further, if $\mu > \omega$, we use $\kappa < \lambda, \nu < \mu$ implies $\lambda > \kappa^\nu$.

We will construct $T = T^\mu$ by constructing $T|\alpha$ and $A = \{a_\beta: \beta \in E\}$, the future large antichain, by induction on $\alpha < \lambda$. As we will use S to fix mac's in each $T^\nu, \nu < \mu$, we assume, via a bijection between λ and $\cup \{\nu\lambda: \nu < \mu\}$, that (ii) holds for $X \subset \cup \{\nu\lambda: \nu < \mu\}$. Our induction hypothesis on α will be

(1) $|T|\alpha| < \lambda, \forall \nu \leq \mu T^\nu \subset \nu\lambda$;

(2) $\forall \beta < \gamma < \alpha, \beta, \gamma \in E, o(a_\beta) = \beta, o(a_\gamma) = \gamma, a_\beta \not\leq a_\gamma$;

(3) $\forall \beta < \alpha \forall \nu < \mu S_\beta$ is a mac in $T^\nu|\beta$ implies S_β is a mac in $T^\nu|\alpha$;

(4) $\forall \beta < \gamma < \alpha \forall x \in T_\beta \exists y \in T_\gamma x < y$;

(5) $\forall \beta < \gamma < \alpha \forall \nu < \mu \forall \sigma \in T_\gamma \forall x \in T_\beta (x|\nu < \sigma \text{ and } \bar{x} \cap A = \emptyset)$ implies that (there are (at least) two distinct $y, z \in T_\gamma x < y, x < z, (\bar{y} \cup \bar{z}) \cap A = \emptyset, \sigma = y|\nu = z|\nu$);

(6) (i) (if λ is inaccessible) $\forall \beta + 1 < \alpha, \beta \notin E$ every β -branch of $T|\beta$ continues;

(ii) (if $\lambda = \kappa^+$) $\forall \beta + 1 < \alpha, \beta$ a limit ordinal $\beta \notin E$, if cf $\beta < \mu$ every β branch of $T|\beta$ continues, if cf $\beta \geq \mu$ the β branch $b(x, \beta)$ continues, as defined below.

We define $b(x, \beta)$ by defining a set of y 's cofinal in it. Let $<'$ well order ${}^\mu\lambda$. Let $\{\gamma(\delta): \delta < \eta\}$ be the monotone enumeration of C_β (from \square_κ^*). Let $\bar{\delta}$ be least such that $x \in T_{\gamma(\bar{\delta})}$. Inductively define $y_\delta, \bar{\delta} \leq \delta < \eta$:

y_δ is the $<'$ least $y \in T_{\gamma(\bar{\delta})}$ such that $x < y$ and $\bar{y} \cap A = \emptyset$ if $\bar{x} \cap A = \emptyset$.

$y_{\delta+1}$ is the $<'$ least $y \in T_{\gamma(\gamma+1)}$ such that $x < y$ and $\bar{y} \cap A = \emptyset$ if $\bar{x} \cap A = \emptyset$.

y_δ for limit δ is the unique $y \in T_{\gamma(\bar{\delta})}$ such that $\bar{\delta} \leq \delta' < \delta$ implies $y_{\delta'} < y_\delta$. (y_δ exists by (iv) and induction hypothesis.) This completes the definition of $b(x, \beta)$ and the statement of the induction hypothesis.

We construct $T|\alpha$ by induction on α .

For $\alpha = 1$ let T_1 be the constant function 0. For α a limit ordinal we must set $T|\alpha = \cup_{\beta < \alpha} T_\beta$. For $\alpha = \beta + 1, \beta = \gamma + 1$, give every element of T_β two immediate successors and add more elements as required by (5).

If $\alpha = \beta + 1, \beta$ a limit ordinal $\beta \notin E$, continue the branches as required by (6). If $\beta \in E$, continue only enough branches to satisfy (4) and (5). If S_β is a mac in $T^\nu|\beta$, we require that these branches contain some $x, x|\nu \in S_\beta$. Then choose an element $a_\beta \in T_\beta$ to satisfy (2).

We verify that in this last case the construction can be done to satisfy the induction hypothesis, the other cases being simpler. By (i) let $\{\gamma(\delta): \delta < \eta\}$ be a monotone enumeration of a set K_β closed, cofinal in $\beta, K_\beta \cap E = \emptyset$. Since S_β is a mac in $T^\nu|\beta$ for every $x \in T|\beta$ there is $\sigma \in S_\beta, \sigma \leq x|\nu$ or $x|\nu < \sigma$. In the latter case by (5) there is $x', x < x', x'|\nu = \sigma$ and $\bar{x}' \cap A = \emptyset$ if $\bar{x} \cap A = \emptyset$. Call such an x correct. For $x \in T|\beta, x$ correct, let $\bar{\delta}$ be least

such that $x \in T_{\gamma(\bar{\delta})}$. We may inductively choose y_δ , $\bar{\delta} \leq \delta < \eta$, $y_\delta \in T_{\gamma(\delta)}$, $y_\delta > x$, using clause 4 of the induction hypothesis if δ is a successor, clause 6 if δ is a limit. Thus clause 4 of the induction hypothesis holds at α . Clause 5 at α may be verified with the same technique. Having verified clause 5 at α , we may use it to choose a_β to satisfy clause 2. Clause 5 continues to hold because we required two distinct elements for each x, σ .

By (2) T has an antichain A of cardinality λ . By (3) and the usual \diamond argument, each T^ν , $\nu < \mu$, is a λ Souslin tree. This completes the construction of a $\lambda, \mu G_\delta$ tree family.

4. There are sup = max-like questions associated with the inequalities A–E. For example, what is the class $\mathcal{C}(\chi, c)$ of cardinal satisfying “ $\chi(X) < \kappa$, $c_\alpha(X) < \kappa$ implies $|X| < \kappa$ for Hausdorff X ?”

(Associated with inequality C is the class $\mathcal{C}(\cdot, c)$ of cardinals κ such that whenever X is a compact T_2 space with $c_\alpha(X) < \kappa$ then every disjoint family of intersections of $\nu < \kappa$ open subsets of X has cardinality less than κ .)

- THEOREM: (a) κ weakly compact implies $\kappa \in \mathcal{C}(\chi, c) \cap C(d\psi, e)$;
 (b) $\lambda < \kappa$, $2^\lambda \geq \kappa$ implies $\kappa \notin \mathcal{C}(\chi, c) \cup \mathcal{C}(d\psi, e)$;
 (c) κ singular strong limit implies $\kappa \in \mathcal{C}(\chi, c) - \mathcal{C}(d\psi, e)$;
 (d) $(V = L)$ κ inaccessible, not weakly compact implies $\kappa \notin \mathcal{C}(\chi, c) \cup \mathcal{C}(d\psi, e)$;
 (e) (a)–(d) hold with $C(\psi, s)$, $\mathcal{C}(\cdot, c)$, or $\mathcal{C}(\kappa, c)$ in place of $\mathcal{C}(\chi, c)$.

PROOF. (a) $\kappa \rightarrow (\kappa)_\alpha^2$, $\alpha < \kappa$.

(b) Consider 2^λ with the usual product topology.

(c) $\kappa \in \mathcal{C}(\chi, c)$ by [Ju, Chapter 3]. The space $\kappa + 1$ where points α , $\alpha < \kappa$ are isolated and the point κ has the neighborhoods from the order topology shows $\kappa \notin \mathcal{C}(d\psi, e)$.

(d) Consider convenient trees and G_δ tree families.

(e) As above. To show (b) for $\mathcal{C}(K, c)$, let λ be least such that $2^\lambda \geq \kappa$. Consider $\bigcup_{\alpha < \lambda} {}^\alpha 2$ with the “tree” topology.

5. The question of whether the product of ccc spaces is ccc is an old question, but most of the results are recent. Kurepa [Kp₁] showed that the product of two Souslin lines has cellularity ω_1 , and that the cellularity of a product of two (in fact, any family of) ccc spaces has cellularity at most c , [Kp₂]. This latter result can be proven by the Erdős-Rado theorem.

An interesting consequence of $MA + \neg CH$ is that the product of ccc spaces is ccc. Galvin [Ga] showed that CH allows the construction of two ccc spaces whose product is not ccc. Roitman [Ro] constructs a space similar to Galvin’s in models obtained by adding a Cohen or random real. In this context Ketonen’s question of whether $c(X) = \omega = c(Y)$ implies $c(X \times Y) \leq \omega_1$ is natural.

In this section we show that adding κ Cohen reals gives ccc spaces X and Y with $c(X \times Y) \geq \kappa$. We relabel the forcing conditions so that Galvin-type spaces are constructed directly from the generic filter.

Let $P: [\kappa]^2 \rightarrow \{0, 1\}$. Let $X(Y)$ be the set of all $H \subset \kappa$ maximal with respect to $P''[H]^2 \subset \{0\}$ ($P''[H]^2 \subset \{1\}$). For $a \in [\kappa]^{<\omega}$, define $B_X a = \{H \in X: a \subset H\}$ and $B_Y a = \{H \in Y: a \subset H\}$. Give $X(Y)$ the topology generated by $\{B_X a: a \in [\kappa]^{<\omega}\}$ ($\{B_Y a: a \in [\kappa]^{<\omega}\}$). It is easy to see that $c(X \times Y) \geq \kappa$, for $\mathcal{U} = \{B_X\{\alpha\} \times B_Y\{\alpha\}: \alpha < \kappa\}$ is a family of disjoint nonempty open sets. $B_X\{\alpha\} \times B_Y\{\alpha\} \neq \emptyset$ because $\{\{\alpha\}\}^2 = \emptyset$, hence $P''[\{\alpha\}]^2 = \emptyset \subset \{0\} \cap \{1\}$. For $\alpha \neq \beta$, $P(\alpha, \beta) = 0$, say, and then $B_Y\{\alpha\} \cap B_Y\{\beta\} = \emptyset$, so \mathcal{U} is a disjoint family. The problem, thus, is to define P so that X and Y are ccc. We accomplish this by defining P generically.

Fix κ . Let $\mathcal{P} = \{p: |p| < \omega, p \text{ is a function, } \text{dom } p \subset [\kappa]^2, \text{ range } p \subset \{0, 1\}\}$. For $p \in \mathcal{P}$, define $\text{supp } p$ to be the smallest $a \subset \kappa$ such that $\text{dom } p \subset [a]^2$. Because later we will use fancier conditions we note that the proof below uses only that \mathcal{P} is ccc and that $\text{supp } p$ is finite.

Let M be a countable transitive model of set theory, let G be an M -generic ultrafilter on \mathcal{P} . Define $P: [\kappa]^2 \rightarrow \{0, 1\}$ by $P = \bigcup G$. Then if X, Y are defined as above $c(X \times Y) \geq \kappa$. We show X is ccc; of course, a parallel proof shows Y is ccc.

If X is not ccc then there is a family of ω_1 disjoint nonempty open sets $B_X a$. We may apply the Δ system lemma to the a 's and, because of the definition of X , throw away the common part. So assume that $p_0 \in \mathcal{P}$, and

$$p_0 \Vdash A = \{a_\nu: \nu < \omega_1\}, \quad \nu < \nu' < \omega_1 \rightarrow a_\nu \cap a_{\nu'} = \emptyset, \\ B_X a_\nu \cap B_X a_{\nu'} = \emptyset. \tag{*}$$

For $n < \omega$, let $\mathcal{Q}_n \subset \mathcal{P}$ be maximal with respect to

- (a) $p \in \mathcal{Q}_n \rightarrow \exists e(p, n) \in M, p \Vdash a_n = e(p, n)$,
- (b) elements of \mathcal{Q}_n are incompatible.

Let $F = \bigcup_{n < \omega, p \in \mathcal{Q}_n} e(p, n) \cup \text{supp } p$. Because \mathcal{P} is ccc, $|\mathcal{Q}_n| = \omega$, hence $|F| = \omega$. Note that $F \in M$. By (*) there is $p_1 < p_0, \nu < \omega_1, e_1 \in M$

$$p_1 \Vdash a_\nu = e_1, \quad a_\nu \cap F = \emptyset.$$

Now because $|\text{supp } p_1| < \omega$ we can extend p_1 , perhaps $|\text{supp } p_1| + 1$ times, to get $p_2 < p_1, n \in \omega, e_2 \in M$, so that $\text{dom } p \subset [F]^2 \cup \text{dom } p_1$ and

$$p_2 \Vdash a_n = e_2, \quad a_n \cap \text{supp } p_1 = \emptyset.$$

Let $\text{dom } p_3 = \{(\alpha, \beta): \alpha \in e_1, \beta \in e_2\}$, $\text{range } p_3 = \{0\}$. Then

$$p_2 \cup p_3 \Vdash B_X\{a_\nu\} \cap B_X\{a_n\} \neq \emptyset$$

which contradicts (*), and establishes that X is ccc.

This sort of generic construction is quite flexible. For example, we can change \mathcal{P} so that the range of $\bigcup G$ is $\{m: m < n\}$. Set X_i to be the set of $H \subset \kappa$ maximal with respect to $\bigcup G''[H]^2 \subset \{m: m < n\} - \{i\}$. Then the product of any $n - 1$ X_i 's is ccc, proved as above, noting that the common value they all can use gives us room to work in. On the other hand, the open sets $\prod_{i < n} B_{X_i}(\alpha), \alpha < \kappa$, show that $c(\prod_{i < n} X_i) = \kappa$. By similar fiddling around we can construct a space W such that $c(W^n) = \omega_{n-1}$.

BIBLIOGRAPHY

- [De] K. Devlin, *Aspects of constructibility*, Lecture Notes in Math., vol. 354, Springer-Verlag, Berlin and New York, 1973.
- [EH] P. Erdős and A. Hajnal, *Unsolved problems in set theory*, Proc. Sympos. Pure Math., vol. 13, Amer. Math. Soc., Providence, R.I., 1971, pp. 17–48.
- [Ga] F. Galvin, *Chain conditions and products*, Fund. Math. (to appear).
- [GW₁] J. Ginsburg and R. G. Woods, *On the cellularity of $\beta X - X$* , Proc. Amer. Math. Soc. **57** (1976), 151–154.
- [GW₂] ———, *A cardinal inequality for topological spaces involving closed discrete sets*, Proc. Amer. Math. Soc. **64** (1977), 357–360.
- [Gr] J. Gregory, *Higher Souslin trees and the generalized continuum hypothesis*, J. Symbolic Logic **41** (1976), 663–671.
- [Je] R. Jensen, *The fine structure of the constructible hierarchy*, Ann. Math. Logic **4** (1972), 229–308.
- [Ju] I. Juhász, *Cardinal functions in topology*, Math. Centrum, Amsterdam, 1971.
- [Kp₁] D. Kurepa, *Ensembles linéaires et une classe de tableaux ramifiés*, Publ. Math. Univ. Belgrade **6** (1936), 129–160.
- [Kp₂] ———, *The Cartesian multiplication and cellularity numbers*, Publ. Inst. Math. (Beograd) (NS) **2** (1962), 121–139.
- [Ro] J. Roitman, *Adding a Cohen or random real: Topological consequences and effect on Martin's axiom*, Fund. Math. (to appear).
- [Ru] M. E. Rudin, *Souslin's conjecture*, Amer. Math. Monthly **76** (1969), 1113–1119.
- [Sh] B. Sapirovskiĭ, *Canonical sets and character. Density and weight in compact spaces*, Soviet Math. Dokl. **15** (1974), 1282–1287.

INSTITUTE FOR MEDICINE AND MATHEMATICS, OHIO UNIVERSITY, ATHENS, OHIO 45701