

Present and past ecological niche models for the Great Tit (*Parus major*)

Fernando Machado-Stredel¹, Gang Song², Ruiying Zhang², Per Alström^{2,3}, Yanhua Qu², Huijie Qiao², Herman Mays⁴, Per G. Ericson⁵, Jon Fjeldså⁶, Fumin Lei², A. Townsend Peterson¹



1) Biodiversity Institute and Department of Ecology and Evolutionary Biology, University of Kansas
 2) Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences
 3) Department of Ecology and Genetics, Animal Ecology, Evolutionary Biology Centre, Uppsala University
 4) Department of Biological Sciences, Marshall University 5) Department of Bioinformatics and Genetics, Swedish Museum of Natural History
 6) Center for Macroecology, Evolution and Climate at the Natural History Museum of Denmark, University of Copenhagen

The Great Tit (*Parus major*) is a charismatic bird found in most of Eurasia. This widespread species is well known by birdwatchers and has been the subject of a plethora of research studies; however, its evolutionary history and systematics are still in debate. There are taxonomic authorities that recognize four species (Clements Checklist), while others suggest that the whole complex is formed by one taxon with 44 subspecies (HBW).

To understand some of the differentiation patterns recovered in a newly well-resolved multi-locus phylogeny (Song et al *in prep.*), we undertook an ecological niche modeling approach, exploring climatic changes since the Pleistocene along the distribution of the whole complex.





Fig 1. Distribution of *P. major* genotyped populations and calibration area for the models (dark gray contour).

METHODS



RESULTS

The best models ($n = 13$) were significant and showed omission rates between 5.0 - 6.8%, generally using a small set of PCs. Most models (11) produced similar predictions; with northern areas being somewhat excluded from the potential distribution. The additional models were broader and consistent among each other, predicting most of Eurasia as suitable for the species, excluding the Arabic Peninsula, southeastern India, northeastern Siberia, and in some cases the Tibetan Plateau (Fig. 2). Finally, models showed greater environmental continuity and larger suitable areas in the LGM (Fig. 3). Most of the suitability loss occurred in the southern range of the study area, contrasting with northwestern regions in Eurasia.




Fig 2. Ecological niche model for *P. major* with test and train records (q & p features, $\beta = 1$, 6 PCs, 6.5% omission).




Fig 3. Climatic turnover from LGM to present along Eurasia and Africa.

CONCLUSIONS

- Climate change since the LGM might have influenced the current distributional patterns and the diversification of *Parus major*.
- At present, there is less suitable areas and a complex landscape.
- We aim to do new models with more samples from Siberia, considering also other past climatic scenarios such as the Last Interglacial.

REFERENCES

- 1.- del Hoyo, J et al. 2018. Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. (<http://www.hbw.com/> on [12/11/2018]).
- 2.- ESRI 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.
- 3.- Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones and A. Jarvis, 2005. Very high resolution interpolated climate surfaces for global land areas. International Journal of Climatology 25: 1965-1978.
- 4.- RStudio Team (2016). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/> [packages: ENMGadgets & kuenm].