

## The depressogenic potential of added dietary sugars

Daniel J. Reis<sup>\*,1,3</sup>, Stephen S. Ilardi<sup>1</sup>, Michael S. Namekata<sup>1</sup>, Erik K. Wing<sup>1</sup>, Carina H. Fowler<sup>1,2</sup>

University of Kansas, United States



### ARTICLE INFO

#### Keywords:

Depression  
Sugar  
Pathophysiology  
Inflammation

### ABSTRACT

Added sugars are ubiquitous in contemporary Western diets. Although excessive sugar consumption is now robustly associated with an array of adverse health consequences, comparatively little research has thus far addressed its impact on the risk of mental illness. But ample evidence suggests that high-dose sugar intake can perturb numerous metabolic, inflammatory, and neurobiological processes. Many such effects are of particular relevance to the onset and maintenance of depressive illness, among them: systemic inflammation, gut microbiota disruption, perturbed dopaminergic reward signaling, insulin resistance, oxidative stress, and the generation of toxic *advanced glycation end-products* (AGEs). Accordingly, we hypothesize that added dietary sugars carry the potential to increase vulnerability to major depressive disorder, particularly at high levels of consumption. The present paper: (a) summarizes the existing experimental and epidemiological research regarding sugar consumption and depression vulnerability; (b) examines the impact of sugar ingestion on known depressogenic physiological processes; and (c) outlines the clinical and theoretical implications of the apparent sugar-depression link. We conclude that the extant literature supports the hypothesized depressogenic impact of added dietary sugars, and propose that an improved understanding of the effects of sugar on body and mind may aid in the development of novel therapeutic and preventative measures for depression.

### Introduction

Over the past century, sugar has become a dietary staple throughout the developed world. Americans currently derive an estimated 14% of all calories from *added sugars*<sup>a</sup> [1]—typically introduced to foods and beverages during their preparation and processing—the equivalent of 18 teaspoons' worth each day. These sweeteners are even found in an estimated 75% of all packaged foods [2]. The consumption of a high-sugar diet has, of course, been implicated as a risk factor for an array of adverse health outcomes, including obesity [3], cardiovascular disease [4], type 2 diabetes mellitus [5], and dental caries [6]. As a result, U.S. dietary guidelines now advise limiting the consumption of added sugars, both to promote better overall health and to help reduce the burgeoning toll of obesity [7].

Although maladaptive dietary habits pose an obvious threat to physical well-being, they also carry the potential to endanger psychological health [8]. In fact, major depressive disorder, one of the most

highly prevalent and disabling forms of mental illness worldwide [9], appears particularly susceptible to unhealthy nutritional influences [10,11]. Dietary interventions, such as those designed to improve nutrition or reduce weight, have been found to reduce depressive symptoms [12]. Similarly, diets high in the consumption of fruits, vegetables, fish, and whole grains are associated with a lower risk of depression onset [13,14], whereas the frequent intake of red meats, refined grains, sweets, and other “unhealthy” foods is linked with increased depression vulnerability [13,15]. Short-term carbohydrate consumption even carries the potential to induce fatigue and decreased alertness [16]. Moreover, healthy and unhealthy diets, respectively, each appear to exert independent effects on mental health [11,17], which suggests that depressogenic processes can be affected both by the relative absence of key nutrients *and* by the excessive presence of harmful foods. Identifying the latter—the specific dietary components that contribute to depression vulnerability—will enhance our understanding of the pathology and treatment of this devastating illness.

\* Corresponding author at: University of Kansas, Department of Psychology, 1415 Jayhawk Blvd., Fraser Hall Room 426, Lawrence, KS 66045, United States.  
E-mail address: [daniel.reis@ku.edu](mailto:daniel.reis@ku.edu) (D.J. Reis).

<sup>1</sup> Department of Psychology, University of Kansas, United States.

<sup>2</sup> Department of Psychology and Neuroscience, Duke University, United States.

<sup>3</sup> ORCID: 0000-0001-7806-3819.

<sup>a</sup> In contrast with the natural sugars contained in whole foods such as fruits, vegetables and dairy products, *added sugars* subsume the large array of sweeteners—fructose, sucrose, maltose, dextrose, high-fructose corn syrup, cane syrup, cane juice, fruit juice concentrate, etc.—commonly added to foods and beverages during processing.

Therefore, given the established link between excessive sugar intake and adverse health outcomes, as well as the broader association between dietary quality and mental health, we hypothesize that added dietary sugars constitute a risk factor for the onset of depression, particularly at high levels of consumption. In the following sections we examine two primary lines of evidence that lend support to the hypothesis: (a) direct investigations of the sugar-depression relationship, including observational studies and animal experiments; and (b) a voluminous literature on the depressogenic physiological effects of added sugar ingestion.

## Sugar consumption and depression risk

### Longitudinal cohort studies

Several large epidemiological studies have explored the sugar-depression link by tracking participants' dietary habits and health outcomes prospectively over a span of several years. The Women's Health Initiative Observational Study, for example, followed 69,954 American women over a 3-year period [18]. Investigators observed that women in the top quintile of added sugar consumption (median 79.2 g daily) were at 23% greater risk of subsequent clinical depression than those in the bottom quintile (17.8 g daily), even after accounting for other relevant dietary factors, disease markers, and demographic variables. Another large trial, the NIH-AARP Diet and Health Study, found that the regular consumption of sugar-sweetened beverages (the leading dietary source of added sugars) conferred a 20% increase in depression incidence over a 4-to-5-year follow-up period [19]. A subsequent study of 15,546 Spanish university graduates likewise observed a heightened risk of depression among the heaviest consumers of sugar over a 10-year follow-up [20], and another recent cohort study also reported a positive association between sugar intake and the onset of future depressive symptoms [21]. Conversely, two recent cohort studies failed to find the hypothesized association [22,23]: Gopinath et al. [22] did not find a statistically significant association between total sugar intake and depressive symptoms, while Vermeulen et al. [23] similarly found no relationship between symptoms and a high-sugar dietary pattern. It should be noted, however, that unlike the aforementioned supporting studies [18,19,20], the two contrary studies [22,24] did not specifically examine the impact of added sugars, per se. Instead, they utilized dietary indices that conflated sugars with other simple carbohydrates—a less precise methodology that may have limited investigators' ability to detect a potential link between added sugars and depression.

### Cross-sectional surveys

Cross-sectional surveys provide additional support for the hypothesized sugar-depression relationship. For example, the incidence of depression for Australian adults was roughly 50% higher among regular versus non-regular soda drinkers [25]. Similarly, Chinese adults who reported drinking over seven ounces of soda daily were significantly more likely to report clinically salient depressive symptoms than those who consumed fewer [26]. Investigators have also found sugar-sweetened beverage intake to be positively associated with depressive symptoms among low-income obese Latino immigrants [27], and among Iranian children and adolescents [28]. Indeed, a recent meta-analysis of cohort and cross-sectional studies found that the consumption of sugar-sweetened beverages was linked to a significantly increased risk of depression [29].

### Animal models

Several published studies suggest that high-sugar diets may elicit depressive behaviors in rats. In comparison with rats fed standard (low-sugar) "rat chow," those provided with free access to high-sucrose food for two days a week over the course of seven weeks displayed

depressive and anhedonic behavioral shifts [30]. Rats given sugar-sweetened water for nine weeks likewise displayed depressive-like behavior in a forced-swim test paradigm [31]. Related work suggests that overfeeding sugar to adolescent rats induces subsequent depressive and anhedonic behavior during adulthood [32,33]. On the other hand, Pyndt Jørgensen et al. [34] have failed to find a relationship between sugar consumption and depressive behavior in mice, which suggests that the depressogenic potential of sugar may differ to some degree across species.

## Depressogenic physiological effects of added sugar ingestion

Although the above-reviewed studies generally support the hypothesized association between excess sugar consumption and the risk of depressive illness, most of the relevant evidence is merely correlational in nature. It remains unclear the degree to which added sugars contribute *causally* to the experience of depression. In fact, we are aware of no well-conducted experimental manipulations of sugar intake to evaluate the sweetener's direct causal impact on human depressive symptomatology.

Nevertheless, one additional line of evidence is worthy of consideration in this context—an extensive, robust research literature on the adverse physiological consequences of excess sugar consumption. Several such pathophysiological effects also exhibit a well-documented potential to induce depression (see Fig. 1), among them: systemic inflammation, gut microbiota disruption, perturbed dopaminergic reward signaling, insulin resistance, oxidative stress, and the generation of toxic *advanced glycation end-products* (AGEs). The impact of sugar ingestion on these depressogenic processes is reviewed in the sections that follow.

### Inflammation

Elevated systemic inflammation is recognized as a potent physiological trigger of depression [35]. Meta-analyses have found the concentrations of several inflammatory factors, for example, to be higher among depressed individuals than among non-depressed controls [36,37]. Inflammation is also associated with several characteristic depressive symptoms, including fatigue, sleep disruption, and appetite changes [38]. Finally, pro-inflammatory drugs carry the potential to induce depressive symptoms and to increase the risk of full-blown depression onset [39], while anti-inflammatory drugs possess anti-depressant therapeutic properties [40].

Dietary sugars can elicit inflammation in both humans and non-human animals. Pro-inflammatory states, marked by elevated levels of inflammatory factors and increased expression of inflammatory genes,




Fig. 1. Schematic overview of the physiological pathways linking sugar to depression.

characterize rats fed a high-fructose diet [41,42]. A similar pro-inflammatory response can be seen in humans following the consumption of sugar-sweetened beverages. Jameel, Phang, Wood, and Garg [43] found that participants' consumption of 50 g of dissolved fructose immediately increased levels of an important inflammatory factor, high-sensitivity c-reactive protein (hs-CRP). In fact, hs-CRP levels have been shown to double after just three weeks of daily exposure to sugar-sweetened beverages [44]. Additionally, a cross-sectional analysis of 17,689 U.S. adults found that measured hs-CRP levels significantly covaried with self-reported sugar consumption [45]. Another cross-sectional study likewise found that higher intake of added sugars was associated with higher levels of inflammation [46]. Added sugars may also trigger specific inflammatory processes localized within the brain (*neuroinflammation*), which are in turn linked with impaired cognitive function [47]. Rats exposed to high-sugar diets display impaired memory, along with increased inflammation and the expression of inflammatory genes within the hippocampus, a major brain nexus of memory consolidation and retrieval [48,49]. Significantly, cognitive deficits are a primary feature of depression, often contributing to the syndrome's characteristic psychosocial impairment [50].

In sum: added sugars have a profound effect on inflammatory processes within the body and brain, and inflammation may serve as a key mediator of sugar-induced depression onset. As discussed in the sections that follow, four additional depressive physiological pathways affected by sugar intake—disruption of the gut-brain axis, oxidative stress, insulin resistance, and the production of toxic advanced glycation end products (AGEs)—are also associated with increased inflammation.

#### Microbiota dysfunction

Extensive interconnections have been discovered between the human central nervous system and the collection of microbes residing within the body—the *microbiota*—and researchers have recently begun to explore the microbiota's potential role in depression and other psychological disorders [51]. The human microbiota is altered by the quantity of microbe-accessible carbohydrates in the diet, including sugars like glucose and fructose [52], and high-sugar diets are thought to permit the proliferation of harmful opportunistic organisms that contribute to *dysbiosis*, a state of maladaptive microbiota imbalance [53]. Dysbiosis, in turn, is linked to numerous physical and psychological pathologies [54].

Although knowledge of the human microbiota remains nascent, animal models suggest that high-sugar diets can disrupt microbiota balance and functioning. Rats fed a high-fructose diet display an altered microbiota, and this disruption is further related to increased disease risk [55]. Magnusson et al. [56] found that mice briefly fed a high-sugar diet also displayed distinct changes within the microbiota, in comparison with mice fed standard or high-fat diets; the sugar-fed mice also evidenced a broad pattern of cognitive impairment. Analyses of gut bacteria in mice have likewise revealed that a long-term, high-fructose diet can substantially alter microbiota composition and induce dysbiosis [57,58]. Additionally, Jena et al. [59] found that a 30-day high-fructose diet suppressed beneficial microbe species in rats. Excessive fructose consumption is also linked to intestinal bacterial overgrowth, a process in which abnormally large numbers of bacteria grow in the small intestine [60].

Sugar-induced dysbiosis may contribute to depression primarily by promoting inflammatory processes throughout the body. In fact, microbiota-induced inflammation has been posited as a primary risk factor for the development of depression [61]. There is growing evidence, for example, that sugar can increase intestinal permeability, resulting in increased passage of bacteria through the intestinal barrier and the subsequent activation of pro-inflammatory signaling pathways [62]. Higher consumption of fructose has also been linked to an increased presence of *endotoxins* (bacterially-produced poisons) in the blood of

humans, indicative of increased intestinal permeability [63]. Bergheim et al. [64] provided mice with sugar-sweetened water for eight weeks, and they found that fructose-fed mice, in comparison with controls, had significantly higher levels of blood endotoxins, as well as sharply increased inflammation. Spruss et al. [60] and Sellmann et al. [65] both found a similar connection in rodents between fructose consumption and increased intestinal permeability. Added sugars may increase intestinal permeability by inhibiting microbes that maintain the intestinal barrier. Microbes of the genus *lactobacillus* have been found to decrease intestinal permeability [66,67], and this same genus seems to be suppressed by exposure to high-fructose diets [59]. There is also evidence that sugar directly affects the structure of the intestinal barrier, as high-fructose diets can interfere with the proteins necessary to maintain healthy intestinal barrier functioning [58,65].

Importantly, excessive intestinal permeability has been implicated in the pathophysiology of depression. Depressed individuals have been found to have an increased passage of bacteria through the intestinal barrier [68] and an enhanced immune and oxidative stress response against such intestinal bacteria [69,70]. Likewise, bacteria of the genus *Alistipes*, microbes associated with intestinal inflammation [71], have been found to be overrepresented in depressed individuals [72,73]. Clinical improvement of depression among patients in treatment—even among those receiving psychotherapy—is also associated with a decrease of both intestinal permeability and inflammation [68].

#### Dopaminergic dysregulation

The neurotransmitter dopamine (DA) plays a central role in brain circuitry that regulates reward-based behavior and key motivational processes. Signaling abnormalities that arise from chronic, stress-induced activation of dopamine-based (*dopaminergic*) pathways seem to contribute to depressive and anhedonic states through a down-regulation of dopaminergic activity [74]. Indeed, depressed humans evince significantly lower baseline dopaminergic brain activity than healthy controls [75]. Targeted interference of DA activity in mice has also been shown to induce depressive-like behavior under stressful conditions [76], and Chang and Grace [77] have demonstrated that rats exposed to chronic, unpredictable stressors manifest both depressive-like behavior and reduced DA activity.

Interestingly, acute sugar consumption tends to stimulate the DA system: even a sip of sugary beverage immediately activates rewards areas within the human brain [78,79]. In rats, Bassareo et al. [80] witnessed a significant burst of DA release in the brain's nucleus accumbens, a region centrally involved in reward and reinforcement, just ten minutes after the animals were fed sugar for the first time. Hajnal, Smith, and Norgren [81] found that DA release in the nucleus accumbens increased as a function of the concentration of ingested sugar, suggesting that added sugars have a dose-dependent impact on DA activity. Furthermore, DA antagonists have been demonstrated to block the reinforcing properties of sugar ingestion [82,83].

In addition to acutely stimulating cerebral DA activity, excessive sugar intake is associated with maladaptive changes in the structure and function of DA pathways—pathology consistent with the observed connection between DA dysregulation and depression [74]. DA in the *striatum*, a component of the brain's reward system, is particularly affected by excessive added sugar consumption. Rats chronically fed a high-sugar diet have been found to exhibit decreased concentrations of striatal DA [84] and reduced expression of the D2 dopamine receptor, which helps regulate reward-based DA activity [85]. Rodents that lose access to sugar after long-term overfeeding similarly display decreased DA levels [86,87]. Additionally, long-term high-sugar diets are associated with decreased striatal D2 receptor binding, further indicative of a sugar-induced reduction of D2 receptor density [88,89].

In sum: a substantial body of evidence suggests that chronic added sugar ingestion can interfere with intrinsic reward systems in a manner capable of inducing anhedonia and motivational deficits. Both are

hallmark symptoms and maintenance factors of depression.

#### Oxidative stress

*Oxidative stress* refers to a physiological state characterized by excess metabolic molecular by-products known as *reactive oxygen species*, which in turn induce toxic effects such as cellular destruction, inflammation, and accelerated aging [90]. Two recent meta-analyses suggest that depression is characterized by elevated levels of oxidative stress [91,92], which has been observed to normalize either with antidepressant drug treatment [92] or interventions such as vitamin D supplementation or even cognitive psychotherapy [93,94]. Notably, direct administration of a compound with antioxidant properties has also been shown to have antidepressant effects [95].

Accumulating research suggests that added sugar consumption can induce oxidative stress. Cosentino et al. [96] found that human cells exposed to high levels of glucose—a state commonly observed following acute sugar ingestion—produced elevated levels of reactive oxygen species. In fact, a heightened oxidative stress response has been observed in vivo among healthy adults immediately following high-dose glucose consumption [97]. Short-term supplementation with fructose-sweetened beverages has also increased oxidative stress in otherwise healthy overweight/obese individuals [98]. Likewise, markers of oxidative stress rapidly rose in preterm neonates after the administration of sucrose for pain management [99].

Added sugars are also associated with oxidative stress in rodents. Markers of oxidative stress are increased in rats chronically fed a high-fructose diet [41,100], while levels of antioxidants (and antioxidative enzymes) are decreased under the same dietary conditions [41,42]. Elevated production of reactive oxygen species has also been found in mice fed excessive sugar [101]. Damage related to fructose-induced oxidative stress has been viewed in the liver of rats [100]. Additionally, Lopes et al. [102] demonstrated that a single injection of fructose can exacerbate oxidative stress within the brains of rats.

In sum: oxidative stress contributes to the dysregulation of various physiological pathways associated with depressive processes. Together with inflammation, it may mediate the depressogenic impact of excessive added sugar consumption.

#### Insulin resistance

Insulin is a key hormone that permits glucose to enter cells for conversion to energy. *Insulin resistance* develops when receptors become less sensitive to the hormone, thereby limiting its effectiveness and in turn depriving cells of adequate fuel. Insulin receptors are present throughout the brain, and they play a central role in regulating its use of glucose for energy [103]. There is even emerging evidence (reviewed in [104]) that depression can directly arise from the brain's insulin resistance and its ensuing disruption of energy utilization. For example, adults with insulin resistance are at heightened risk for the development of future depression [105], while cross-sectional epidemiological work suggests a low- to moderate-strength association between insulin resistance and depression incidence [106].

Excessive sugar consumption is strongly implicated as a potential driver of insulin resistance. In fact, high-sugar diets are frequently used to induce insulin resistance in rodent models. Pagliassotti, Prach, Koppenhafer, and Pan [107] demonstrated that a high-sugar diet fed to rats can elicit insulin resistance in the liver and muscles within one and eight weeks, respectively. Fructose consumption has also been shown to down-regulate the expression of insulin receptors and other insulin-related proteins in rats [108]. Humans are similarly susceptible to sugar-induced insulin dysregulation. Couchevin et al. [109], for example, found that a short-term, high-fructose diet increased fasting glucose levels in healthy men and women, indicative of impaired insulin functioning. Stanhope et al. [110] likewise demonstrated that consumption of fructose-sweetened beverages decreased insulin

sensitivity over the course of ten weeks. In fact, even one week of fructose overfeeding is sufficient to decrease insulin sensitivity [111,112].

Significantly, insulin resistance interacts with other depressogenic processes. Inflammation and oxidative stress are both involved in the pathogenesis of insulin resistance [113,114]. In turn, insulin has anti-inflammatory and antioxidant properties which may become disrupted following insulin resistance [115,116]. Insulin resistance is also associated with perturbed dopamine signaling in rodents [116,117]. The bidirectional relationships between insulin dysregulation and key depressive processes highlight the detrimental systemic impact of added sugar consumption.

#### Advanced glycation end-products (AGEs)

*Advanced glycation end-products* (AGEs) are toxic molecules that form when sugar molecules react with proteins, lipids (fats), or other compounds in the body. Accumulating evidence suggests that excessive added sugar consumption promotes AGE formation by increasing the availability of glucose and fructose, thereby allowing these sugars to be irreversibly converted to AGEs within the body [118]. Animal models have demonstrated that high-fructose and high-glucose diets increase the concentration of AGEs in both blood and tissue [119,120]. Once formed, AGEs are associated with a cascade of adverse biological reactions, including oxidative stress, inflammation, and neurocognitive dysfunction [121,122,123].

In light of these broadly deleterious effects, AGEs have recently been proposed as novel markers of lifestyle-related diseases [124]—a category that includes depression [125]. In fact, a recent landmark investigation by van Dooren et al. [126] lends support to the hypothesized role of AGEs in depressive illness. Among a well-characterized sample of 862 Dutch adults (Maastricht Study), those with elevated AGE accumulations in skin tissue evidenced a 42% increase in the risk of depressive disorder [126]. Study investigators also observed a significant positive association between measured AGE levels and the severity of depressive symptoms.

Although the Maastricht Study is the only large-scale investigation to date of the depressive impact of AGEs, several additional lines of evidence lend support to the finding. First, depressed individuals have been observed to have low levels of circulating protective molecules that bind and neutralize AGEs before they are able to cause damage [127,128]. Depressed individuals also appear to be deficient in the expression of glyoxalase-1, an antioxidant enzyme that helps degrade AGEs and minimize their toxic impact [129]. Further, Franklin et al. [130] demonstrated that mice without functioning AGE receptors are resistant to stress-induced depressive-like behavior. Finally, rodents exposed to high AGE levels have been found to experience impaired neuronal growth (*neurogenesis*) in the hippocampus—a central neurological feature of depression—and this reduced neurogenesis was reversed by administration of an anti-AGE drug [131]. Taken together, these findings provide substantial support for the hypothesis that AGEs exert depressogenic effects.

#### Conclusions and future directions

Multiple distinct lines of evidence generally converge in suggesting that the consumption of added sugars may induce depressogenic effects. The strongest evidence, by far, comes from the extensive literature on the pathophysiological consequences of added sugar ingestion, which collectively serve as candidate mediational mechanisms through which added sugars may adversely influence well-being. As reviewed, sugar intake promotes numerous maladaptive processes capable of inducing depressive symptomatology (depicted in Fig. 1): microbiota dysfunction (an altered *gut-brain axis*), disordered dopamine signaling, oxidative stress, insulin resistance, and the generation of advanced glycation end-products (AGEs). Most of these processes also promote pathological

inflammation—itself a particularly robust, well-established risk factor for depression [35]. Although the existing evidence regarding the hypothesized sugar-depression link is still neither definitive nor conclusive due to the absence of rigorous experimental manipulations of added sugar intake among sufficiently large and randomly assigned participant samples, we nonetheless regard the extant body of relevant research as persuasive.

Much more extensive investigation will be necessary, of course, to fully elucidate the sweetener's hypothesized depressogenic potential in humans. Ideally, such work would not only feature the experimental control of sugar intake at varying levels, but also the inclusion of relevant biomarkers of sugar-induced depressogenic processes. In a similar vein, it may prove valuable to evaluate the *psychological* impact of dietary interventions specifically designed to help reduce the consumption of added sugars—and thereby to reduce the occurrence of inflammation, oxidative stress, gut dysbiosis, and so on. Such interventions may carry the potential to reduce depressive symptoms (or to prevent their future occurrence), at least among the subset of patients with high baseline sugar consumption.

It is also necessary to clarify the amount (“dosage”) of sugar required to induce psychopathological consequences. Gangwisch et al. [18] observed the lowest risk of depression onset among those in the lowest quintile (20%) of added sugar intake, who had a median consumption of 17.8 g daily. Notably, this amount is not far below the suggested daily limit<sup>b</sup> recommended by the American Heart Association [132], and we believe it serves as a reasonable placeholder guideline while additional research is conducted—particularly for those battling depression or at high risk of future onset of depressive illness. There likely also exists considerable individual variation—derived from genetic, epigenetic, and gut microbial differences—in the physiological response to added sugars. That is, some people are probably far more sensitive than others to the sweetener's depressogenic effects. The discovery of any such reliable individual differences would likely be of both clinical and theoretical importance to the field.

Finally, to the degree that the connection between added sugars and depressive illness is robustly established by further research, it may desirable to develop low-sugar dietary interventions that can be effectively implemented at a population level. Unlike existing low-carb clinical diets, which typically require the extremely limited consumption of any carbohydrates [133,134], depression-focused diets could aim specifically to reduce the intake of added sugars while simultaneously ignoring other types of carbohydrates, thereby eliminating the need for restrictive (or expensive) dietary changes that act as a barrier to adherence [135,136]. One simple change that could have a substantial impact on added sugar intake is the reduction or elimination of sugar-sweetened beverages. Such beverages are the single leading source of added sugars [137] and they have limited nutritional value, making them a prime candidate for reduction or elimination. Educational and behavioral interventions (e.g., substituting water for sodas) have been shown to successfully reduce intake of sugar-sweetened beverages in both children and adults, as have public policy strategies such as targeted sugary beverage taxation [138,139,140]. However, the long-term effectiveness and sustainability of such strategies remain largely unknown [138], as do their potential protective benefit vis-à-vis depressive illness. Other high-sugar, low-nutrition foods, such as candy, should also be considered prime candidates for reduction. Regardless of the intervention strategy, excessive restriction (e.g., the banning of all sugar products) would likely prove counter-productive, inasmuch as restrictive dietary patterns are associated with disordered eating [141]. Moreover, there is some early evidence that reducing sugar intake can actually increase its reinforcement value—an important caveat for investigators to bear in mind during the development of novel dietary interventions [142].

<sup>b</sup> The suggested daily limit is 25 g for women and 36 g for men.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- [1] Martinez Steele E, Baraldi LG, Louzada ML, Moubarac JC, Mozaffarian D, Monteiro CA. Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open* 2016;6(3):e009892 <https://doi.org/10.1136/bmjopen-2015-009892>.
- [2] Ng SW, Slining MM, Popkin BM. Use of caloric and noncaloric sweeteners in US consumer packaged foods, 2005–2009. *J Acad Nutr Diet* 2012;112(11). <https://doi.org/10.1016/j.jand.2012.07.009>.
- [3] Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 2012;346:e7492 <https://doi.org/10.1136/bmj.e7492>.
- [4] Yang Q, Zhang Z, Gregg EW, Flanders WD, Merritt R, Hu FB. Added sugar intake and cardiovascular diseases mortality among US adults. *JAMA Intern Med* 2014;174(4):516–24. <https://doi.org/10.1001/jamainternmed.2013.13563>.
- [5] Imamura F, O'Connor L, Ye Z, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ* 2015;351:h3576 <https://doi.org/10.1136/bmj.h3576>.
- [6] Peres MA, Sheiham A, Liu P, et al. Sugar consumption and changes in dental caries from childhood to adolescence. *J Dent Res* 2016;95(4):388–94. <https://doi.org/10.1177/0022034515625907>.
- [7] U.S. Department of Health and Human Services and U.S. Department of Agriculture (December 2015). 2015–2020 Dietary Guidelines for Americans. 8th Edition. Retrieved from <http://health.gov/dietaryguidelines/2015/guidelines/>.
- [8] Logan AC, Jacka FN. Nutritional psychiatry research: an emerging discipline and its intersection with global urbanization, environmental challenges and the evolutionary mismatch. *J Physiol Anthropol* 2014;33:22. <https://doi.org/10.1186/1880-6805-33-22>.
- [9] World Health Organization (2017). Depression and other common mental disorders: Global health estimates Retrieved from [http://www.who.int/mental\\_health/management/depression/prevalence\\_global\\_health\\_estimates/en/](http://www.who.int/mental_health/management/depression/prevalence_global_health_estimates/en/).
- [10] Meyer BJ, Kolanu N, Griffiths DA, Grounds B, Howe PR, Kreis IA. Food groups and fatty acids associated with self-reported depression: an analysis from the Australian National Nutrition and Health Surveys. *Nutrition* 2013;29(7–8):1042–7. <https://doi.org/10.1016/j.nut.2013.02.006>.
- [11] Marx W, Moseley G, Berk M, Jacka F. Nutritional psychiatry: the present state of the evidence. *Proc Nutr Soc* 2017;1–10. <https://doi.org/10.1017/s0029665117002026>.
- [12] Firth J, Marx W, Dash S, et al. The effects of dietary improvement on symptoms of depression and anxiety: a meta-analysis of randomized controlled trials. *Psychosom Med* 2019;81(3):265–80. <https://doi.org/10.1097/psy.0000000000000673>.
- [13] Lassale C, Batty GD, Baghdadli A, et al. Healthy dietary indices and risk of depressive outcomes: a systematic review and meta-analysis of observational studies. *Mol Psychiatry* 2019;24(7):965–86. <https://doi.org/10.1038/s41380-018-0237-8>.
- [14] Lai JS, Hiles S, Bisquera A, Hure AJ, McEvoy M, Attia J. A systematic review and meta-analysis of dietary patterns and depression in community-dwelling adults. *Am J Clin Nutr* 2014;99(1):181–97. <https://doi.org/10.3945/ajcn.113.069880>.
- [15] Li Y, Lv MR, Wei YJ, et al. Dietary patterns and depression risk: a meta-analysis. *Psychiatry Res* 2017;253:373–82. <https://doi.org/10.1016/j.psychres.2017.04.020>.
- [16] Mantantzis K, Schlaghecken F, Sunram-Lea SI, Maylor EA. Sugar rush or sugar crash? A meta-analysis of carbohydrate effects on mood. *Neurosci Biobehav Rev* 2019;101:45–67. <https://doi.org/10.1016/j.neubiorev.2019.03.016>.
- [17] Jacka FN. Nutritional psychiatry: where to next? *EBioMedicine* 2017;17:24–9. <https://doi.org/10.1016/j.ebiom.2017.02.020>.
- [18] Gangwisch JE, Hale L, Garcia L, et al. High glycemic index diet as a risk factor for depression: analyses from the Women's Health Initiative. *Am J Clin Nutr* 2015;102(2):454–63. <https://doi.org/10.3945/ajcn.114.103846>.
- [19] Guo X, Park Y, Freedman ND, et al. Sweetened beverages, coffee, and tea and depression risk among older US adults. *PLoS ONE* 2014;9(4).
- [20] Sanchez-Villegas A, Zazpe I, Santiago S, Perez-Cornago A, Martinez-Gonzalez MA, Lahortiga-Ramos F. Added sugars and sugar-sweetened beverage consumption, dietary carbohydrate index and depression risk in the Seguimiento Universidad de Navarra (SUN) Project. *Br J Nutr* 2018;119(2):211–21. <https://doi.org/10.1017/s0007114517003361>.
- [21] Knüppel A, Shipley MJ, Llewellyn CH, Brunner EJ. Sugar intake from sweet food and beverages, common mental disorder and depression: prospective findings from the Whitehall II study. *Sci Rep* 2017;7(1):6287. <https://doi.org/10.1038/s41598-017-05649-7>.
- [22] Gopinath B, Flood VM, Burlutsky G, Louie JC, Mitchell P. Association between carbohydrate nutrition and prevalence of depressive symptoms in older adults. *Br J Nutr* 2016;116(12):2109–14. <https://doi.org/10.1017/s0007114516004311>.
- [23] Vermeulen E, Stronks K, Snijder MB, et al. A combined high-sugar and high-saturated-fat dietary pattern is associated with more depressive symptoms in a multi-ethnic population: the HELIUS (Healthy Life in an Urban Setting) study. *Public*

- Health Nutr 2017;20(13):2374–82. <https://doi.org/10.1017/s1368980017001550>.
- [24] Vermeulen E, Knuppel A, Shipley MJ, et al. High-sugar, high-saturated-fat dietary patterns are not associated with depressive symptoms in middle-aged adults in a prospective study. *J Nutr* 2018;148(10):1598–604. <https://doi.org/10.1093/jn/nxy154>.
- [25] Shi Z, Taylor AW, Wittert G, Goldney R, Gill TK. Soft drink consumption and mental health problems among adults in Australia. *Public Health Nutr* 2010;13(7):1073–9. <https://doi.org/10.1017/s1368980009993132>.
- [26] Yu B, He H, Zhang Q, et al. Soft drink consumption is associated with depressive symptoms among adults in China. *J Affect Disord* 2015;172:422–7. <https://doi.org/10.1016/j.jad.2014.10.026>.
- [27] Drieling RL, Goldman Rosas L, Ma J, Stafford RS. Community resource utilization, psychosocial health, and sociodemographic factors associated with diet and physical activity among low-income obese Latino immigrants. *J Acad Nutr Diet* 2014;114(2):257–65. <https://doi.org/10.1016/j.jand.2013.07.025>.
- [28] Zahedi H, Kelishadi R, Heshmat R, et al. Association between junk food consumption and mental health in a national sample of Iranian children and adolescents: the CASPIAN-IV study. *Nutrition* 2014;30(11–12):1391–7. <https://doi.org/10.1016/j.nut.2014.04.014>.
- [29] Hu D, Cheng L, Jiang W. Sugar-sweetened beverages consumption and the risk of depression: a meta-analysis of observational studies. *J Affect Disord* 2019;245:348–55. <https://doi.org/10.1016/j.jad.2018.11.015>.
- [30] Iemolo A, Valenza M, Tozier L, et al. Withdrawal from chronic, intermittent access to a highly palatable food induces depressive-like behavior in compulsive eating rats. *Behav Pharmacol* 2012;23(5–6):593–602. <https://doi.org/10.1097/FBP.0b013e328357697f>.
- [31] Lemos C, Rial D, Goncalves FQ, et al. High sucrose consumption induces memory impairment in rats associated with electrophysiological modifications but not with metabolic changes in the hippocampus. *Neuroscience* 2016;315:196–205. <https://doi.org/10.1016/j.neuroscience.2015.12.018>.
- [32] Harrell CS, Burgado J, Kelly SD, Johnson ZP, Neigh GN. High-fructose diet during periadolescent development increases depressive-like behavior and remodels the hypothalamic transcriptome in male rats. *Psychoneuroendocrinology* 2015;62:252–64. <https://doi.org/10.1016/j.psneuen.2015.08.025>.
- [33] Vendruscolo LF, Gueye AB, Darnaudey M, Ahmed SH, Cador M. Sugar over-consumption during adolescence selectively alters motivation and reward function in adult rats. *PLoS One* 2010;5(2):e9296. <https://doi.org/10.1371/journal.pone.0009296>.
- [34] Pyndt Jørgensen B, Hansen JT, Krych L, et al. A possible link between food and mood: dietary impact on gut microbiota and behavior in BALB/c mice. *PLoS One* 2014;9(8):e103398. <https://doi.org/10.1371/journal.pone.0103398>.
- [35] Slavich GM, Irwin MR. From stress to inflammation and major depressive disorder: a social signal transduction theory of depression. *Psychol Bull* 2014;140(3):774–815. <https://doi.org/10.1037/a0035302>.
- [36] Dowlati Y, Herrmann N, Swardfager W, et al. A meta-analysis of cytokines in major depression. *Biol Psychiatry* 2010;67(5):446–57. <https://doi.org/10.1016/j.biopsych.2009.09.033>.
- [37] Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom Med* 2009;71(2):171–86. <https://doi.org/10.1097/PSY.0b013e3283181907c1b>.
- [38] Jokela M, Virtanen M, Batty GD, Kivimaki M. Inflammation and specific symptoms of depression. *JAMA Psychiatry* 2016;73(1):87–8. <https://doi.org/10.1001/jamapsychiatry.2015.1977>.
- [39] Capuron L, Miller AH. Cytokines and psychopathology: lessons from interferon-alpha. *Biol Psychiatry* 2004;56(11):819–24. <https://doi.org/10.1016/j.biopsych.2004.02.009>.
- [40] Köhler O, Benros ME, Nordentoft M, et al. Effect of anti-inflammatory treatment on depression, depressive symptoms, and adverse effects: a systematic review and meta-analysis of randomized clinical trials. *JAMA Psychiatry* 2014;71(12):1381–91. <https://doi.org/10.1001/jamapsychiatry.2014.1611>.
- [41] Castro MC, Massa ML, Arbelaez LG, Schinella G, Gagliardino JJ, Francini F. Fructose-induced inflammation, insulin resistance and oxidative stress: a liver pathological triad effectively disrupted by lipolic acid. *Life Sci* 2015;137:1–6. <https://doi.org/10.1016/j.lfs.2015.07.010>.
- [42] Kovacevic S, Nestorov J, Matic G, Elakovic I. Fructose-enriched diet induces inflammation and reduces antioxidative defense in visceral adipose tissue of young female rats. *Eur J Nutr* 2015. <https://doi.org/10.1007/s00394-015-1065-0>.
- [43] Jameel F, Phang M, Wood LG, Garg ML. Acute effects of feeding fructose, glucose and sucrose on blood lipid levels and systemic inflammation. *Lipids Health Dis* 2014;13:195. <https://doi.org/10.1186/1476-511x-13-195>.
- [44] Aeberli I, Gerber PA, Hochuli M, et al. Low to moderate sugar-sweetened beverage consumption impairs glucose and lipid metabolism and promotes inflammation in healthy young men: a randomized controlled trial. *Am J Clin Nutr* 2011;94(2):479–85. <https://doi.org/10.3945/ajcn.111.013540>.
- [45] Mazidi M, Kengne AP, Mikhailidis DP, Cicero AF, Banach M. Effects of selected dietary constituents on high-sensitivity C-reactive protein levels in U.S. adults. *Ann Med* 2017;1–6. <https://doi.org/10.1080/07853890.2017.1325967>.
- [46] O'Connor L, Imamura F, Brage S, Griffin SJ, Wareham NJ, Forouhi NG. Intakes and sources of dietary sugars and their association with metabolic and inflammatory markers. *Clin Nutr* 2017. <https://doi.org/10.1016/j.clnu.2017.05.030>.
- [47] Carvalho AF, Miskowiak KK, Hyphantis TN, et al. Cognitive dysfunction in depression – pathophysiology and novel targets. *CNS Neurol Disord Drug Targets* 2014;13(10):1819–35.
- [48] Beilharz JE, Maniam J, Morris MJ. Short-term exposure to a diet high in fat and sugar, or liquid sugar, selectively impairs hippocampal-dependent memory, with differential impacts on inflammation. *Behav Brain Res* 2016;306:1–7. <https://doi.org/10.1016/j.bbr.2016.03.018>.
- [49] Hsu TM, Konanur VR, Taing L, et al. Effects of sucrose and high fructose corn syrup consumption on spatial memory function and hippocampal neuroinflammation in adolescent rats. *Hippocampus* 2015;25(2):227–39. <https://doi.org/10.1002/hipo.22368>.
- [50] McIntyre RS, Cha DS, Soczynska JK, et al. Cognitive deficits and functional outcomes in major depressive disorder: determinants, substrates, and treatment interventions. *Depress Anxiety* 2013;30(6):515–27. <https://doi.org/10.1002/da.22063>.
- [51] Kim YK, Shin C. The microbiota-gut-brain axis in neuropsychiatric disorders: pathophysiological mechanisms and novel treatments. *Curr Neuropharmacol* 2017. <https://doi.org/10.2174/1570159x15666170915141036>.
- [52] Sonnenburg ED, Sonnenburg JL. Starving our microbial self: the deleterious consequences of a diet deficient in microbiota-accessible carbohydrates. *Cell Metab* 2014;20(5):779–86. <https://doi.org/10.1016/j.cmet.2014.07.003>.
- [53] Payne AN, Chassard C, Lacroix C. Gut microbial adaptation to dietary consumption of fructose, artificial sweeteners and sugar alcohols: Implications for host-microbe interactions contributing to obesity. *Obes Rev* 2012;13(9):799–809. <https://doi.org/10.1111/j.1467-789X.2012.01009.x>.
- [54] de Vos WM, de Vos EA. Role of the intestinal microbiome in health and disease: from correlation to causation. *Nutr Rev* 2012;70(Suppl 1):S45–56. <https://doi.org/10.1111/j.1753-4887.2012.00505.x>.
- [55] Wei X, Song M, Yin X, et al. Effects of dietary different doses of copper and high fructose feeding on rat fecal metabolome. *J Proteome Res* 2015. <https://doi.org/10.1021/acs.jproteome.5b00596>.
- [56] Magnusson KR, Hauck L, Jeffrey BM, et al. Relationships between diet-related changes in the gut microbiome and cognitive flexibility. *Neuroscience* 2015;300:128–40. <https://doi.org/10.1016/j.neuroscience.2015.05.016>.
- [57] Ferrere G, Leroux A, Wrzosek L, et al. Activation of Kupffer cells is associated with a specific dysbiosis induced by fructose or high fat diet in mice. *PLoS One* 2016;11(1):e0146177. <https://doi.org/10.1371/journal.pone.0146177>.
- [58] Wagnerberger S, Spruss A, Kanuri G, et al. Toll-like receptors 1–9 are elevated in livers with fructose-induced hepatic steatosis. *Br J Nutr* 2012;107(12):1727–38. <https://doi.org/10.1017/s0007114511004983>.
- [59] Jena PK, Singh S, Prajapati B, Nareshkumar G, Mehta T, Seshadri S. Impact of targeted specific antibiotic delivery for gut microbiota modulation on high-fructose-fed rats. *Appl Biochem Biotechnol* 2014;172(8):3810–26. <https://doi.org/10.1007/s12010-014-0772-y>.
- [60] Spruss A, Kanuri G, Wagnerberger S, Haub S, Brischoff SC, Bergheim I. Toll-like receptor 4 is involved in the development of fructose-induced hepatic steatosis in mice. *Hepatology* 2009;50(4):1094–104. <https://doi.org/10.1002/hep.23122>.
- [61] Berk M, Williams LJ, Jacka FN, et al. So depression is an inflammatory disease, but where does the inflammation come from? *BMC Med* 2013;11:200. <https://doi.org/10.1186/1741-7015-11-200>.
- [62] Piya MK, Harte AL, McTernan PG. Metabolic endotoxaemia: is it more than just a gut feeling? *Curr Opin Lipidol* 2013;24(1):78–85. <https://doi.org/10.1097/MOL.0b013e32835b4431>.
- [63] Thuy S, Ladurner R, Volynets V, et al. Nonalcoholic fatty liver disease in humans is associated with increased plasma endotoxin and plasminogen activator inhibitor 1 concentrations and with fructose intake. *J Nutr* 2008;138(8):1452–5.
- [64] Bergheim I, Weber S, Vos M, et al. Antibiotics protect against fructose-induced hepatic lipid accumulation in mice: role of endotoxin. *J Hepatol* 2008;48(6):983–92. <https://doi.org/10.1016/j.jhep.2008.01.035>.
- [65] Sellmann C, Priebs J, Landmann M, et al. Diets rich in fructose, fat or fructose and fat alter intestinal barrier function and lead to the development of nonalcoholic fatty liver disease over time. *J Nutr Biochem* 2015;26(11):1183–92. <https://doi.org/10.1016/j.jnutbio.2015.05.011>.
- [66] Liu Z, Zhang P, Ma Y, et al. Lactobacillus plantarum prevents the development of colitis in IL-10-deficient mouse by reducing the intestinal permeability. *Mol Biol Rep* 2011;38(2):1353–61. <https://doi.org/10.1007/s11033-010-0237-5>.
- [67] Zareie M, Johnson-Henry K, Jury J, et al. Probiotics prevent bacterial translocation and improve intestinal barrier function in rats following chronic psychological stress. *Gut* 2006;55(11):1553–60. <https://doi.org/10.1136/gut.2005.080739>.
- [68] Kéri S, Szabo C, Kelemen O. Expression of Toll-like receptors in peripheral blood mononuclear cells and response to cognitive-behavioral therapy in major depressive disorder. *Brain Behav Immun* 2014;40:235–43. <https://doi.org/10.1016/j.bbi.2014.03.020>.
- [69] Maes M, Kubera M, Leunis JC, Berk M. Increased IgA and IgM responses against gut commensals in chronic depression: further evidence for increased bacterial translocation or leaky gut. *J Affect Disord* 2012;141(1):55–62. <https://doi.org/10.1016/j.jad.2012.02.023>.
- [70] Maes M, Kubera M, Leunis JC, Berk M, Geffard M, Bosmans E. In depression, bacterial translocation may drive inflammatory responses, oxidative and nitrosative stress (O&NS), and autoimmune responses directed against O&NS-damaged neopeptides. *Acta Psychiatr Scand* 2013;127(5):344–54. <https://doi.org/10.1111/j.1600-0447.2012.01908.x>.
- [71] Qiu X, Zhang F, Yang X, et al. Changes in the composition of intestinal fungi and their role in mice with dextran sulfate sodium-induced colitis. *Sci Rep* 2015;5:10416. <https://doi.org/10.1038/srep10416>.
- [72] Jiang H, Ling Z, Zhang Y, et al. Altered fecal microbiota composition in patients with major depressive disorder. *Brain Behav Immun* 2015;48:186–94. <https://doi.org/10.1016/j.bbi.2015.03.016>.
- [73] Naserifabrouei A, Hestad K, Avershina E, et al. Correlation between the human fecal microbiota and depression. *Neurogastroenterol Motil* 2014;26(8):1155–62. <https://doi.org/10.1111/nmo.12378>.

- [74] Grace AA. Dysregulation of the dopamine system in the pathophysiology of schizophrenia and depression. *Nat Rev Neurosci* 2016;17(8):524–32. <https://doi.org/10.1038/nrn.2016.57>.
- [75] Lambert G, Johansson M, Agren H, Friberg P. Reduced brain norepinephrine and dopamine release in treatment-refractory depressive illness: evidence in support of the catecholamine hypothesis of mood disorders. *Arch Gen Psychiatry* 2000;57(8):787–93.
- [76] Tye KM, Mirzabekov JJ, Warden MR, et al. Dopamine neurons modulate neural encoding and expression of depression-related behaviour. *Nature* 2013;493(7433):537–41. <https://doi.org/10.1038/nature11740>.
- [77] Chang CH, Grace AA. Amygdala-ventral pallidum pathway decreases dopamine activity after chronic mild stress in rats. *Biol Psychiatry* 2014;76(3):223–30. <https://doi.org/10.1016/j.biopsych.2013.09.020>.
- [78] Burger KS, Stice E. Neural responsiveness during soft drink intake, anticipation, and advertisement exposure in habitually consuming youth. *Obesity (Silver Spring)* 2014;22(2):441–50. <https://doi.org/10.1002/oby.20563>.
- [79] Stice E, Burger KS, Yokum S. Relative ability of fat and sugar tastes to activate reward, gustatory, and somatosensory regions. *Am J Clin Nutr* 2013;98(6):1377–84. <https://doi.org/10.3945/ajcn.113.069443>.
- [80] Bassareo V, Cucca F, Musio P, Lecca D, Frau R, Di Chiara G. Nucleus accumbens shell and core dopamine responsiveness to sucrose in rats: role of response contingency and discriminative/conditioned cues. *Eur J Neurosci* 2015;41(6):802–9. <https://doi.org/10.1111/ejn.12839>.
- [81] Hajnal A, Smith GP, Norgren R. Oral sucrose stimulation increases accumbens dopamine in the rat. *Am J Physiol Regul Integr Comp Physiol* 2004;286(1):R31–7. <https://doi.org/10.1152/ajpregu.00282.2003>.
- [82] Kraft TT, Yakubov Y, Huang D, et al. Dopamine D1 and opioid receptor antagonists differentially reduce the acquisition and expression of fructose-conditioned flavor preferences in BALB/c and SWR mice. *Physiol Behav* 2015;151:213–20. <https://doi.org/10.1016/j.physbeh.2015.07.028>.
- [83] Dela Cruz JA, Coke T, Icaza-Cukali D, Khalifa N, Bodnar RJ. Roles of NMDA and dopamine D1 and D2 receptors in the acquisition and expression of flavor preferences conditioned by oral glucose in rats. *Neurobiol Learn Mem* 2014;114:223–30. <https://doi.org/10.1016/j.nlm.2014.07.004>.
- [84] Ahmed S, Kashem MA, Sarker R, Ahmed EU, Hargreaves GA, McGregor IS. Neuroadaptations in the striatal proteome of the rat following prolonged excessive sucrose intake. *Neurochem Res* 2014;39(5):815–24. <https://doi.org/10.1007/s11064-014-1274-6>.
- [85] Levy A, Marshall P, Zhou Y, et al. Fructose:glucose ratios—A study of sugar self-administration and associated neural and physiological responses in the rat. *Nutrients* 2015;7(5):3869–90. <https://doi.org/10.3390/nu7053869>.
- [86] Kim S, Shou J, Abera S, Ziff EB. Sucrose withdrawal induces depression and anxiety-like behavior by Kir2.1 upregulation in the nucleus accumbens. *Neuropharmacology* 2018;130:10–7. <https://doi.org/10.1016/j.neuropharm.2017.11.041>.
- [87] Avena NM, Bocarsly ME, Rada P, Kim A, Hoebel BG. After daily bingeing on a sucrose solution, food deprivation induces anxiety and accumbens dopamine/acetylcholine imbalance. *Physiol Behav* 2008;94(3):309–15. <https://doi.org/10.1016/j.physbeh.2008.01.008>.
- [88] Colantuoni C, Schwenker J, McCarthy J, et al. Excessive sugar intake alters binding to dopamine and mu-opioid receptors in the brain. *NeuroReport* 2001;12(16):3549–52.
- [89] Bello NT, Lucas LR, Hajnal A. Repeated sucrose access influences dopamine D2 receptor density in the striatum. *NeuroReport* 2002;13(12):1575–8.
- [90] Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Curr Biol* 2014;24(10):R453–62. <https://doi.org/10.1016/j.cub.2014.03.034>.
- [91] Black CN, Bot M, Scheffer PG, Cuijpers P, Penninx BW. Is depression associated with increased oxidative stress? A systematic review and meta-analysis. *Psychoneuroendocrinology* 2015;51:164–75. <https://doi.org/10.1016/j.psyneuen.2014.09.025>.
- [92] Jimenez-Fernandez S, Gurpegui M, Diaz-Atienza F, Perez-Costillas L, Gerstenberg M, Correll CU. Oxidative stress and antioxidant parameters in patients with major depressive disorder compared to healthy controls before and after antidepressant treatment: results from a meta-analysis. *J Clin Psychiatry* 2015;76(12):1658–67. <https://doi.org/10.4088/JCP.14r09179>.
- [93] Sepelhrmanesh Z, Kolahdooz F, Abedi F, et al. Vitamin D supplementation affects the Beck Depression Inventory, insulin resistance, and biomarkers of oxidative stress in patients with major depressive disorder: a randomized, controlled clinical trial. *J Nutr* 2016;146(2):243–8. <https://doi.org/10.3945/jn.115.218883>.
- [94] Kaufmann FN, Gazal M, Mondin TC, et al. Cognitive psychotherapy treatment decreases peripheral oxidative stress parameters associated with major depression disorder. *Biol Psychol* 2015;110:175–81. <https://doi.org/10.1016/j.biopsycho.2015.08.001>.
- [95] Berk M, Copolov DL, Dean O, et al. N-acetyl cysteine for depressive symptoms in bipolar disorder—a double-blind randomized placebo-controlled trial. *Biol Psychiatry* 2008;64(6):468–75. <https://doi.org/10.1016/j.biopsych.2008.04.022>.
- [96] Cosentino F, Eto M, De Paolis P, et al. High glucose causes upregulation of cyclooxygenase-2 and alters prostanoid profile in human endothelial cells: role of protein kinase C and reactive oxygen species. *Circulation* 2003;107(7):1017–23.
- [97] Choi HJ, Jeon SY, Hong WK, et al. Effect of glucose ingestion in plasma markers of inflammation and oxidative stress: analysis of 16 plasma markers from oral glucose tolerance test samples of normal and diabetic patients. *Diabetes Res Clin Pract* 2013;99(2):e27–31. <https://doi.org/10.1016/j.diabres.2012.01.005>.
- [98] Hokayem M, Blond E, Vidal H, et al. Grape polyphenols prevent fructose-induced oxidative stress and insulin resistance in first-degree relatives of type 2 diabetic patients. *Diab Care* 2013;36(6):1454–61. <https://doi.org/10.2337/dc12-1652>.
- [99] Asmerom Y, Slater L, Boskovic DS, et al. Oral sucrose for heel lance increases adenosine triphosphate use and oxidative stress in preterm neonates. *J Pediatr* 2013;163(1):29–35.e21. <https://doi.org/10.1016/j.jpeds.2012.12.088>.
- [100] Cioffi F, Senese R, Lasala P, et al. Fructose-rich diet affects mitochondrial DNA damage and repair in rats. *Nutrients* 2017;9(4). <https://doi.org/10.3390/nu9040323>.
- [101] Choi Y, Abdelmegeed MA, Song BJ. Diet high in fructose promotes liver steatosis and hepatocyte apoptosis in C57BL/6J female mice: role of disturbed lipid homeostasis and increased oxidative stress. *Food Chem Toxicol* 2017;103:111–21. <https://doi.org/10.1016/j.fct.2017.02.039>.
- [102] Lopes A, Vilela TC, Taschetto L, et al. Evaluation of the effects of fructose on oxidative stress and inflammatory parameters in rat brain. *Mol Neurobiol* 2014;50(3):1124–30. <https://doi.org/10.1007/s12035-014-8676-y>.
- [103] Ghasemi R, Haeri A, Dargahi L, Mohamed Z, Ahmadiani A. Insulin in the brain: sources, localization and functions. *Mol Neurobiol* 2013;47(1):145–71. <https://doi.org/10.1007/s12035-012-8339-9>.
- [104] Yamagata AS, Mansur RB, Rizzo LB, Rosenstock T, McIntyre RS, Brietzke E. Selfish brain and selfish immune system interplay: a theoretical framework for metabolic comorbidities of mood disorders. *Neurosci Biobehav Rev* 2017;72:43–9. <https://doi.org/10.1016/j.neubiorev.2016.11.010>.
- [105] Ford AH, Flicker L, Hankey GJ, et al. Insulin resistance and depressive symptoms in older men: the health in men study. *Am J Geriatr Psychiatry* 2015;23(8):872–80. <https://doi.org/10.1016/j.jagp.2014.10.010>.
- [106] Kan C, Silva N, Golden SH, et al. A systematic review and meta-analysis of the association between depression and insulin resistance. *Diab Care* 2013;36(2):480–9. <https://doi.org/10.2337/dc12-1442>.
- [107] Pagliassotti MJ, Prach PA, Koppenhafer TA, Pan DA. Changes in insulin action, triglycerides, and lipid composition during sucrose feeding in rats. *Am J Physiol* 1996;271(5 Pt 2):R1319–26.
- [108] Meepprom A, Sompong W, Suwannaphet W, Yibchok-anun S, Adisakwattana S. Grape seed extract supplementation prevents high-fructose diet-induced insulin resistance in rats by improving insulin and adiponectin signalling pathways. *Br J Nutr* 2011;106(8):1173–81. <https://doi.org/10.1017/s0007114511001589>.
- [109] Couchepin C, Le KA, Bortolotti M, et al. Markedly blunted metabolic effects of fructose in healthy young female subjects compared with male subjects. *Diab Care* 2008;31(6):1254–6. <https://doi.org/10.2337/dc07-2001>.
- [110] Stanhope KL, Schwarz JM, Keim NL, et al. Consuming fructose-sweetened, not glucose-sweetened, beverages increases visceral adiposity and lipids and decreases insulin sensitivity in overweight/obese humans. *J Clin Invest* 2009;119(5):1322–34. <https://doi.org/10.1172/jci37385>.
- [111] Lecoultrre V, Egli L, Carrel G, et al. Effects of fructose and glucose overfeeding on hepatic insulin sensitivity and intrahepatic lipids in healthy humans. *Obesity (Silver Spring)* 2013;21(4):782–5. <https://doi.org/10.1002/oby.20377>.
- [112] Le KA, Ith M, Kreis R, et al. Fructose overconsumption causes dyslipidemia and ectopic lipid deposition in healthy subjects with and without a family history of type 2 diabetes. *Am J Clin Nutr* 2009;89(6):1760–5. <https://doi.org/10.3945/ajcn.2008.27336>.
- [113] Asghar A, Sheikh N. Role of immune cells in obesity induced low grade inflammation and insulin resistance. *Cell Immunol* 2017;315:18–26. <https://doi.org/10.1016/j.cellimm.2017.03.001>.
- [114] Tiganis T. Reactive oxygen species and insulin resistance: the good, the bad and the ugly. *Trends Pharmacol Sci* 2011;32(2):82–9. <https://doi.org/10.1016/j.tips.2010.11.006>.
- [115] Dandona P, Chaudhuri A, Ghanim H, Mohanty P. Proinflammatory effects of glucose and anti-inflammatory effect of insulin: relevance to cardiovascular disease. *Am J Cardiol* 2007;99(4a):15b–26b. <https://doi.org/10.1016/j.amjcard.2006.11.003>.
- [116] Kleinridders A, Cai W, Cappellucci L, et al. Insulin resistance in brain alters dopamine turnover and causes behavioral disorders. *Proc Natl Acad Sci USA* 2015;112(11):3463–8. <https://doi.org/10.1073/pnas.1500877112>.
- [117] Morris JK, Bomhoff GL, Gorres BK, et al. Insulin resistance impairs nigrostriatal dopamine function. *Exp Neurol* 2011;231(1):171–80. <https://doi.org/10.1016/j.expneurol.2011.06.005>.
- [118] Aragno M, Mastrocola R. Dietary sugars and endogenous formation of advanced glycation endproducts: emerging mechanisms of disease. *Nutrients* 2017;9(4). <https://doi.org/10.3390/nu9040385>.
- [119] Mastrocola R, Collino M, Rogazzo M, et al. Advanced glycation end products promote hepatosteatosis by interfering with SCAP-SREBP pathway in fructose-drinking mice. *Am J Physiol Gastrointest Liver Physiol* 2013;305(6):G398–407. <https://doi.org/10.1152/ajpgi.00450.2012>.
- [120] Cannizzaro L, Rossoni G, Savi F, et al. Regulatory landscape of AGE-RAGE-oxidative stress axis and its modulation by PPAR $\gamma$  activation in high fructose diet-induced metabolic syndrome. *Nutr Metab (Lond)* 2017;14:5. <https://doi.org/10.1186/s12986-016-0149-z>.
- [121] Li J, Liu D, Sun L, Lu Y, Zhang Z. Advanced glycation end products and neurodegenerative diseases: mechanisms and perspective. *J Neurol Sci* 2012;317(1–2):1–5. <https://doi.org/10.1016/j.jns.2012.02.018>.
- [122] Van Puyvelde K, Mets T, Njemini R, Beyer I, Bautmans I. Effect of advanced glycation end product intake on inflammation and aging: a systematic review. *Nutr Rev* 2014;72(10):638–50. <https://doi.org/10.1111/nure.12141>.
- [123] Yamagishi S, Maeda S, Matsui T, Ueda S, Fukami K, Okuda S. Role of advanced glycation end products (AGEs) and oxidative stress in vascular complications in diabetes. *Biochim Biophys Acta* 2012;1820(5):663–71. <https://doi.org/10.1016/j.bbagen.2011.03.014>.
- [124] Takeuchi M. Serum levels of toxic AGEs (TAGE) may be a promising novel biomarker for the onset/progression of lifestyle-related diseases. *Diagnostics (Basel)*

- 2016;6(2). <https://doi.org/10.3390/diagnostics6020023>.
- [125] Hidaka BH. Depression as a disease of modernity: explanations for increasing prevalence. *J Affect Disord* 2012;140(3):205–14. <https://doi.org/10.1016/j.jad.2011.12.036>.
- [126] van Dooren FE, Pouwer F, Schalkwijk CG, et al. Advanced glycation end product (AGE) accumulation in the skin is associated with depression: the Maastricht study. *Depress Anxiety* 2016. <https://doi.org/10.1002/da.22527>.
- [127] Chen G, Wu Y, Wang T, et al. Association between serum endogenous secretory receptor for advanced glycation end products and risk of type 2 diabetes mellitus with combined depression in the Chinese population. *Diab Technol Ther* 2012;14(10):936–42. <https://doi.org/10.1089/dia.2012.0072>.
- [128] Emanuele E, Martinelli V, Carlin MV, Fugazza E, Barale F, Politi P. Serum levels of soluble receptor for advanced glycation endproducts (sRAGE) in patients with different psychiatric disorders. *Neurosci Lett* 2011;487(1):99–102. <https://doi.org/10.1016/j.neulet.2010.10.003>.
- [129] Fujimoto M, Uchida S, Watanuki T, et al. Reduced expression of glyoxalase-1 mRNA in mood disorder patients. *Neurosci Lett* 2008;438(2):196–9. <https://doi.org/10.1016/j.neulet.2008.04.024>.
- [130] Franklin TC, Wohleb ES, Zhang Y, Fogaca M, Hare B, Duman RS. Persistent increase in microglial RAGE contributes to chronic stress-induced priming of depressive-like behavior. *Biol Psychiatry* 2017. <https://doi.org/10.1016/j.biopsych.2017.06.034>.
- [131] Wang SH, Sun ZL, Guo YJ, Yuan Y, Yang BQ. Diabetes impairs hippocampal function via advanced glycation end product mediated new neuron generation in animals with diabetes-related depression. *Toxicol Sci* 2009;111(1):72–9. <https://doi.org/10.1093/toxsci/kfp126>.
- [132] Johnson RK, Appel LJ, Brands M, et al. Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation* 2009;120(11):1011–20. <https://doi.org/10.1161/circulationaha.109.192627>.
- [133] Turton JL, Raab R, Rooney KB. Low-carbohydrate diets for type 1 diabetes mellitus: a systematic review. *PLoS One* 2018;13(3):e0194987. <https://doi.org/10.1371/journal.pone.0194987>.
- [134] Bolla AM, Caretto A, Laurenzi A, Scavini M, Piemonti L. Low-carb and ketogenic diets in type 1 and type 2 diabetes. *Nutrients* 2019;11(5). <https://doi.org/10.3390/nu11050962>.
- [135] Hammarstrom A, Wiklund AF, Lindahl B, Larsson C, Ahlgren C. Experiences of barriers and facilitators to weight-loss in a diet intervention – a qualitative study of women in northern Sweden. *BMC Womens Health* 2014;14:59. <https://doi.org/10.1186/1472-6874-14-59>.
- [136] Munt AE, Partridge SR, Allman-Farinelli M. The barriers and enablers of healthy eating among young adults: a missing piece of the obesity puzzle: a scoping review. *Obes Rev* 2017;18(1):1–17. <https://doi.org/10.1111/obr.12472>.
- [137] Bailey RL, Fulgoni VL, Cowan AE, Gaine PC. Sources of added sugars in young children, adolescents, and adults with low and high intakes of added sugars. *Nutrients* 2018;10(1). <https://doi.org/10.3390/nu10010102>.
- [138] Kirkpatrick SI, Raffoul A, Maynard M, Lee KM, Stapleton J. Gaps in the evidence on population interventions to reduce consumption of sugars: a review of reviews. *Nutrients* 2018;10(8). <https://doi.org/10.3390/nu10081036>.
- [139] Abdel Rahman A, Jomaa L, Kahale LA, Adair P, Pine C. Effectiveness of behavioral interventions to reduce the intake of sugar-sweetened beverages in children and adolescents: a systematic review and meta-analysis. *Nutr Rev* 2018;76(2):88–107. <https://doi.org/10.1093/nutrit/nux061>.
- [140] Cabrera Escobar MA, Veerman JL, Tollman SM, Bertram MY, Hofman KJ. Evidence that a tax on sugar sweetened beverages reduces the obesity rate: a meta-analysis. *BMC Public Health* 2013;13:1072. <https://doi.org/10.1186/1471-2458-13-1072>.
- [141] Linardon J, Mitchell S. Rigid dietary control, flexible dietary control, and intuitive eating: evidence for their differential relationship to disordered eating and body image concerns. *Eat Behav* 2017;26:16–22. <https://doi.org/10.1016/j.eatbeh.2017.01.008>.
- [142] Flack KD, Uffholz K, Casperson S, Jahns L, Johnson L, Roemmich JN. Decreasing the consumption of foods with sugar increases their reinforcing value: a potential barrier for dietary behavior change. *J Acad Nutr Diet* 2019;119(7):1099–108. <https://doi.org/10.1016/j.jand.2018.12.016>.