



# High nitrous oxide fluxes from rice indicate the need to manage water for both long- and short-term climate impacts

Kritee Kritee<sup>a,1</sup>, Drishya Nair<sup>b,c</sup>, Daniel Zavala-Araiza<sup>a</sup>, Jeremy Proville<sup>a</sup>, Joseph Rudek<sup>a</sup>, Tapan K. Adhya<sup>b,d</sup>, Terrance Loecke<sup>e</sup>, Tashina Esteves<sup>b,c</sup>, Shalini Balireddygarif, Obulapathi Dava<sup>f</sup>, Karthik Ram<sup>g</sup>, Abhilash S. R.<sup>9</sup>, Murugan Madasamy<sup>h</sup>, Ramakrishna V. Dokka<sup>i</sup>, Daniel Anandaraj<sup>h</sup>, D. Athiyaman<sup>9</sup>, Malla Reddy<sup>f</sup>, Richie Ahuja<sup>a</sup>, and Steven P. Hamburg<sup>a</sup>

<sup>a</sup>Environmental Defense Fund, New York, NY 10010; <sup>b</sup>Environmental Defense Fund, New Delhi 110001, India; <sup>c</sup>Fair Climate Network, Bagepalli 561 207, Karnataka, India; <sup>d</sup>School of Biotechnology, Kalinga Institute of Industrial Technology, Bhubaneswar 751 024, Odisha, India; <sup>e</sup>University of Kansas, Lawrence, KS 66045; <sup>f</sup>Accion Fraternal Ecology Centre, Anantapur 515 003, Andhra Pradesh, India; <sup>g</sup>Bharath Environment Seva Team, Pudukkottai 622 101, Tamil Nadu, India; <sup>h</sup>Palmyrah Workers Development Society, Tirunelveli 627 452, Tamil Nadu, India; and <sup>i</sup>Timbaktu Collective, Chennekothapalli 515 101, Andhra Pradesh, India

Edited by Paul G. Falkowski, Rutgers, The State University of New Jersey, New Brunswick, NJ, and approved August 3, 2018 (received for review June 11, 2018)

Global rice cultivation is estimated to account for 2.5% of current anthropogenic warming because of emissions of methane (CH<sub>4</sub>), a short-lived greenhouse gas. This estimate assumes a widespread prevalence of continuous flooding of most rice fields and hence does not include emissions of nitrous oxide (N<sub>2</sub>O), a long-lived greenhouse gas. Based on the belief that minimizing CH<sub>4</sub> from rice cultivation is always climate beneficial, current mitigation policies promote increased use of intermittent flooding. However, results from five intermittently flooded rice farms across three agroecological regions in India indicate that N<sub>2</sub>O emissions per hectare can be three times higher (33 kg-N<sub>2</sub>O·ha<sup>-1</sup>·season<sup>-1</sup>) than the maximum previously reported. Correlations between N<sub>2</sub>O emissions and management parameters suggest that N<sub>2</sub>O emissions from rice across the Indian subcontinent might be 30–45 times higher under intensified use of intermittent flooding than under continuous flooding. Our data further indicate that comanagement of water with inorganic nitrogen and/or organic matter inputs can decrease climate impacts caused by greenhouse gas emissions up to 90% and nitrogen management might not be central to N<sub>2</sub>O reduction. An understanding of climate benefits/drawbacks over time of different flooding regimes because of differences in N<sub>2</sub>O and CH<sub>4</sub> emissions can help select the most climate-friendly water management regimes for a given area. Region-specific studies of rice farming practices that map flooding regimes and measure effects of multiple comanaged variables on N<sub>2</sub>O and CH<sub>4</sub> emissions are necessary to determine and minimize the climate impacts of rice cultivation over both the short term and long term.

rice | nitrous oxide | methane | alternate wetting and drying | water

Rice (*Oryza sativa*) is a staple for nearly one-half of the world's seven billion people (1) and thus deserves special attention with respect to interactions with a changing climate. Rice farming provides a livelihood to ~145 million households (1), who in turn utilize for 11% of arable land, one-third of irrigation water (1), and at least one-seventh of fertilizers globally (2). Rice cultivation results in enhanced methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) emissions (hereafter, rice-CH<sub>4</sub> and rice-N<sub>2</sub>O, respectively), both potent greenhouse gases (GHGs) that contribute to climate change.

Rice cultivation is currently estimated to emit ~36 MMT CH<sub>4</sub> and contribute 2.5% (~0.1 W·m<sup>-2</sup>) to radiative forcing (3–7). These climate impacts of rice-CH<sub>4</sub> are projected to double by 2100 (8). Nitrous oxide (N<sub>2</sub>O) traps more heat over all time frames compared with CH<sub>4</sub> on a weight basis [100-y global warming potential (GWP<sub>100</sub>) of 298 vs. 34; GWP<sub>20</sub> of 268 vs. 86] and has a longer atmospheric lifetime (121 vs. 12 y) (9). While recent scientific research recognizes that rice-N<sub>2</sub>O needs to be addressed (3, 7, 10–12), policies on climate impacts of rice continue to assume that rice-N<sub>2</sub>O is negligible or small at <10%

of the total CO<sub>2</sub>e<sub>100y</sub> even under intermittently flooded conditions (13–15). None of the major rice-producing countries, including the two leading rice producers, China and India (16, 17), officially report rice-N<sub>2</sub>O or related emission factors in their national GHG inventories submitted to the United Nations (3). Crucially, most policy recommendations on rice management that include consideration of climate impacts focus on reducing rice-CH<sub>4</sub> by alternate wetting and drying (AWD), also called intermittent flooding. Water levels during intermittent flooding are typically allowed to fall to 15 cm below the soil surface before another round of irrigation (13–15). The only notable global policy guidance document to recognize rice-N<sub>2</sub>O is a recent modeling-based report (18), which suggested that, globally, neglecting contribution of soil carbon, rice-N<sub>2</sub>O contributes 25% to the GHG impact of rice cultivation on a CO<sub>2</sub>e<sub>100y</sub> basis (9).

Many factors including redox, bioavailable N, and organic C affect the extent of N<sub>2</sub>O formation that occurs primarily due to microbial nitrification–denitrification. Most research done to capture rice-N<sub>2</sub>O to date has been performed at farms with

## Significance

Methane from global rice cultivation currently accounts for one-half of all crop-related greenhouse gas emissions. Several international organizations are advocating reductions in methane emissions from rice by promoting intermittent flooding without accounting for the possibility of large emissions of nitrous oxide (N<sub>2</sub>O), a long-lived greenhouse gas. Our experimental results suggest that the Indian subcontinent's N<sub>2</sub>O emissions from intermittently flooded rice fields could be 30–45 times higher than reported under continuous flooding. Net climate impacts of rice cultivation could be reduced by up to 90% through comanagement of water, nitrogen, and carbon. To do this effectively will require a careful ongoing global assessment of N<sub>2</sub>O emissions from rice, or we will risk ignoring a very large source of climate impact.

Author contributions: K.K., T.K.A., T.L., D. Anandaraj, D. Athiyaman, M.R., and R.A. designed research; D.N., S.B., O.D., K.R., A.S.R., M.M., and R.V.D. performed research; K.K., D.Z.-A., and J.P. contributed new reagents/analytic tools; K.K., D.N., D.Z.-A., J.P., T.L., T.E., S.B., O.D., K.R., and A.S.R. analyzed data; and K.K., D.Z.-A., J.P., J.R., R.A., and S.P.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

<sup>1</sup>To whom correspondence should be addressed. Email: kritee@edf.org.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1809276115/-/DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1809276115/-/DCSupplemental).

Published online September 10, 2018.

continuous or mild-intermittent flooding under the assumption that these flooding regimes are representative of most rice cultivation, given their weed and pest control benefits (1). Under continuous flooding, redox conditions are conducive for methanogenesis, but not ideal for formation of N<sub>2</sub>O. Midseason drainage (a form of mild-intermittent flooding that causes a single long aeration event) brings redox conditions to levels that limit methanogenesis but are still lower than suitable for large amounts of N<sub>2</sub>O formation (19, 20). However, more intensely intermittent flooding includes multiple aeration events (i.e., drying and wetting of soils) leading to higher pulsed microbial activity, enhanced mineralization and nitrification–denitrification, as well as more redox cycles (21–23). All of these shifts increase the potential for high N<sub>2</sub>O emissions (21–23). Such multiple aeration events are common in both irrigated and rainfed rice farms in India, Pakistan, Nepal, Bangladesh, China, and South America as a result of high evapo-transpiration rates, unreliable water/electricity supply, rainfall regimes, soil characteristics, and topography (see *SI Appendix, SI Text, section 1* for details). However, very few studies have examined intermittently flooded rice farms, especially at a sampling intensity sufficient to accurately capture the high temporal variability in N<sub>2</sub>O fluxes. About 40 recent Indian studies on rice-GHG measured emissions on an average of 12% days in a rice growing season, a potentially insufficient number to accurately characterize N<sub>2</sub>O fluxes (*Dataset S1, Table 1*).

Our goal in the research reported here was to intensively study GHG emissions at rice farms that conventionally deploy a range of noncontinuous flooding regimes. We hypothesized that comanagement of N, water, and/or organic matter (OM) will result in a reduction in net climate impacts of rice. We made measurements at five farmer-managed farms (*Dataset S1, Table 2*) across three agroecological regions in India between 2012 and 2014 with sampling on 35–65% of days per season. We compared rice GHG emissions from two broad categories of treatments. “Baseline” practices (BPs) were identified via surveys of conventional farmers (*Dataset S1, Table 3*). Farm-specific “alternate” practices (APs) were potential climate-smart farming

practices, included completely organic practices at two farms and were identified by complex stakeholder processes (Table 1, *SI Appendix, SI Text, section 1*, and Fig. S1, and *Dataset S1, Tables 1–29*). Recommendations for alternate treatments included shifting flooding regimes closer to mild-intermittent flooding compared with BPs. However, farmers managed irrigation and it was only monitored unlike other management parameters that were both managed and monitored. There were a total of 13 treatments, with three replicates each, from the five farms (*Methods* and *SI Appendix, SI Text, sections 1–4*). We also examined potential correlations of N<sub>2</sub>O and CH<sub>4</sub> with 25 parameters including temperature characteristics, several water use variables, organic/inorganic inputs, soil organic carbon (24), texture (25), pH, and electrical conductivity.

### High N<sub>2</sub>O Fluxes

Fluxes of N<sub>2</sub>O at our farms varied from 0 to 33 kg N<sub>2</sub>O·ha<sup>-1</sup>·season<sup>-1</sup> (*Dataset S1, Table 30*) and –200 to 15,000 μg N<sub>2</sub>O·m<sup>-2</sup>·h<sup>-1</sup> among replicates from different treatments (*SI Appendix, Figs. S3–S8*). Our highest seasonal or hourly N<sub>2</sub>O fluxes are ~325–700% higher than the maximum previously reported rice-N<sub>2</sub>O fluxes (~10 kg N<sub>2</sub>O·ha<sup>-1</sup>·season<sup>-1</sup> and 2,100 μg·m<sup>-2</sup>·h<sup>-1</sup> measured in Italy under intermittent flooding conditions that were similar to our mild-intermittent regimes) (12). Depending on the mineralization rates of added OM, the proportion of applied N converted to N<sub>2</sub>O could be as high as 15–30% (*Dataset S1, Tables 24–30*), or 1–2 orders of magnitude greater than previously reported (10–12, 19, 26–28). The range of rice-CH<sub>4</sub> varied from 1 to 340 kg·ha<sup>-1</sup>·season<sup>-1</sup> (*SI Appendix, SI Text, section 6*, and *Dataset S1, Table 31*). When expressed in terms of long-term climate impacts, the contribution of N<sub>2</sub>O to net CO<sub>2</sub>e<sub>100</sub> ranged from zero to as high as 99% with a mean of ~35% (*Dataset S1, Table 32*).

Rice-N<sub>2</sub>O measured in our study is high for farms (Table 1, Fig. 1, and *SI Appendix, Fig. S1*) that underwent multiple aeration events as a result of fluctuating water levels and low

**Table 1. Farm-specific baseline (business as usual), APs, and GHG emissions**

Farm/year and treatment	Inorganic nitrogen,* kg·ha <sup>-1</sup>	Carbon input,† t·ha <sup>-1</sup>	Water index,‡ cm	Flood events§	Intermittent flooding regime¶	N <sub>2</sub> O, kg·ha <sup>-1</sup>	CH <sub>4</sub> , kg·ha <sup>-1</sup>	Yield, t·ha <sup>-1</sup>
Agroecological region# 3.0 (seed variety BPT 5204)								
Farm 1 2012								
Baseline	91	3.9–4.5	–555 (85)	1	Medium	13.1 (6.03)	66.5 (38.4)	4.8
Alternate	0	4.1–4.8	–580 (144)	1	Medium	4.7 (1.53)	81.1 (69.7)	4.6
Farm 2 2013								
Baseline	243	5.6–6.8	–0.7 (33)	3	Mild	0.62 (0.47)	105 (7.23)	4.8
Alternate	0	8.4–10.0	–152 (16)	3	Mild	0.10 (0.20)	98.3 (74.5)	2.7
Agroecological Region# 8.3 (seed variety ADT 39)								
Farm 3 2012								
Baseline	219	0.0–0.0	–486 (10)	0	Medium	22.7 (7.47)	3.98 (4.89)	4.2
Alternate	61	2.7–3.7	–416 (81)	0	Medium	2.51 (0.69)	4.6 (0.39)	2.7
Farm 3 2013								
Baseline	202	0.6–0.8	–1,036 (16)	3	Intense	17.4 (15.4)	108 (11.2)	5.6
Alternate	20	2.5–3.0	–858 (52)	3	Intense	11.5 (9.55)	112 (33.9)	4.0
Farm 4 2014								
Baseline	174	1.0–1.2	–212 (63)	3	Mild/medium	0.88 (0.83)	141 (19.3)	3.5
Alternate	91	1.1–1.4	–316 (147)	5	Mild/medium	0.02 (0.2)	154 (54.3)	3.2
Agroecological Region# 8.1 (seed variety ASD 16)								
Farm 5 2013								
Baseline	121	0.0–0.0	15 (65)	3	Mild	1.39 (1.66)	286 (49.1)	6.5
Alternate	99	0.01–0.02	–155 (91)	4	Mild	2.47 (1.16)	216 (88.1)	6.5

All errors in parentheses represent the ±95% confidence intervals ( $n = 3$ ).

\*The ranges for mineralized organic nitrogen and emission factors for each replicate are presented in *Dataset S1, Table 30*.

†Organic C content range as estimated via literature review (*Dataset S1, Table 4–9*).


‡Cumulative extent of flooding as determined by FWTs (*SI Appendix, SI Text, section 3*).

§Number of times a replicate had flooding for >3 d.

¶*SI Appendix, Fig. S1* presents our definitions of flooding regimes.

#See *SI Appendix, Fig. S2*, for a regional map.

||The methane flame ionization detector behaved anomalously in this cropping season, likely causing unusually low methane emissions.



**Fig. 1.** Average N<sub>2</sub>O and CH<sub>4</sub> fluxes. The GWP of N<sub>2</sub>O is three and nine times higher than CH<sub>4</sub> over 20 and 100 y, respectively. Therefore, the climate impacts of N<sub>2</sub>O are more dominant than those of CH<sub>4</sub> in the longer term (i.e., 100 vs. 20 y). The error bars represent the ±95% confidence interval.

cumulative flooding as observed using field-water tubes (*Methods*, *SI Appendix*, *SI Text*, section 3, and Figs. S3–S14, and *Dataset S1*, Table 32). Our high-intensity sampling allowed us to see “delayed” N<sub>2</sub>O peaks even 30 d after N addition (*SI Appendix*, *SI Text*, section 4) potentially caused by N made bioavailable via mineralization through successive aeration events.

### Parameters Affecting Rice-N<sub>2</sub>O

When individual management and soil characteristics were considered, rice-N<sub>2</sub>O was positively correlated with added inorganic N and soil texture, and negatively correlated with extent of flooding and added OM (two variables usually positively correlated with rice-CH<sub>4</sub>) (11, 27) (*SI Appendix*, Figs. S17–S22 and *Dataset S1*, Tables 32 and 34). However, the following multiple-regression model explained most of the observed variability in seasonal rice-N<sub>2</sub>O (*P* value < 0.001, adjusted *R*<sup>2</sup> = 0.80; *SI Appendix*, *SI Text*, section 5, and Fig. S32):

$$\begin{aligned} \text{N}_2\text{O} = & -0.01 * (\text{water index}) - 0.91 * (\text{flood events}_{>3 \text{ days}}) \\ & + 0.02 * \text{N}_{\text{inorganic}} + \epsilon_1. \end{aligned} \quad [1]$$

Here, N<sub>2</sub>O represents emissions in kilograms-N-hectare<sup>-1</sup>·season<sup>-1</sup>, flood events<sub>>3 d</sub> is the number of times a plot had flooding (>0-cm water level) for >3 d, N<sub>inorganic</sub> is inorganic N input in kilograms-hectare<sup>-1</sup>, and  $\epsilon_1$  is statistical error (*SI Appendix*, Fig. S29 and *Dataset S1*, Table 35). Water index, a measure of cumulative extent of flooding and the sum of daily water levels in a vertical field water tube (FWT), emerged as the most significant predictor of N<sub>2</sub>O. Flood events<sub>>3 d</sub>, another water use variable, described the number of multiple aeration events for a given water index. When there were frequent long (>3 d) flood events but lesser short (<3 d) flood events, there was a reduction in aeration events and rice-N<sub>2</sub>O. The variable flood events<sub>>3d</sub> is noncorrelated with water index (*SI Appendix*, Fig. S23). Given the focal importance of water management regimes to rice-N<sub>2</sub>O, we are introducing definitions of mild-, medium-, and intense-intermittent flooding regimes based on the ranges of water indices and number of flood events in *SI Appendix*, Fig. S1.

Data from individual farms clearly indicate that OM addition suppresses and/or delays the emergence of a N<sub>2</sub>O peak despite low water index (*SI Appendix*, *SI Text*, section 5, and Fig. S20 and *Dataset S1*, Table 30). In addition, many N<sub>2</sub>O and CH<sub>4</sub> peaks were associated with drainage events (*Dataset S1*, Tables 30 and 31), but N<sub>2</sub>O flux at some farms with high OM inputs did not increase with drainage. However, added OM was not included in our final model because it did not add any additional statistical power to the best-fit multiple regression model (*Methods*). Organic inputs are known to decrease N<sub>2</sub>O flux for both rice and nonrice farms under N-limitation by delaying mineralization of mineral-N when the C/N ratio of OM is high, improving either N-incorporation in microbial biomass or promoting conversion of N<sub>2</sub>O to N<sub>2</sub> (29–31).

### The Risk of Enhanced Rice-N<sub>2</sub>O in the Indian Subcontinent

Because intermittent flooding is being actively promoted to reduce rice-CH<sub>4</sub> through policy frameworks at national and international levels (13–15), our research should be replicated in other regions to determine the implications of our findings on the potential magnitude of global rice-N<sub>2</sub>O. While extrapolation of region-specific findings to additional agroecological regions should be done with caution (*SI Appendix*, *SI Text*, section 8), we examine the potential implications of policies which ignore large rice-N<sub>2</sub>O emissions from intermittently flooded farms on the Indian subcontinent.

We investigated potential rice-N<sub>2</sub>O by exploring the impact of deploying three hypothetical flooding scenarios (continuous, medium-intermittent, and intense-intermittent flooding for irrigated farms; *SI Appendix*, Fig. S1) on the Indian subcontinent using our multiple-regression model (Eq. 1). We explored the climate implications among 12 classes of water management regimes in the subcontinent (*SI Appendix*, Fig. S36) (32) using spatially explicit data detailing rice-specific inorganic fertilizer use (33). *Dataset S1*, Table 38 presents water index and flooding events<sub>>3 d</sub> assumptions for each management class and each flooding scenario.

As expected, our results suggest that rainfed and upland farms are at risk for elevated rice-N<sub>2</sub>O, while deepwater and wetland rice cropping systems are much less susceptible to such emissions (Fig. 2). Two recent modeling studies of India suggest emissions of 18,000 tons N<sub>2</sub>O·y<sup>-1</sup>, assuming 90% of rice production is under continuous flooding (34), and 250,000 tons N<sub>2</sub>O·y<sup>-1</sup>, assuming 70% is under midseason drainage (18). When we use the same rate of N addition (69 kg N·ha<sup>-1</sup>) and similar water management (i.e., mild-intermittent flooding; *Dataset S1*, Table 38) as used by the earlier model-based study (18), our model suggests Indian rice-N<sub>2</sub>O at ~230,000 tons N<sub>2</sub>O·y<sup>-1</sup> close to the estimate of 250,000 tons N<sub>2</sub>O·y<sup>-1</sup> under midseason drainage. However, under medium- or intense-intermittent flooding regimes, which are more common than previously acknowledged and might be becoming more frequent due to water stress and AWD guidelines, our model predicts a higher range of 530,000–790,000 tons N<sub>2</sub>O·y<sup>-1</sup> for rice-N<sub>2</sub>O in India (*Methods* and *Dataset S1*, Tables 12 and 13). Similarly, our estimates of rice-N<sub>2</sub>O for the entire Indian subcontinent under more intensely intermittent flooding conditions are 1.5–2 times higher than under mild-intermittent flooding (18) and 30–45 times higher than under continuous flooding (34) (*Dataset S1*, Tables 39 and 40). Rice-N<sub>2</sub>O from the Indian subcontinent according to our model is higher than previously reported as a result of (i) high N<sub>2</sub>O fluxes under intensely intermittent flooding, (ii) higher number of water management classes (32) that assume intense forms of intermittent flooding compared with an assumption of continuous flooding (34) or mid-season drainage (18), and (iii) a higher and geospatially variable inorganic N addition rate of 102 ± 48 (SD) kg N·ha<sup>-1</sup> based on more up-to-date data (33) compared with a fixed quantity of 69 kg N·ha<sup>-1</sup> (18). Even without any geospatial modeling, the emission factors for intermittently flooded farms developed in this study




Fig. 2. Qualitative risk of elevated  $N_2O$  emissions from the Indian subcontinent under three flooding scenarios: continuous (A), medium-intermittent (B), and intense-intermittent (C) flooding. The maps depicts rice growing areas in India, Nepal, Pakistan, Sri Lanka, and Bangladesh across 12 water management regimes. For assumed water indices and flooding events, see Dataset S1, Table 38. For a quantitative assessment of emissions, see Dataset S1, Tables 41 and 42.

suggest that net Indian rice- $N_2O$  and rice- $CH_4$  emissions are equivalent to  $\sim 245$  million  $tCO_2e_{100}y^{-1}$ , more than two times higher than previous estimates (17, 35) (SI Appendix, SI Text, section 10).

Given the International Rice Research Institute's latest global estimate that  $\sim 60\%$  of global rice area is irrigated (36) and thus susceptible to high rice- $N_2O$  under intensely intermittent flooding regimes, there is a need for further research to fully understand the net climate benefits of promoting intermittent flooding for short-term climate mitigation.

#### Parameters Affecting Rice- $CH_4$

In contrast to rice- $N_2O$ , rice- $CH_4$  was positively correlated with parameters that reflect flooding extent and amount of soil OM (Dataset S1, Table 36), consistent with past findings that the lowest  $CH_4$  fluxes are recorded on farms with multiple aeration events and poor soils (37). The following best-fit model explained our seasonal rice- $CH_4$  data ( $P$  value  $< 0.001$ , adjusted  $R^2 = 0.91$ ):

$$CH_4 = 34 * (\text{flood events}_{>3d}) + 88 * \text{SOM} + \epsilon_2. \quad [2]$$

Here,  $CH_4$  represents emissions in kilograms  $CH_4 \cdot \text{hectare}^{-1} \cdot \text{season}^{-1}$ ,  $\text{flood events}_{>3d}$  is the number of times a plot had  $>0$ -cm water level for  $>3$  d, SOM is soil OM in percentage, and  $\epsilon_2$  is statistical error (SI Appendix, Fig. S30 and Dataset S1, Table 37). Unlike SOM, we did not observe a consistently positive correlation of rice- $CH_4$  with organic inputs corroborating previous studies on intermittently flooded farms (27) (SI Appendix, SI Text, section 6, and Fig. S28).

#### Mitigation Potential of APs

Compared with the baseline plots at the same farms, management of multiple parameters at alternate plots shows average mitigation of up to  $70 \text{ kg } CH_4 \cdot \text{ha}^{-1}$  ( $2.4 \text{ tCO}_2e_{100} \cdot \text{ha}^{-1}$ ) and up to  $20 \text{ kg } N_2O \cdot \text{ha}^{-1}$  ( $6 \text{ tCO}_2e_{100} \cdot \text{ha}^{-1}$ ) (SI Appendix, SI Text, section 7, and Figs. S31–S35 and Dataset S1, Table 33). The range of rice- $CH_4$  mitigation observed ( $-0.50$ – $2.4 \text{ tCO}_2e_{100} \cdot \text{ha}^{-1} \cdot \text{season}^{-1}$ ; Dataset S1, Table 33) is similar to the potential noted by the Intergovernmental Panel on Climate Change (IPCC) ( $-0.55$ – $2.8 \text{ tCO}_2e_{100} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ ) (3). However, the IPCC suggests, without specifying how this range might be different for rice vs. nonrice farms, that fertilizer management leads to a smaller and narrower range of  $N_2O$  mitigation ( $0.01$ – $0.32 \text{ tCO}_2e_{100} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ ) relative to what we observed ( $-0.3$ – $6 \text{ tCO}_2e_{100} \cdot \text{ha}^{-1} \cdot \text{season}^{-1}$ ) (3).

An analysis based on Eqs. 1 and 2 shows that when water management shifts from continuous to mild-intermittent flooding and N is reduced from  $250$  to  $150 \text{ kg} \cdot \text{ha}^{-1}$ , a  $60\%$  reduction in net climate impacts can be achieved (Dataset S1, Tables 39 and 40). Compared with BPs, APs provided a  $10$ – $90\%$  ( $0.4$ – $6.0$


$tCO_2e_{100} \cdot \text{ha}^{-1} \cdot \text{season}^{-1}$ ) net reduction in climate impacts for five out of six seasons with a small increase in net climate impacts in the sixth season examined (Dataset S1, Table 33). Many of the APs examined in this study produced significantly lower yields than our BPs, but reduction in yields is not correlated with reduction in net climate impacts (Table 1 and SI Appendix, Fig. S38). More research is required to optimize inorganic N, OM, or water inputs such that climate impacts per unit yield are minimized.

Notably, existing AWD-based guidelines to mitigate climate impacts of rice assume that rice- $N_2O$  can be controlled primarily by efficient fertilizer use (3, 15). Our data, however, show that reducing fertilizer use might not be central to managing rice- $N_2O$  (SI Appendix, Figs. S18, S21, and S24). Our model suggests that, as the extent of intermittent flooding increases (i.e., water index and flood events $_{>3d}$  decrease), the contribution of fertilizer-N to  $N_2O$  decreases (Eq. 1, Compare Dataset S1, Tables 39 and 40). In farms with very high N use, reducing N bioavailability by decreasing N or increasing OM use will still be crucial to reducing rice- $N_2O$  (SI Appendix, SI Text, section 7). Previous work shows that addition of N right before prolonged flooding can significantly reduce rice- $N_2O$  (38), but the prolonged flooding option is not easily available in water-stressed areas. With respect to OM addition, recommendations are frequently based on the well-documented impact of OM on rice- $CH_4$  under continuous flooding (27). Our results provide a basis for developing OM management recommendations to limit rice- $N_2O$  under intermittently flooded conditions (SI Appendix, SI Text, section 7).

#### Climate Impacts of Rice- $N_2O$ and Rice- $CH_4$ over Time

An updated assessment of net climate impacts of water management at rice farms is required as rice- $N_2O$  can be higher than previously assumed and with an overall trend of an inverse relationship between rice- $CH_4$  and rice- $N_2O$  (SI Appendix, Figs. S15 and S16). Because the climate impacts of  $CH_4$  and  $N_2O$  differ significantly over time, the goal of rice management should be to reduce net radiative forcing over both the long term and short term, instead of focusing on minimizing climate impacts over only the long term by reducing  $N_2O$  or only the short term by reducing  $CH_4$ . The standard practice of determining climate impacts among GHGs is through GWPs, which compare a given GHG against  $CO_2$  at a single arbitrarily selected time (e.g., 100 y). Reporting the implications of specific mitigation options over both the short-term GWP (20 y) and long-term GWP (100 y) gives a more complete picture of climate impacts.

Moving beyond the evaluation of climate impacts at two distinct times, the technology warming potential (TWP) framework (39) integrates GWPs over time and allows an easy way to visualize trade-offs between GHGs with different radiative forcing and residence times. Here, we extend the use of the TWP framework to rice cultivation. Fig. 3 presents the relative cumulative climate impacts of four hypothetical flooding regimes compared with a



**Fig. 3.** Temporal analysis of climate impacts of four hypothetical irrigated water management classes. Each water management regime is represented by a fixed water index and range of flood events<sub>>3 d</sub> and is presented relative to a fixed “base case” (continuous flooding, water index = 500, flood events<sub>>3 d</sub> = 6; represented by the red line). The ratios of cumulative radiative forcing relative to the base case are shown on the y axis, and continuous flooding regimes (red band; water index = 500, flood event 5–8) are compared with mild (blue band; water index = –100, flood events 2–6), medium (green band; water index = –600, flood events 0–5), and intense (purple band; water index = –1,200, flood events 0–3) intermittent regimes in A–C, respectively. The ratio of cumulative radiative forcing values below 1 (red line) represent climate benefit relative to the fixed base case with the width of the shaded regions reflecting the variability in climate impacts for a given water index depending on the number of flood events. The lowest number of flood events are at the lower band edge, and the highest number of flood events at the top edge, because the less flood events<sub>>3 d</sub> cause net lower GWP (Eqs. 1 and 2). These ratios of cumulative radiative forcings change with time on x axis. Intense intermittent regimes cross over and have more cumulative climate impact than our base case within 60–100 y. Medium intermittent regimes with many flood events<sub>>3 d</sub> could cross over as early as 40 y. However, medium intermittent scenarios with very few flood events<sub>>3 d</sub> or mild intermittent scenarios might never have more climate impact than the chosen base case.

continuous-flooding “base case,” assuming a constant and continuous flux of both N<sub>2</sub>O and CH<sub>4</sub> for 200 y (*SI Appendix, SI Text, section 9*, and *Dataset S1, Tables 39 and 40*). For each flooding regime, the climate impacts of N<sub>2</sub>O continue to add to the long-term radiative forcing because it is a long-term climate forcer as opposed to CH<sub>4</sub> whose climate impacts are predominately in the short term.

The extent of climate impacts of different flooding regimes compared with the base case of continuous flooding varies over time and with water management. The comparison of continuous flooding regimes with different intermittent flooding regimes shows that, in general, relatively shallow (e.g., mild-intermittent, water index ~ –100) flooding can reduce the long- and short-term climate impacts of rice cultivation compared with continuous flooding regimes (Fig. 3A). At lower water indices (Fig. 3B and C), however, the climate impacts of reducing CH<sub>4</sub> through water management could be more than offset by N<sub>2</sub>O fluxes within 30 y, especially if the number of flood-events<sub>>3 d</sub> are high.

Regardless of the relative importance of water, nitrogen, and carbon in impacting rice-N<sub>2</sub>O, a temporal analysis of management options for each region can be a powerful tool to visualize climate impacts over both the short term and long term.

## Implications

**Intensive Mapping of Flooding Regimes and Measurement of Rice-N<sub>2</sub>O Is Critical.** Our empirical data show high N<sub>2</sub>O fluxes at medium- and intense-intermittently flooded rice farms, and extrapolation of these observations suggests that many, but not all, rice-growing regions in the Indian subcontinent (and potentially globally) could potentially be experiencing significant rice-N<sub>2</sub>O and concomitant climate impacts (Figs. 1 and 2 and *Dataset S1, Tables 41, 42, and 44*). Increasing pressure on limited water resources, AWD water management, and a changing climate (i.e., higher temperatures and evapo-transpiration rates) could make additional regions susceptible to high N<sub>2</sub>O fluxes. Thus, if we are to understand the climate implications and realistic mitigation potential of climate-smart rice production practices, it is important that rice-N<sub>2</sub>O be intensively measured (*Dataset S1, Table 43*) along with the mapping of actual flooding regimes. We expect rice-N<sub>2</sub>O to be significantly higher than present estimates.

**AWD Is Not Always Climate Beneficial, Especially in the Long Term.** While multiple parameters including carbon and fertilizer use influenced GHG emissions, flooding regimes emerged as the

strongest predictor of the net climate impacts of farm-specific BPs and APs in our study (Eqs. 1 and 2). Two key strategies often proposed to reduce rice-CH<sub>4</sub> [i.e., limiting water and C input (11, 40)] could stimulate N<sub>2</sub>O production (*SI Appendix, Figs. S17–S22 and Dataset S1, Tables 30 and 34*). It is crucial to understand under what conditions this disbenefit of water and C input reduction is important. The assumption by policymakers that AWD with some adjustments in fertilizer use will significantly reduce the net climate impact of rice farms will not always be true (Fig. 3). We need to intensify the study of farm-specific integrated management of inorganic N, OM, and water use with a focus on maximizing rice yields and farm profits while minimizing short- and long-term climate impacts (*SI Appendix, SI Text, section 11*). Based on these data, policies can be adopted that allow robust large-scale implementation of integrated climate-protecting and production-maximizing practices (11, 27, 38).

## Methods

**BPs and APs.** Both baseline and alternative treatments were farm and year specific (Table 1 and *SI Appendix, SI Text, sections 1 and 2*, and *Dataset S1, Tables 4 and 9*). BPs represented management practices implemented by the majority of conventional small-holder rice farmers in the previous year as determined by region-specific farmer surveys before the beginning of each season (*Dataset S1, Tables 3 and 10–23*). The surveys indicated that the farmers were using fertilizer at rates significantly different from those recommended by the local governments and/or academic institutions. The APs were chosen by a consortium of local agronomists, farmers, and nongovernmental organization partners as previously described and has been previously described (41).

**Measurement of GHG Emissions.** Samples collected through a modified manual chamber were analyzed by gas chromatograph to measure N<sub>2</sub>O and CH<sub>4</sub> on 35–65% of days in a growing season with an average minimum detection limit of 18 mg N<sub>2</sub>O·h<sup>–1</sup>·m<sup>–2</sup> and 37 mg CH<sub>4</sub>·h<sup>–1</sup>·m<sup>–2</sup> (*SI Appendix, SI Text, section 2*, and *Dataset S1, Tables 4–9 and 24–29*). The complete methodology including details of unique vertically stacked chambers, access bridges, and temperature and volume corrections is summarized in *SI Appendix, SI Text* (41).

**Water Index.** Water index is the sum of daily water levels (in centimeters) in a FWT in a growing season relative to the soil surface. Water levels were observed between 8 and 11 AM once a day (sampling intensity, 55–100% days in a season; *Dataset S1, Table 2*). The daily water levels represent a snapshot because they dropped quickly (4–15 cm within 24 h) after irrigation (*SI Appendix, SI Text, section 3*).

**Multiple Regression.** Each farm with a different treatment ( $n = 13$  treatments) was considered an independent observation, and the mean of each parameter (3 replicates) was used to represent each farm in the regression analysis. To select the “best-fit” multiple regression model for  $N_2O$  and  $CH_4$ , we looked to minimize the Akaike information criterion and checked for model significance after adding/removing parameters.

**Estimation of Rice- $N_2O$  Flux from Indian Subcontinent.** Multiple regression coefficients were extrapolated using spatially explicit datasets of rice-specific inorganic N fertilizer inputs (in kilograms per hectare) (33) and high-resolution rice management classes for the subcontinent (32). The N fertilizer dataset and the rice management classes were available at 5-arc-min (~10-km) (33) and 500-m (32) grid cell resolutions, respectively. For each class of rice irrigation management, we assigned a range of representative water index and flood event values. See *SI Appendix, SI Text, section 8*, and *Figs. S36 and S37* and *Dataset S1, Table 38* for details.

**TWPs.** A framework developed by Alvarez et al. (39) was used to calculate TWPs, which at each point in time represent the ratio of cumulative radiative forcing from two different management practices. The choice of the denominator (water index = 500; flood event = 6) is a benchmark against which all other management practices are compared. Our analysis assumes

that both climate pollutants ( $N_2O$  and  $CH_4$ ) are emitted continuously and indefinitely at a constant rate  $E_{i,j}$  for 200 y. Thus, the TWP used to compare  $CH_4$  and  $N_2O$  emissions from two management practices is expressed as follows:

$$TWP = \frac{E_{1,CH_4}TRF_{CH_4}(t) + E_{1,N_2O}TRF_{N_2O}(t)}{E_{2,CH_4}TRF_{CH_4}(t) + E_{2,N_2O}TRF_{N_2O}(t)}$$

where  $E_{i,j}$  represents the emission rate (in kilograms per hectare) of climate pollutant  $j$  from management practice  $i$ , and  $TRF_j(t)$  represents the total radiative forcing values of each pollutant  $j$ . The selection of management practices scenarios and the estimation of emission rates is presented in *Dataset S1, Table 39*. The derivation of  $TRF_j(t)$  values is provided in *SI Appendix, SI Text, section 9*.

**ACKNOWLEDGMENTS.** We thank two anonymous reviewers for their insightful comments and thorough suggestions. This work would not have been possible without collaboration with five rice farmers in peninsular India. The Institute on the Environment at University of Minnesota and the International Rice Research Institute helped us access crop-specific nitrous oxide emission factors from their study and geospatial information on rice management classes, respectively. This research was partially supported by funding from ICCO Foundation and Brot fuer die Welt.

1. Global Rice Science Partnership (2013) *Rice Almanac* (International Rice Research Institute, Los Baños, Philippines), 4th Ed.
2. Heffer P (2009) *Assessment of Fertilizer Use by Crop at the Global Level 2006–2007* (International Fertilizer Industry Association, Paris).
3. Smith P, et al. (2007) Agriculture. *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Metz B, Davidson OR, Bosch PR, Dave R, Meyer LA (Cambridge Univ Press, Cambridge, UK).
4. Ciais P, et al. (2014) Carbon and other biogeochemical cycles. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Stocker TF, et al. (Cambridge Univ Press, Cambridge, UK).
5. Kirschke S, et al. (2013) Three decades of global methane sources and sinks. *Nat Geosci* 6:813–823.
6. Turner A, et al. (2016) A large increase in US methane emissions over the past decade inferred from satellite data and surface observations. *Geophys Res Lett* 43:2218–2224.
7. Carlson KM, et al. (2017) Greenhouse gas emissions intensity of global croplands. *Nat Clim Change* 7:63–68.
8. van Groenigen KJ, Van Kessel C, Hungate BA (2013) Increased greenhouse-gas intensity of rice production under future atmospheric conditions. *Nat Clim Change* 3:288–291.
9. Myhre G, et al. (2013) Anthropogenic and natural radiative forcing. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds Stocker TF, et al. (Cambridge Univ Press, Cambridge, UK).
10. Li X, Yuan W, Xu H, Cai Z, Yagi K (2011) Effect of timing and duration of midseason aeration on  $CH_4$  and  $N_2O$  emissions from irrigated lowland rice paddies in China. *Nutr Cycling Agroecosyst* 91:293–305.
11. Linquist BA, Adviento-Borbe MA, Pittelkow CM, van Kessel C, van Groenigen KJ (2012) Fertilizer management practices and greenhouse gas emissions from rice systems: A quantitative review and analysis. *Field Crops Res* 135:10–21.
12. Lagomarsino A, et al. (2016) Alternate wetting and drying of rice reduced  $CH_4$  but triggered  $N_2O$  peaks in a clayey soil of central Italy. *Pedosphere* 26:533–548.
13. Climate and Clean Air Coalition (2014) *Methane Mitigation in Rice Paddies: A New CCAC Component. A Joint Announcement Brochure by CGIAR Research Program on Climate Change, Agriculture, and Food Security (CCAFS) and Climate and Clean Air Coalition (CCAC) to Reduce Short-Lived Climate Pollutants* (International Rice Research Institute, Los Baños, Philippines).
14. CCAFS (2017) Mitigation strategies in rice production, in collaboration with the Climate and Clean Air Coalition (CCAC). *Low Emissions Development Research Flagship, The CGIAR Research Program on Climate Change Agriculture Food Security (Wageningen University and Research, Wageningen, The Netherlands)*.
15. Richards M, Sander BO (2014) *Alternate Wetting and Drying in Irrigated Rice: Implementation Guide for Policymakers and Investors as Practice Brief on Climate Smart Agriculture* (CGIAR Research Program on Climate Change, Agriculture and Food Security, Copenhagen).
16. National Development and Reform Commission (2012) *Second National Communication on Climate Change of the People's Republic of China* (National Development and Reform Commission, Beijing).
17. Ministry of Environment, Forest and Climate Change (2012) *Government of India's Second National Communication to the United Nations Framework Convention on Climate Change* (Ministry of Environment, Forest and Climate Change, New Delhi).
18. Environmental Protection Agency (2013) *Global Mitigation of Non- $CO_2$  Greenhouse Gases: 2010–2030* (United States Environmental Protection Agency, Office of Atmospheric Programs, Washington, DC), pp 19–42.
19. Johnson-Beebout SE, Angeles OR, Alberto MCR, Buresh RJ (2009) Simultaneous minimization of nitrous oxide and methane emission from rice paddy soils is improbable due to redox potential changes with depth in a greenhouse experiment without plants. *Geoderma* 149:45–53.
20. Hou AX, Chen GX, Wang ZP, Van Cleemput O, Patrick WH (2000) Methane and nitrous oxide emissions from a rice field in relation to soil redox and microbiological processes. *Soil Sci Soc Am J* 64:2180–2186.
21. Lagomarsino A, et al. (2016) Past water management affected GHG production and microbial community pattern in Italian rice paddy soils. *Soil Biol Biochem* 93:17–27.
22. Liang LL, Grant DA, Jenerette GD (2016) Multivariate regulation of soil  $CO_2$  and  $N_2O$  pulse emissions from agricultural soils. *Glob Change Biol* 22:1286–1298.
23. Harrison-Kirk T, Beare MH, Meenken ED, Condon LM (2013) Soil organic matter and texture affect responses to dry/wet cycles: Effects on carbon dioxide and nitrous oxide emissions. *Soil Biol Biochem* 57:43–55.
24. Walkley A, Black IA (1934) An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci Soc Am J* 79:239–250.
25. Kettler TA, Doran JW, Gilbert TL (2001) Simplified method for soil particle-size determination to accompany soil-quality analyses. *Soil Sci Soc Am J* 65:849–852.
26. Sun W, Huang Y (2012) Synthetic fertilizer management for China's cereal crops has reduced  $N_2O$  emissions since the early 2000s. *Environ Pollut* 160:24–27.
27. Sanchis E, Ferrer M, Torres AG, Cambra-López M, Calvet S (2012) Effect of water and straw management practices on methane emissions from rice fields: A review through a meta-analysis. *Environ Eng Sci* 29:1053–1062.
28. Adviento-Borbe MA, et al. (2015) Methane and nitrous oxide emissions from flooded rice systems following the end-of-season drain. *J Environ Qual* 44:1071–1079.
29. Dalal RC, Gibson I, Allen DE, Menzies NW (2010) Green waste compost reduces nitrous oxide emissions from feedlot manure applied to soil. *Agric Ecosyst Environ* 136:273–281.
30. Iqbal J, Parkin TB, Helmers MJ, Zhou X, Castellano MJ (2015) Denitrification and nitrous oxide emissions in annual croplands, perennial grass buffers, and restored perennial grasslands. *Soil Sci Soc Am J* 79:239–250.
31. Zou J, Huang Y, Jiang J, Zheng X, Sass RL (2005) A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: Effects of water regime, crop residue, and fertilizer application. *Global Biogeochem Cycles* 19:GB2021.
32. Gumma MK, Nelson A, Thenkabail PS, Singh AN (2011) Mapping rice areas of South Asia using MODIS multitemporal data. *J Appl Remote Sens* 5:053547.
33. Mueller ND, et al. (2012) Closing yield gaps through nutrient and water management. *Nature* 490:254–257.
34. Gerber JS, et al. (2016) Spatially explicit estimates of  $N_2O$  emissions from croplands suggest climate mitigation opportunities from improved fertilizer management. *Glob Change Biol* 22:3383–3394.
35. Bhatia A, Jain N, Pathak H (2013) Methane and nitrous oxide emissions from Indian rice paddies, agricultural soils and crop residue burning. *Greenhouse Gas Sci Technol* 3:196–211.
36. International Rice Research Institute (2011) Global rice area: Data obtained from International Rice Research Institute's Social Sciences Division in January 2016, derived from World Rice Statistics. Available at [ricestat.irri.org:8080/wrs2/entrypoint.htm](http://ricestat.irri.org:8080/wrs2/entrypoint.htm). Accessed January 5, 2016.
37. Sass RL (2003)  $CH_4$  emissions from rice agriculture. *IPCC Expert Meetings on Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (Institute for Global Environmental Strategies/IPCC, Hayama, Japan), pp 399–417.
38. Linquist BA, et al. (2015) Reducing greenhouse gas emissions, water use, and grain arsenic levels in rice systems. *Glob Change Biol* 21:407–417.
39. Alvarez RA, Pacala SW, Winebrake JJ, Chameides WL, Hamburg SP (2012) Greater focus needed on methane leakage from natural gas infrastructure. *Proc Natl Acad Sci USA* 109:6435–6440.
40. Ortiz-Monasterio I, et al. (2010) Greenhouse gas mitigation in the main cereal systems: Rice, wheat and maize. *Climate Change and Crop Production*, ed Reynolds MP (CAB International, Oxfordshire, UK), Vol 13.
41. Kritee K, et al. (2015) Groundnut cultivation in semi-arid peninsular India for yield scaled nitrous oxide emission reduction. *Nutr Cycling Agroecosyst* 103:115–129.