



Research Paper

Impact of genus (*Geotrigona*, *Melipona*, *Scaptotrigona*) on the targeted ¹H-NMR organic profile, and authenticity test by interphase emulsion of honey processed in cerumen pots by stingless bees in Ecuador

Patricia Vit^{a,*}, Jane van der Meulen^b, Maria Diaz^b, Silvia R.M. Pedro^c, Isabelle Esperança^d, Rahimah Zakaria^e, Gudrun Beckh^b, Favian Maza^f, Gina Meccia^g, Michael S. Engel^{h,i}

^a Food Science Department, Faculty of Pharmacy and Bioanalysis, Universidad de Los Andes, Mérida, 5101, Venezuela

^b Quality Services International GmbH, 28199, Bremen, Germany

^c Biology Department, Universidade de São Paulo, Ribeirão Preto, Brazil

^d Institute of Chemistry, Universidad Federal de Rio de Janeiro, Rio de Janeiro, RJ, 21945970, Brazil

^e Department of Physiology, School of Medical Sciences, Universiti Sains Malaysia, Kota Bharu, Malaysia

^f Faculty of Agricultural and Livestock Sciences, Universidad Técnica de Machala, Machala, El Oro province, Ecuador

^g Research Institute, Faculty of Pharmacy and Bioanalysis, Universidad de Los Andes, Mérida 5101, Venezuela

^h Division of Entomology, Natural History Museum, Department of Ecology & Evolutionary Biology, 1501 Crestline Drive-Suite 140, University of Kansas, Lawrence, KS, USA

ⁱ Division of Invertebrate Zoology, American Museum of Natural History, Central Park West at 79th Street, New York, NY, 10024, USA



ARTICLE INFO

Keywords:

Geotrigona

HATIE

HBT

Melipona

Meliponini

Targeted ¹H-NMR

Pot-honey

Scaptotrigona

Stingless bees

ABSTRACT

The biodiversity of Ecuadorian stingless bees is almost 200 species. Traditional pot-honey harvest in Ecuador is mostly done from nests of the three genera selected here *Geotrigona* Moure, 1943, *Melipona* Illiger, 1806, and *Scaptotrigona* Moure, 1942. The 20 pot-honey samples collected from cerumen pots and three ethnic honeys “abeja de tierra”, “bermejo”, and “cushillomishki” were analyzed for qualitative and quantitative targeted ¹H-NMR honey profiling, and for the Honey Authenticity Test by Interphase Emulsion (HATIE). Extensive data of targeted organic compounds (41 parameters) were identified, quantified, and described. The three honey types were compared by ANOVA. Amino acids, ethanol, hydroxymethylfurfural, aliphatic organic acids, sugars, and markers of botanical origin. The number of phases observed with the HATIE were one in *Scaptotrigona* and three in *Geotrigona* and *Melipona* honeys. Acetic acid (19.60 ± 1.45 g/kg) and lactic acid (24.30 ± 1.65 g/kg) were particularly high in *Geotrigona* honey (in contrast to 1.3 g/kg acetic acid and 1.6 g/kg lactic acid in *Melipona* and *Scaptotrigona*), and with the lowest fructose + glucose (18.39 ± 1.68) g/100g honey compared to *Melipona* (52.87 ± 1.75) and *Scaptotrigona* (52.17 ± 0.60). Three local honeys were tested using PCA (Principal Component Analysis), two were assigned with a correct declared bee origin, but “bermejo” was not a *Melipona* and grouped with the *Scaptotrigona* cluster. However after HCA (Hierarchical Cluster Analysis) the three honeys were positioned in the *Melipona-Scaptotrigona* cluster. This research supports targeted ¹H-NMR-based profiling of pot-honey metabolomics approach for multi-parameter visualization of organic compounds, as well as descriptive and pertained multivariate statistics (HCA and PCA) to discriminate the stingless bee genus in a set of *Geotrigona*, *Melipona* and *Scaptotrigona* honey types. The NMR characterization of Ecuadorian honey produced by stingless bees emphasizes the need for regulatory norms. A final note on stingless bee markers in pot-honey metabolites which should be screened for those that may extract phylogenetic signals from nutritional traits of honey. *Scaptotrigona vitorum* honey revealed biosurfactant activity in the HATIE, originating a fingerprint Honey Biosurfactant Test (HBT) for the genus in this set of pot-honeys.

* Corresponding author.

E-mail address: vitolivier@gmail.com (P. Vit).

<https://doi.org/10.1016/j.crfs.2022.11.005>

Received 30 December 2021; Received in revised form 8 September 2022; Accepted 2 November 2022

Available online 12 November 2022

2665-9271/© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The tribe Meliponini (Hymenoptera: Apidae: Apinae) is that lineage encompassing the stingless bee entomological group (Michener, 2000) known to store and process honey and pollen in cerumen pots. The scientific literature refers to their honey as SBH (stingless bee honey). The term pot-honey was proposed to draw attention to the interactions of nectar to honey transformations within the cerumen container made up of stingless bee wax and plant resins (Vit et al., 2013). Given their peculiar chemical composition with higher water content and free acidity relative to honey extracted from *Apis mellifera* honeycomb, analytical comparisons were initiated with honeys produced by species of *Melipona* and *Scaptotrigona*, and extracted from cerumen pots in Brazil (Gonnet et al., 1964). After reviewing some 500 honeys from 66 species of stingless bees, Ávila et al. (2018) emphasized that meliponine honey positive health effects and market potential represented an innovative beacon in food, pharmaceutical, and cosmetic industries, in a world otherwise dominated by *A. mellifera* honey. Health promoting bioactive properties with bee-and-plant origin sustain the nutraceutical and medicinal uses of pot-honey (Pimentel et al., in press). According to the Codex Standard definition, honey is produced only by honey bees, and is described as “Honey consists essentially of different sugars predominantly glucose and fructose ...” (CODEX STAN, 1987), and water. A precious but unregulated product (Braghini et al., 2021). There are few regulatory quality factors for these pot-honeys, with the first national norm established for kelulut –all stingless bees in Malaysia (Department of Standards Malaysia, 2017) and the second from Argentina for ‘Yatef’ bees *Tetragonisca fiebrigi* (Secretaría de Regulación y Gestión Sanitaria y Secretaría de Alimentos y Bioeconomía, 2019).

Of the approximately 600 species of stingless bees, almost 400 species were found in the New World tropics (Camargo and Pedro, 2007). Substantial discoveries were predicted by Michener (2013), and the current diversity of meliponines is distributed in: 1. The Neotropics (ca. 500 species) from 34.90°S in Montevideo, Uruguay up to 27.03°N in Álamos, Sonora, Mexico, 2. Africa (ca. 16 species) from 28.54°S in Eshowe, South Africa up to 18.00°N in Njala, Sierra Leone, and 3. Indo-Malaysia/Australasia (ca. 100 species) from 36.41°S in Australia up to 24.23°N in Taiwan; exceptionally up to 4000 m.a.s.l. in Peru and Bolivia (Roubik and Vergara, 2021). The greatest global biodiversity of Meliponini is in Ecuador, 100 species in 64 km² of the Amazonian forest located in the Yasuní Biosphere (Roubik, 2018). Three of the key genera in the region are *Melipona* (73 spp.), *Scaptotrigona* (22 spp.) and *Geotrigona* (22 spp.) less represented in biodiversity collections, see number of species in Camargo and Pedro (2007) chapter and *Melipona* review (Engel, 2021). They are abundant bees, and represent critical components of the stingless bee fauna in the region of study. Sustainable meliponiculture increases the resilience of small-scale stingless bee keepers, obtains good quality products, and underpins productive crops due to pollination (Baumung et al., 2021). Stingless bee management and honey processing by meliponiculturists also imparts sanitary quality to this product (Heard, 2016).

The stingless bee keeping with *Melipona beecheii* by ancient Mayan peoples is well documented in Mexico (Weaver and Weaver, 1981). Despite the enormous biodiversity of 200 species of stingless bees in Ecuador (Vit et al., 2018), their ancient records of meliponiculture are unknown. However, the indigenous peoples are certainly aware of the bees and exploit these products, likely having done so for hundreds of years. The honey produced by stingless bees has been widely relished in the tropics (Schwarz, 1948), and interested the Brazilian Court of the International Exhibition (1862) to display *Melipona* and *Trigona* bees, along with samples of their, honey and wax (Smith, 1863). Nonetheless, but the request to showcase Meliponini in the Ecuadorian Pavillion of the Milan EXPO 2015 was declined. Honey pots have different volumes and shapes (Aguilar et al., 2013), *Melipona* generally has larger pots than those produced by species of *Scaptotrigona*, and *Geotrigona*, which are more elongated sausage-like pots –using palynological descriptors for

pollen morphology P/E (ratio polar/equatorial view of a pollen grain) a perprolate pot for the underground bee, subprolate for *Melipona* and spheroidal for *Scaptotrigona* (Vit P, personal observations), as illustrated in Fig. 1.

Bacteria, molds and yeasts found naturally in honey are non-pathogenic, and new taxa of microbiota with commensal or mutualistic ecological roles inside the stingless bee nests may offer sources of bioactive metabolites (Gilliam, 1997; Morais et al., 2013). Lactic acid is produced during the fermentation of honey (Vit et al., 2011). Little is known about relationships of honey and other products inside the stingless bee nest itself, e.g. *Tetragonisca angustula* from Venezuela (Pérez-Pérez et al., 2013), conversion of raw nectar into pot-honey (Leonhardt and Kaltenpoth, 2014). A seminal metabolomics approach investigated materials of the stingless bee nest: propolis, cerumen and honey produced by *Meliponula ferruginea* from Tanzania, storage modifications, and comparison with *A. mellifera* (Popova et al., 2021), and supported future scientific research to answer how pot-honey is produced and kept within the cerumen pots.

Ecuadorian *A. mellifera* honey marketed in Quito (Salvador et al., 2019) and pot-honeys from twelve species of stingless bees (Villacrés-Granda et al., 2021) were assessed for physico-chemical quality. Nuclear magnetic resonance (NMR) based metabolomics was approached to fingerprint the entomological origin (Vit et al., 2015; Razali et al., 2018) and to detect adulteration (Yong et al., 2022) of meliponine honey. The advantage of NMR spectroscopy is the chemical profiling of a number of organic functional groups, to simultaneously quantify mono-, di-, tri-saccharides, amino and organic acids, nucleobases, ethanol, HMF, and other characteristic constituents in a honey spectra (Popova et al., 2021). A systematic comparison between physico-chemical parameters and corresponding targeted NMR-based profile was done as a control for NMR in Chilean honeys (Fuentes Molina et al., 2020), and Iranian honeys (Khansaritoreh et al., 2021). The large number of organic compounds identified and quantified by targeted NMR expands descriptive categories as a complement to the honey quality control analyses. Walker et al. (2022) considered the challenge of multiple analytical techniques leading to complex reports on sophisticated honey adulteration.

The aim of this study was to apply ¹H NMR spectroscopy and a Honey Authenticity Test by Interphase Emulsion (HATIE) to compare the impact of the stingless bee genus in the organic matrix of their pot-honey. The phylogenetic origin (i.e., in terms of three monophyletic genera) on the honey each lineage produces. For this preliminary targeted metabolomics approach, twenty honey samples of *Geotrigona*, *Melipona* and *Scaptotrigona* were characterized with the content of 10 sugars, HMF and ethanol, 10 aliphatic organic acids, 10 amino acids, and 9 markers of botanical origin. Analysis of variance (ANOVA),




Fig. 1. Diagram of equatorial view of honey pots from Ecuadorian *Geotrigona*, *Melipona*, and *Scaptotrigona* (left to right).

hierarchical cluster analysis (HCA) and principal component analysis (PCA) based on NMR profiles were used to detect similarities and differences between the honey produced by *Geotrigona*, *Melipona*, and *Scaptotrigona* in Ecuador, and to position three test pot-honey samples at the genus-level in the database. An idea on entomological markers and searching phylogenetic traits in honey is given at the end. A generic approach to study the current role and interactions of stingless bees and their associated microbiota on sugary resources is to further in understanding of environmental factors and putative phylogenetic relatedness of honey.

2. Materials and methods

2.1. Pot-honey sampling

The harvest of pot-honeys was done in seven provinces of Ecuador: El Oro, Loja, Manabí, Napo, Orellana, Pastaza, and Santa Elena (Fig. 2). The three biomes were represented: Coast (El Oro, Manabí, Santa Elena), Andes mountains known as Sierra (Loja), and Amazonian forest (Napo, Orellana, Pastaza). Twenty samples of mature honey from stingless bee colonies were extracted by suction with a plastic syringe connected with a rubber tube, after cutting a circular section with a syringe needle, from sealed cerumen pots –this means the nectar or the honeydew has been transformed into honey, similarly to the operculated *Apis mellifera* honeycombs. Two samples of *Melipona* were collected by Kichwa. The honey was transported at environmental temperature and kept frozen in the dark until analysis in the laboratory. The genus of stingless bee specimens deposited in the collection Prof. J.M.F. Camargo (RPSP) was

identified by S.R.M. Pedro at Universidade de São Paulo, Ribeirão Preto, Brazil. A further discovery of a new species was done by Engel (2022) on specimens submitted to C.D. Michener and deposited in the Snow Entomological Museum Collection (SEMC) at the University of Kansas, Lawrence, Kansas, USA.

2.2. NMR

In nuclear magnetic resonance (NMR) spectroscopy, the chemical shift is the resonant frequency of a nucleus relative to a standard in a magnetic field. Often the position and number of chemical shifts are diagnostic of the structure of a molecule.

2.2.1. NMR reference sample and honey sample preparation, targeted quantitative $^1\text{H-NMR}$ spectra acquisition and processing

All chemicals used in NMR were of analytical grade (>99% purity). The HI 44518 - QuantRefA-NMR-Tube 5 mm, 600 μL (Reference Sample for quantification in food applications) was the NMR mixture analysis. All NMR spectra were acquired immediately after honey dilutions as previously described (Vit et al., 2022), on a Bruker Ascend TM 400 MHz FoodScreener (BrukerBiospin, Rheinstetten, Germany) equipped with a 5 mm PA BBI 400SI H-BB-D-05 Z probe and using Bruker SampleXpress (BrukerBiospin, Rheinstetten, Germany) for automatic sample change. The compounds were quantified by using the Honey-Profiling routine (release 1.0, BrukerBiospin, Rheinstetten, Germany) by automatic integration of the peak area calculated with an external standard (Spraul et al., 2009).



Fig. 2. Locations of pot-honey harvested in seven provinces of Ecuador: 1. El Oro, 2. Loja, 3. Manabí, 4. Napo, 5. Orellana, 6. Pastaza, and 7. Santa Elena.




Fig. 3. Number of honey phases after the HATIE *Geotrigona* (left), *Melipona* (center), and *Scaptotrigona vitorum* (right). After: Vit (2022a).




Fig. 4. Number of phases in the honey authenticity test From: Vit (2022a).

2.3. Honey authenticity test by interphase emulsion (HATIE)

This method was protected by two patents (Vit, 1995, 1999). It was created to detect fake honey (Vit, 1998). A disposable syringe was used to measure 0.5 mL honey and dilute it with 0.5 mL distilled water, 1 mL of diethyl ether was added and shaken by up and down wrist movements for 20 s, let stand still for 1 min, and the number of phases was observed, and noted. It was repeated with a 10 mL cylinder using 1 g honey, diluted with 1 g distilled water, diethyl ether was added to duplicate the volume of the dilution, and vortexed. It was let stand still for 1 min, and the number of phases was observed, and noted. Two patterns with one, or three phases were reported with this test for the genuine pot-honeys studied here, and illustrated in Fig. 3 (Vit, 2022a). The unique phase of *Scaptotrigona vitorum* Engel, 2022 honey was produced by an active biosurfactant of suspected microbial origin.

Vit (2022a) presented a diagram for the number of phases in six Ecuadorian honey types: 1. One phase (*Scaptotrigona vitorum*), 2. Two phases (fake honey), and 3. Three phases (*Apis*, *Geotrigona*, *Melipona*, *Tetragonisca*) (see Fig. 4).

2.4. Statistical analysis

The three entomological honey types were analyzed by ANOVA one way ($P < 0.05$), and post-hoc Scheffé test was performed to compare organic chemical components based on targeted NMR quantifications of pot-honey produced by *Geotrigona*, *Melipona* and *Scaptotrigona*, using the version 26 IBM SPSS Statistics software (IBM Corp, 2019). Hierarchical cluster analysis (HCA) for sugars and organic acids were generated using the squared Euclidean distance between groups and using the Ward's method. Pearson correlation matrix ($P < 0.05$), and Principal component analysis (PCA) were conducted using (XLSTAT Software Premium, 2021). Microsoft Power Point 2018 was used to prepare, integrate, and edit the figures composed from multiple sources.

3. Results and discussion


The following 41 parameters (10 sugars, HMF and ethanol, 9 organic acids, 10 amino acids, and 10 markers of botanical origin) were quantified in the $^1\text{H-NMR}$ spectra of Ecuadorian pot-honey samples: 1. Sugars (fructose, glucose, sucrose, gentiobiose, maltose, maltotriose, mannose, melezitose, raffinose, and turanose), 2. HMF and ethanol, 3. Aliphatic organic acids (acetic acid, citric acid, formic acid, fumaric acid, lactic acid, malic acid, pyruvic acid, quinic acid, and succinic acid), 4. Amino acids (alanine, aspartic acid, glutamine, isoleucine, leucine, phenylalanine, proline, pyroglutamic acid, tyrosine, valine), 5. Markers of botanical origin (acetoin, 2,3-butanediol, dihydroxyacetone, kynurenic acid, methylglyoxal, methylglyoxal dihydrate, methylglyoxal monohydrate, 3-phenyllactic acid, shikimic acid, and trigonelline). Quinic acid, isoleucine and kynurenic acid were not detected in the twenty honeys analyzed. See a Supplementary Table S1 with the molecular formula, chemical structure, regions of the $^1\text{H-NMR}$ spectra (ppm), and signal type for the 41 organic metabolites.

In the study of commercial Manuka honey the aliphatic components of honey (lactic acid, acetaldehyde, methylglyoxal dihydrate and methylglyoxal monohydrate, pyruvaldehyde, and pyruvic acid) showed signals in the first spectroscopic region (1.3–2.3 ppm), and phenyllactic acid in the aromatic region (6.0–8.0 ppm) of the $^1\text{H NMR}$ spectra (Gresley et al., 2012). In a comparison of commercial Ecuadorian *Apis mellifera*, *Geotrigona* and *Scaptotrigona* honeys, a central region (3.5–4.5 ppm) of the $^{13}\text{C NMR}$ spectra was characteristic for each bee genus, because the endogenous lipophilic markers of the bees obtained with chloroform extracts were independent of floral and geographical origin (Vit et al., 2015), as described.

The organic chemical profile (41 metabolites) for the honey produced by three genera of stingless bees had remarkable differences and similarities. The variability of the honey they produced, was a biodiversity trait. Comparative $^1\text{H-NMR}$ spectra at stingless bee genus level were presented in Figs. 5 and 6. The tabulated quantitative data were described and discussed accordingly in each section of metabolites (3.1 to 3.5). Targetted $^1\text{H NMR}$ was a powerful technique for multi-parametric analysis. Qualitative identifications of selected metabolites in the reference standard and their corresponding quantitative measurements were possible.

A considerably high number of honey metabolites was detected in each $^1\text{H-NMR}$ spectrum with a complex profile with only few isolated and identifiable signals from specific metabolites. The analysis of the spectrum with overlapped signals may hinder both peak identification and concentration determination. The $^1\text{H-NMR}$ spectra of three representative pot-honey of *Geotrigona*, *Melipona* and *Scaptotrigona* were shown in Fig. 5. Differences in the number and/or intensity of the peaks was correlated with the stingless bee genus. *Geotrigona* had very high intensities in the aliphatic region, showing the high concentrations of acetic acid and lactic acid. The signals of carbohydrates with anomeric and ring protons were comprised in a very narrow and overcrowded region of the $^1\text{H-NMR}$ spectrum (3.0–5.5 ppm). Signals of highly concentrated sugars like fructose and glucose were easily identified because of their high intensities. The aliphatic $^1\text{H-NMR}$ region of the spectra (3.0–0.1 ppm) contained AOs and amino acids. The aromatic region (10–6 ppm) comprised signals from other honey metabolites such as aromatic amino acids, and terpenoids used as botanical markers.

A more detailed view of the spectra was provided in Fig. 6. In the *Geotrigona* pot-honey, the sugar region of the spectrum showed a very characteristic profile with a high content of raffinose of 5.6 g/100g. Fructose and glucose signals were identifiable but the concentrations were under reference values of *Apis mellifera* honey. The low intensity (concentration) of glucose was remarkable in contrast to the very high concentrations of acetic acid and lactic acid. Signals of side chains of amino acids were easily identified in the aliphatic region of the spectrum, and in the aromatic region for the aromatic amino acids phenylalanine and tyrosine. In the *Melipona* pot-honey the intensity of raffinose



Metabolites: AcA acetic acid, EtOH ethanol, F fructose, G glucose, LacA lactic acid, M maltose, Mt maltotriose, R raffinose, S sucrose.

Fig. 5. Targeted $^1\text{H-NMR}$ spectra of *Geotrigona*, *Melipona*, and *Scaptotrigona* pot-honeys (400 MHz, 300K, PBS buffer, pH 3.1).

Metabolites: AcA acetic acid, EtOH ethanol, F fructose, G glucose, LacA lactic acid, M maltose, Mt maltotriose, R raffinose, S sucrose.


peaks was clearly lower than in *Geotrigona* and the quantification revealed a concentration of 0.3g/100g, which was still higher than the raffinose concentration of *Apis mellifera* honey. Unusual mannose peaks were easily identifiable. The aromatic region comprised peaks from aromatic amino acids, and some additional signals corresponded to other aromatic compounds. In the *Scaptotrigona* pot-honey the aliphatic region showed an intense signal of ethanol, acetic acid, and lactic acid, but lower than *Geotrigona*. These metabolites are typical of fermentation processes.

3.1. Sugars

The signals of glucose and fructose have a key role to discriminate *A. mellifera* honey (Lolli et al., 2008; Ohmenhaeuser et al., 2013). It is essential overcoming the honey matrix interference and the two main sugars with much higher peak intensities that may cover remaining

signals of other compounds. The targeted sugar profile consisted of monosaccharides (fructose, glucose, and mannose), disaccharides (gentiobiose, maltose, sucrose, and turanose), and trisaccharides (maltotriose, melezitose, and raffinose). The contents of the ten targeted sugars, the fructose + glucose value and the fructose/glucose ratio (Table 1) have differences and similarities between the three honey types. The total content of sugars (g/100 g honey) varied from 25.07 *Geotrigona* < 54.73 *Melipona* < 57.75 *Scaptotrigona*.

The ratio fructose/glucose (F/G) is used to predict slow crystallization for values < 1.3. This ratio was similar in the three genera (1.3–1.4), lower than the unprecedented above 2 reported for the Tanzanian *Meliponula ferruginea* (Popova et al., 2021), but similar to the Australian *Tetragonula carbonaria* (Persano Oddo et al., 2008). The F/G was 2.06 in one *Geotrigona* honey. However the low fructose content of *Geotrigona* (10.58 g/100 g) was almost triplicated in the other two genera, as well as the glucose content of *Geotrigona* (7.81 g/100 g) was triplicated in the



Metabolites: AcA acetic acid, Ala alanine, EtOH ethanol, F fructose, G glucose, Ile isoleucine, LacA lactic acid, Leu leucine, M maltose, Ma mannose, Mt maltotriose, Phe phenylalanine, R raffinose, S sucrose, T turanose, Tyr tyrosine Val valine. TSP trimethylsilylpropanoic acid (chemical shift reference).

Fig. 6. Targeted $^1\text{H-NMR}$ spectra of *Geotrigona*, *Melipona* and *Scaptotrigona* pot-honey (400 MHz, 300K, PBS buffer, pH 3.1), and expansions of aromatic, sugar and aliphatic regions (plots 1–4). The intensities in vertical scales of the NMR regions have been expanded to show signals of less concentrated metabolites. **Metabolites:** AcA acetic acid, Ala alanine, EtOH ethanol, F fructose, G glucose, Ile isoleucine, LacA lactic acid, Leu leucine, M maltose, Ma mannose, Mt maltotriose, Phe phenylalanine, R raffinose, S sucrose, T turanose, Tyr tyrosine Val valine. TSP trimethylsilylpropanoic acid (chemical shift reference).

Table 1
Sugar contents in Ecuadorian pot-honey quantified with ¹H NMR.

Sugars (g/ 100g honey)	Stingless bee genus		
	<i>Geotrigona</i> (n = 8)	<i>Melipona</i> (n = 4)	<i>Scaptotrigona</i> (n = 8)
Fructose	10.58 ± 0.79 ^a [7.83–13.65]	29.53 ± 1.84 ^b [25.36–33.93]	29.85 ± 0.49 ^b [27.92–31.81]
Glucose	7.81 ± 0.94 ^a [4.9–11.92]	23.23 ± 0.79 ^b [21.65–25.30]	22.33 ± 0.29 ^b [21.31–23.42]
Sucrose	0.15 ± 0.02 ^a [0.07–0.28]	0.10 ± 0.05 ^a [0.02–0.24]	0.20 ± 0.06 ^a [0.01–0.45]
Gentiobiose	0.20 ± 0.07 ^a [0.00–0.44]	0.02 ± 0.01 ^a [0.01–0.04]	0.17 ± 0.11 ^a [0.01–0.89]
Maltose	0.49 ± 0.05 ^a [0.27–0.69]	0.83 ± 0.32 ^a [0.27–1.47]	2.02 ± 0.21 ^b [0.92–2.75]
Maltotriose	0.10 ± 0.01 ^a [0.03–0.15]	0.05 ± 0.02 ^a [0.01–0.09]	0.10 ± 0.01 ^a [0.03–0.14]
Mannose	n.d.	0.04 ± 0.02 [0.00–0.08]	n.d.
Melezitose	0.07 ± 0.00 ^a [0.05–0.08]	0.12 ± 0.03 ^a [0.08–0.21]	0.25 ± 0.03 ^b [0.16–0.38]
Raffinose	5.03 ± 0.31 ^b [4.13–6.67]	0.45 ± 0.06 ^a [0.34–0.62]	1.24 ± 0.09 ^a [0.88–1.69]
Turanose	0.64 ± 0.05 ^a [0.37–0.78]	0.36 ± 0.05 ^a [0.27–0.50]	1.58 ± 0.28 ^b [0.00–2.54]
Total	25.07	54.73	57.75
quantified sugars			
Fructose + Glucose	18.39 ± 1.68 ^a [13.08–24.50]	52.87 ± 1.75 ^b [49.12–56.10]	52.17 ± 0.60 ^b [49.71–54.96]
Fructose/ Glucose	1.42 ± 0.10 ^a [1.05–2.06]	1.27 ± 0.11 ^a [1.07–1.57]	1.34 ± 0.03 ^a [1.24–1.48]

Results are expressed as means ± standard error and [minimum - maximum] values. n.d. not detected. Different superscripts in a row show significant difference between G-M-S honeys with ANOVA (Scheffé, $P < 0.05$).

Melipona and *Scaptotrigona* honeys, possibly indicating this underground bee has more active enzymes to catabolize fructose and glucose from the nectar, or its associated microbiota does it. Consequently the fructose + glucose was also lower in the *Geotrigona* (18.4) than *Melipona* (52.9) and *Scaptotrigona* (52.2). Sucrose was not statistically different but *Melipona* (0.10) < *Geotrigona* (0.15) < *Scaptotrigona* (0.20). Very low values compared to the maximum 5% permitted for *A. mellifera* honey (CODEX, 1987). Average gentiobiose varied from 0.02 in *Melipona* and 0.17 to 0.20 in the other two honey types.

Persano Oddo et al. (2008) observed a putative maltose (15.3–22.8 g/100 g) of the Australian *Tetragonula carbonaria* honey not perfectly coincident with the HPLC maltose standard, and suggested a further investigation, similar to the previous observations (2.5–3.2 g/100 g *Melipona* and 19.7–32.3 g/100 g other genera) by Bogdanov et al. (1996) in Venezuelan pot-honeys. Therefore, the maltose content of *Tetragonula carbonaria* reported in 2008, could correspond to the trehalulose recently discovered by Fletcher et al. (2020) in pot-honey harvested from four genera of stingless bees in Australia (*Tetragonula*), Brazil (*Tetragonisca*), and Malaysia (*Geniotrigona* and *Heterotrigona*). A new international regulation for honey including trehalulose is mandatory (Zawawi et al., 2022), and also the inclusion of trehalulose in the reference sample for targeted NMR Bruker's Honey-Profilng. However the trehalulose content of 28.38 g/100 g honey for the Australian and Malaysian species, and the maltose content that varied from 0.49 to 2.02 g/100g in the Ecuadorian honey would not match with that quantity. It will not match either with the lower quantities of maltose in *Melipona* and *Scaptotrigona* species from Venezuela, it could be similar to values of other stingless bees (*Friesoemellita nigra paupera*, *Nannotrigona* aff. *chapadana*, *Tetragonisca angustula*) but it was higher for two *Nannotrigona* aff. *varia* (38.7–56.4) g/100 g honey, with corresponding fructose (16.1–13.9) and glucose (13.9–6.7) (Vit et al., 1998a). Maltose

content of *Meliponula ferruginea* honey from Tanzania was low (1.56) and trehalulose lower than that (0.35), their fructose (26.9) high and lower glucose (12.8) causing the distinctive F/G > 2 (Popova et al., 2021). The fructose + glucose content was intermediate to Ecuadorian honey, approximately 20 < 30 < 40 < 50 g/100g comparing *Geotrigona* < Australia-Malaysian *Geniotrigona-Heterotrigona-Tetragonula* < African *Meliponula* < Neotropical *Melipona* and *Scaptotrigona*. The trehalulose was also suggested as an additional marker of honeydew honey in Spain, instead of quercitol (de la Fuente et al., 2007), and similarly in some Bulgarian and Macedonian honeydews (Gerginova et al., 2020). Some statistical differences are significant for a higher maltose in *Scaptotrigona* (2.02), but the ranges of individual honeys overlap between *Melipona* [0.27–1.47] and *Scaptotrigona* [0.92–2.75].

The remaining five sugars were quantified in minor concentrations. Mannose was detected only in *Melipona* (0.04) and not in the other two genera. A lower maltotriose in *Melipona* (0.05), higher melezitose in *Scaptotrigona* (0.25), higher raffinose in *Geotrigona* (5.03), and higher turanose in *Scaptotrigona* (1.58). If the post-hoc ANOVA is done with Duncan test, less conservative than the Scheffé we used, raffinose is the unique sugar to differentiate the honey produced in our collection of honeys produced by *Geotrigona*, *Melipona* and *Scaptotrigona*. The total quantified sugars (g/100 g) varied from 25.07 for *Geotrigona*, 54.73 for *Melipona* and 57.75 for *Scaptotrigona*, a hallmark for pot-honey compared to higher concentrations of sugars in *A. mellifera* honey – a suggested standard of reducing sugars above 50g/100g for *Melipona* and *Scaptotrigona* was valid for the fructose + glucose values in Table 1 (Vit et al., 2004). The melezitose could be an indicator of honeydew honey (Persano-Oddo and Piro, 2004), and the raffinose too (Consonni et al., 2012). Interestingly, positive Pearson correlations were found between contents of glucose and fructose (0.968), maltose and melezitose (0.853), maltose and turanose (0.801). Negative correlations were observed between raffinose and fructose (–0.941), and glucose (–0.970). Raffinose (consists on galactose, glucose, and fructose) can be hydrolyzed to sucrose and galactose by the enzyme α -galactosidase (α -GAL). This may explain its negative Pearson correlations with fructose and glucose. Raffinose family of oligosaccharides (RFOs) are α -1, 6-galactosyl extensions of sucrose. This group of plant oligosaccharides is known to serve as desiccation protectant in seeds, as a transport sugar in phloem sap, and as a storage sugar. Humans lack the α -GAL enzyme to break down this trisaccharide in their small intestine.

Nineteen sugars from aqueous extracts were identified in Italian honey of acacia, chestnut, rhododendron, multiflora and mountain by NMR: fructose, glucose, gentiobiose, isomaltose, kojibiose, maltose, maltulose, melibiose, nigerose, palatinose, sucrose, turanose, erlose, isomaltotriose, kestose, maltotriose, melezitose, raffinose, and maltotetraose (Consonni et al., 2012). Including also 11 unknown sugars, these authors suggested characteristic sets as identity cards for the Italian honeys: acacia (sucrose, turanose, fructose), chestnut (nigerose, kojibiose, raffinose, isomaltose, isomaltotriose, gentiobiose), rhododendron (melezitose, α -glucose, maltose, erlose, kestose), multiflora (β -glucose), and mountain (melezitose, α -glucose). Jang et al. (2016) studied 23 sugars (16 disaccharides and 8 trisaccharides) in Korean acacia, chestnut, multiflora, and artificial honey. Raffinose was more frequent and abundant in chestnut honey (0.01–0.17) mg/100 g.

All sugars separated the three honey types in distinctive clusters (Fig. 7A). The trisaccharide raffinose quantified in honey by NMR (see Table 1) was used to create a dendrogram, where a separation of *Geotrigona* from a cluster of *Scaptotrigona* and *Melipona* honeys was achieved (See Fig. 7B). *Melipona* and *Scaptotrigona* pot-honeys were separated by discriminant analysis of ten quality factors on Venezuelan species using only reducing sugars, sucrose, diastase activity and nitrogen, after factor reduction (Vit et al., 1998). Recently (Ávila et al., 2019) confirmed two distinctive clusters of Brazilian *Melipona* and *Scaptotrigona* honey by PCA of normalized mineral content and physicochemical analysis (moisture,




Fig. 7. Dendrograms based on the content of all sugars (A) and raffinose (B) in honey produced by *Geotrigona* (Geo), *Melipona* (Mel) and *Scaptotrigona* (Sca). Observe that total sugars (A) generated alternated Mel-Sca honeys in the Mel & Sca cluster, but the raffinose (B) has ordered Mel first, followed by Sca in the root of the dendrogram. HCA by Euclidean distance using Ward's method.

Aw, electrical conductivity, total acidity, °BRIX, pH, colour, ash and total minerals). On a botanical approach, two clusters of *Heterotrigona itama* honeydew and blossom honey were revealed by PCA of physico-chemical (ash content, hydrogen peroxide, free acidity, total mineral elements, K, Mg, Ca, total phenolics), and antioxidant (ferric reducing power) parameters in Malaysia (Ng et al., 2021). The honey produced by two species (*Tetragonula carbonaria* and *Tetragonula hockingsi*) of Australian stingless bees was separated from honey of two species (*Heterotrigona itama* and *Geniotrigona thoracica*) from Malaysia by HCA and PCA of moisture, electrical conductivity, pH, free acidity, color, glucose, fructose, trehalulose, total phenolics, total minerals, and 14 mineral elements (Zawawi et al., 2022).

3.2. Hydroxymethylfurfural and ethanol

The hydroxymethylfurfural (HMF) is a degradation product of fructose caused by heating and aging, but also increases in acidic environments like that of honey. The presence of ethanol in honey derives from fermentation.

Both HMF and ethanol contents of honey were given in Table 2. The HMF content was higher in the *Geotrigona* (40.8 mg/kg) honey than the other two genera (7.2 *Scaptotrigona* and 20.9 *Melipona*), but ethanol was lower in this honey with substantial acetic and lactic acid concentrations caused by fermentation. The content of ethanol was also higher than *A. mellifera* in *Meliponula ferruginea* honey from Tanzania (Popova et al., 2021).

Table 2
HMF and ethanol contents in Ecuadorian pot-honey quantified with ¹H NMR.

Ethanol and HMF (mg/kg honey)	Stingless bee genus		
	<i>Geotrigona</i> (n = 8)	<i>Melipona</i> (n = 4)	<i>Scaptotrigona</i> (n = 8)
Ethanol	440.24 ± 100.60 ^a [157.47–902.74]	1553.66 ± 928.10 ^a [17.56–3783.05]	1753.17 ± 299.86 ^a [998.5–3419.2]
HMF	40.76 ± 5.81 ^b [7.14–64.63]	20.90 ± 13.70 ^{ab} [0.00–60.21]	7.21 ± 2.70 ^a [1.64–19.71]

Results are expressed as means ± standard error and [minimum - maximum] values. Different superscripts in a row show significant difference between G-M-S honeys with ANOVA (Scheffé, P < 0.05).

3.3. Aliphatic organic acids (AOA)

Natural aliphatic organic acids in honey are intermediates or final products of plant and microbiota Krebs cycle (citric, succinic, glutaric, fumaric, and oxaloacetic), fermentation (lactic, acetic), bee enzymes (gluconic), indicators of aerobic or anaerobic processes.

The content of AOA in pot-honey is given in Table 3a. The total average content of nine organic acids varied from 0.29 to 4.43 g/100 g honey. Gluconic acid, lactic acid, and malic acid were studied in pot-honeys of *Tetragonula carbonaria* from Australia (Persano-Oddo et al., 2008), and *Melipona favosa* from Venezuela (Sancho et al., 2013), and from Malaysia (Shamsudin et al., 2019). Organic acids confer the organoleptic sour taste to honey and a distinctive acetic acid smell based on its concentration. Acetic acid and formic acid were volatiles identified in honey (Siegmund et al., 2018). All aliphatic organic acids were grouped in the targeted NMR approach of Chilean honeys (Fuentes Molina et al., 2020), here pyroglutamic acid was in the amino acids group, while kynurenic acid, 3-phenyllactic acid and shikimic acid were considered markers. Aliphatic organic acids in honey are of interest as indicators of fermentation or bioactivity. They were suggested as authenticity markers of bracing honeydew honey adulterated with floral honey from Brazil (Seraglio et al., 2021).

In our study, acetic acid 19.60 g/kg and lactic acid 24.30 g/kg from *Geotrigona* were statistically higher than *Melipona* (1.33 and 1.56) and *Scaptotrigona* (1.21 and 1.55) honeys respectively. Indeed, the most remarkable sensory feature of *Geotrigona* honey is the very strong sour taste and acetic acid smell. Acetic (3.05 g/100 g) and lactic (2.38 g/100 g) acids for the Tanzanian *Meliponula ferruginea* honey (Popova et al., 2021), similar lactic acid but acetic acid above *Geotrigona* in our study. The family Acetobacteriaceae are acetic acid bacteria (AAB) and break down carbohydrates under acidic media, they occurred in *Tetragonula* but not in *Austroplebeia australis* (Leonhardt and Kaltenpoth, 2014). Therefore, from our data, would be of interest to identify AAB in the Ecuadorian stingless bee honey.

Other organic acids were present at much lower concentrations in honey (mg/kg). Citric acid range was (20.6–284.2) and malic acid (0.0–210.9). The *Geotrigona* citric acid was almost triple of that in *Melipona* and double than *Scaptotrigona*. For malic acid *Geotrigona* (102.6) triplicated *Melipona* (35.7) in contrast with the low concentration found in *Scaptotrigona* (8.0). Three organic acids were evaluated in pot-honey of *Tetragonula carbonaria* from Australia by enzymatic

methods: citric acid (0.23) and malic acid (0.12) were similar to *A. mellifera* honeys, but D-gluconic acid (9.9) was higher in *T. carbonaria* honey (Persano Oddo et al., 2008). In the honey of *Melipona favosa* from Venezuela citric acid (0.05) and lactic acid (0.03) (Sancho et al., 2013) were lower than the Ecuadorian *Melipona*. Gluconic acid is known to be the main acid responsible for the free acidity of *A. mellifera* honey and may indicate a high activity of glucose oxidase at ripening. Formic acid was also more concentrated in the *Geotrigona* (54.6) than *Melipona* (11.6) and *Scaptotrigona* (12.9), lower than values reported for *Apis mellifera* unifloral honeys 199–506 *Castanea*, 50–128 *Eucalyptus*, and 186–209 mg/kg multifloral (Suarez-Luque et al., 2006). *Geotrigona* and *Scaptotrigona* generate alarm signals near to their nests, but formic acid is used by *Oxytrigona* in caustic defensive secretion (Roubik, 1989). Formic acid, the simplest of fatty acids is the alarm pheromone in ant species (Blum and Brand, 1972). Although fumaric acid was higher in *Scaptotrigona* (3.4) than *Geotrigona* (1.2) and *Melipona* (1.0), the variations caused no statistical difference between them. Pyruvic acid in *Geotrigona* (23.2) doubled the content of *Melipona* (16.1) and *Scaptotrigona* (10.0). Succinic acid was also higher in *Geotrigona* (61.0), some X5 of *Melipona* (12.0) and double of *Scaptotrigona* (25.2). Higher values were found in the honeydew honey from Brazil (0.5–0.7 g/100g (Brugnerotto et al., 2019). The quinic acid was absent in honey of the three genera, and the D-gluconic acid was not included in the NMR-based profiling, despite its importance in honey formation. It should be noted that gluconic acid has 16 stereoisomers, and has not been quantified in NMR-based profiles of honey so far because it is not easily measurable. D-gluconic acid can have theoretically 5 forms - one open, 2 pyranose, 2 furanose and all of them depending on pH could exist as acid or anion, all together 10 different forms in water (Simova S, personal communication). Gluconic acid accounted for 64.6–99.8% of total organic acids quantified by LC-MS/MS in *Apis mellifera* honey, and considered to be produced by worker bee enzymes such as glucose oxidase and the TCA cycle (Suto et al., 2020).

Lactic acid bacteria (LAB) microbiota are in symbiosis with *A. mellifera* (Olofsson et al., 2016) but Apini preserve their honey more dehydrated than Meliponini –fermentation by symbiont microbes such as *Zygosaccharomyces* and *Starmarella* species frequently isolated from their nests, have a role in pot-honey conservation (de Paula et al., 2021). Besides the LAB also acetic acid bacteria (AAB) are in symbiosis with the Australian *Tetragonula carbonaria* but not *Austroplebeia australis* (Leonhardt and Kaltenpoth, 2014). *Acetobacter*-like operational taxonomic unit (OTU) are associated with some stingless bees gut microbiota (Kwong et al., 2017). A cerumen pot made up of resins and stingless bee wax is rather soft and elastic, as needed to absorb the impact of anaerobic gas-related volume changes during nectar/honeydew fermentative processes to produce pot-honey (Vit P, personal observation), and additionally fibers of filamentous fungi evidenced in *Scaptotrigona depilis* brood cells (Menezes et al., 2015) may contribute to the material design with an organic holding mesh featuring its properties, and contributing to e.g. strength, elasticity, waterproof, bio-structure. A sealed honey pot is therefore, an ideal anaerobic container to maintain a wet environment in the warm nest temperature, needed for microbial proliferation in the sugary substrate harvested, carried and stored by the bees to fill open pots. *Bacillus cereus* was the most frequently isolated species in a *Melipona fasciata* nest from Panama (Gilliam et al., 1990). *Bacillus* species were the major bacteria found in a *Tetragonisca angustula* nest from Venezuela (Pérez-Pérez et al., 2013), and a *Heterotrigona itama* nest from Malaysia (Ngalimat et al., 2019). These spore forming bacteria secrete esterases, lipases, proteases, aminopeptidases, phosphatases, and glycosidases that transform food substrates into more digestible products. *Bacillus* species also secrete antibiotics and fatty acids to inhibit competing microorganisms which could spoil pot-honey or pot-pollen (Gilliam et al., 1990). The research of bee-microbes to protect pollinators may capitalize the discovery of novel bioactive compounds (Gilliam, 1997). Microbiome of stingless bees is highly variable. Wider bacteriological profiles of indoor surface were caused by a diversity of

Table 3aAliphatic organic acid contents in Ecuadorian pot-honey quantified with ¹H NMR.

Aliphatic organic acids (mg/kg honey)	Stingless bee genus		
	<i>Geotrigona</i> (n = 8)	<i>Melipona</i> (n = 4)	<i>Scaptotrigona</i> (n = 8)
Acetic acid (g/kg)	19.60 ± 1.45 ^b	1.33 ± 0.33 ^a	1.21 ± 0.34 ^a
Monocarboxylic acid	[15.23–25.56]	[0.46–2.06]	[0.21–2.85]
Citric acid	145.71 ± 27.20 ^b	27.07 ± 3.01 ^a	97.71 ± 25.41 ^b
Tricarboxylic acid	[77.16–284.20]	[20.20–32.71]	[30.88–224.74]
Formic acid	54.61 ± 15.57 ^b	11.62 ± 4.26 ^a	12.86 ± 2.45 ^a
Monocarboxylic acid	[8.83–131.38]	[3.72–23.02]	[4.67–25.09]
Fumaric acid	1.17 ± 0.47 ^a	0.98 ± 0.45 ^a	3.39 ± 1.38 ^a
Dicarboxylic acid	[0.00–2.74]	[0.0–2.13]	[0.0–11.68]
Lactic acid (g/kg)	24.30 ± 1.65 ^b	1.56 ± 0.45 ^a	1.55 ± 0.39 ^a
Monocarboxylic acid	[19.28–30.45]	[0.41–2.63]	[0.43–3.39]
Malic acid	102.59 ± 25.92 ^b	35.68 ± 20.60 ^{ab}	7.95 ± 7.95 ^a
Dicarboxylic acid	[0.00–210.91]	[0.0–72.78]	[0.00–63.61]
Pyruvic acid	23.18 ± 2.55 ^a	16.13 ± 9.79 ^a	9.98 ± 0.64 ^a
Monocarboxylic acid	[14.79–34.48]	[3.86–45.27]	[7.55–13.02]
Quinic acid	n.d.	n.d.	n.d.
Monocarboxylic acid			
Succinic acid	61.03 ± 6.20 ^b	12.88 ± 1.16 ^a	25.21 ± 5.74 ^a
Dicarboxylic acid	[37.08–92.1]	[10.73–15.79]	[11.64–56.18]
Total aliphatic organic acids	44294.73	2972.83	2950.11

Results are expressed as means ± standard error and [minimum - maximum] values, n.d. not detected. Different superscripts in a row show significant difference between G-M-S honeys with ANOVA (Scheffé, P < 0.05).

materials –of animal, plant, or soil origin– used by diverse species of stingless bees to build their nests (de Sousa, 2021).

A tentative set of aliphatic or non-aromatic organic acids was proposed for the *Geotrigona* exceptionally high free acidity and sour taste, in descending order of average concentration: 1. Lactic acid (24.3 g/kg), 2. Acetic acid (19.6 g/kg), 3. Citric acid (145.7 mg/kg), 4. Malic acid (102.6 mg/kg), 5. Succinic acid (61.0 mg/kg), and 6. Formic acid (54.6 mg/kg).

The organic acids quantified in honey (see Table 3a) were used for HCA to create a dendrogram, where the *Geotrigona* cluster was separated from a cluster of *Scaptotrigona* and *Melipona* honey in Fig. 8. Similar clusters were obtained in the dendrogram with all the sugars and the sugar raffinose, Fig. 7.

In Table 3b harmonized units of g organic acid/100 g honey were used to express the data from Table 3a (g/kg and mg/kg) compared with the literature: 1. *Tetragonula carbonaria* (Tcar) from Australia (Persano Oddo et al., 2008) Unifloral and multifloral *Apis mellifera* (AmS) Spanish floral honey by capillary zone electrophoresis CZE (Mato et al., 2006), 2. *Apis mellifera* (AmB) Brazilian bracinga honeydew honey by capillary electrophoresis CE (Seraglio et al., 2021) published as mg/100 g, 3. *Meliponula ferruginea* (Mul) honey from Tanzania, already using g/100 g (Popova et al., 2021). There are advantages and disadvantages in choosing any type of unit to express the content of organic acids in honey, because their range is wide. Using mg/kg is not practical for the hypervalues of most concentrated acids in pot-honey such as acetic acid and lactic acid. The visibility of g/100 g is reduced or even lost for the less concentrated acids, but these are usual units in food composition. Possibly mg/100 g honey would be an intermediate option.

The total aliphatic organic acids varied from 0.29 g/100 g honey for *Melipona* and *Scaptotrigona* in Ecuador to 5.76 for *Meliponula* in Tanzania (Popova et al., 2021), and 1.04–1.53 g/100g *Apis mellifera* honey in Table 3b. The *Geotrigona* had the highest content of 4.44 g AOA/100 g honey from Ecuador, which was also reflected in the free acidity of this remarkably sour honey with more than 500 mEq/kg honey (Vit, 2022b). In the twelve species of stingless bees studied by (Villacrés-Granda et al.




Fig. 8. Dendrogram based on the content of all aliphatic organic acids in honey produced by *Geotrigona* (Geo), *Melipona* (Mel) and *Scaptotrigona* (Sca). HCA by Euclidean distance using Ward’s method.

Table 3b

Contents of aliphatic organic acids (g/100 g) in *Apis mellifera* (AmS) honey from Spain (Mato et al., 2006), (AmB) from Brazil (Seraglio et al., 2021), and pot-honeys produced by *Tetragonula carbonaria* (Tcar) from Australia (Persano Oddo et al., 2008), *Meliponula ferruginea* (Mul) from Tanzania (Popova et al., 2021), *Geotrigona* (Geo), *Melipona* (Mel) and *Scaptotrigona* (Sca) from Ecuador in this work.

Bee	AcA	CitA	FoA	FuA	GluA	LacA	MalA	PyrA	ShA	SucA	Total
AmS	0.02	–	0.03	–	0.95	0.02	0.02	n.d.	–	0.00	1.04
AmB	0.02	0.05	0.02	–	0.96	0.17	0.14	–	–	0.13	1.53
Geo	1.96	0.02	0.01	0.01	–	2.43	0.01	0.00	0.00	0.00	4.44
Mel	0.13	0.00	0.00	0.00	–	0.16	0.00	0.00	0.00	0.00	0.29
Mul	3.05	0.18	0.01	–	–	2.38	0.12	0.00	0.02	0.00	5.76
Sca	0.12	0.01	0.00	0.00	–	0.16	0.00	0.00	0.00	0.00	0.29
Tcar	–	0.01	–	–	0.99	–	0.02	–	–	–	1.02

AcA acetic acid, CitA citric acid, FoA formic acid, FuA fumaric acid, GluA gluconic acid, LacA lactic acid, MalA malic acid, PyrA pyroglutamic acid, ShA shikimic acid, SucA succinic acid.

2021), the total organic acids measured by HPLC varied from 0.44 mg/100 g honey *Melipona mimetica* and 3.00 mg/100 g for *Tetragonisca angustula* from Ecuador. These authors measured acetic, citric, lactic and oxalic acids. Those values could be possibly in g/100 g honey. Their data is a great contribution with 27 pot-honeys, it was not only limited to all the honey quality standards, but also investigated the content of vitamin C, amino acids leucine and proline, organic acids, contaminant and nutritional metals, besides antibacterial activity. Acetic acid was the most abundant in *Scaptotrigona polysticta* and *Tetragonisca angustula* honey, but oxalic acid –a possible veterinary residue for *Varroa destructor* treatment of *Apis mellifera*, was the major organic acid in 10/12 stingless bee species. Its origin is unknown in pot-honey and may have a sensory impact on this already sour honey. In a study with eight Austrian unifloral honeys, a very unlikely set for floral volatiles (thujone isomers, eucalyptol, camphor, eugenol, thymol, and carvacrol) was suspected to originate in the use of essential oils by bee keepers (Siegmund et al., 2018). However, oxalic acid remained unchanged after varroa treatments (11–119 mg/kg) in Swiss honeys (Bogdanov et al., 2002), and similar values (14–114 mg/kg) in honeys from Spain (Mato et al., 2006).

From the eight organic acids studied by HPLC in *Geniotrigona thoracica* (Gt) and *Heterotrigona itama* (Hi) honey from Malaysia, formic acid was not detected (Shamsudin et al., 2019). These authors analyzed three monofloral types (acacia, starfruit and gelam) to observe further variations of AOA contents (g/kg) caused by the botanical origin respectively: Acetic acid (0.09, 0.08, 0.06) Gt, (0.30, 0.06, 0.01) Hi; citric acid (0.04, 0.05, 0.03) Gt, (0.04, 0.06, 0.09) Hi; gluconic acid (0.48, n.d., 0.55) Gt, (0.90, 0.39, 1.48) Hi; lactic acid (0.20, 0.03, 0.17) Gt, (0.15, 0.18, 0.18) Hi; malic acid (0.30, n.d., 0.30) Hi only; succinic acid (0.52, 0.07, n.d.) Gt (0.32, 0.38, 0.34) Hi, and tartaric acid (0.04, n.d., n.d.) Gt, (0.06, 0.03, n.d.) Hi. The succinic acid was not present in Gelam honey, the other AOA varied according to the bee species fed by different nectars. They found that gluconic acid was lower in starfruit than in the acacia and gelam honeys. Malaysian *H. itama* averages of floral and honeydew honey were similar and low for gluconic acid (0.47) and acetic acid (0.10) g/kg measured with enzymatic methods (Ng et al., 2021). The concentrations of these aliphatic organic acids given in (g/kg) should be multiplied by 10 to scale them into (g/100 g) as in Table 3b. All the Asian values will be very low compared with the African, Australian and South American pot-honeys, even for the gluconic

acid that was measured enzymatically in Australian *T. carbonaria*, by CE in *A. mellifera* from Brazil and Spain, but not by NMR.

3.4. Amino acids

The origin of amino acids in honey is mostly undescribed in literature, except proline. The content of amino acids in pot-honey was given in Table 4. Proline and phenylalanine were the major amino acids in the three genera studied here, followed by alanine and pyroglutamic acid. In *Meliponula ferruginea* honey from Tanzania the major amino acid after proline was pyroglutamic acid, rarely found in *Apis mellifera* honey (Popova et al., 2021). In the Ecuadorian honeys proline varied from 34.41 to 565.77 mg/kg and averages were statistically different between *Geotrigona* (416.58) and *Scaptotrigona* (286.91), and *Melipona* (81.37) the lowest. Isoleucine was absent in all the honeys. The total average content of amino acids varied from 277.82 to 1183.10 mg/kg honey.

Bumble bees do not differentiate between different amino acids, but between different concentrations of the same amino acid (Leonhardt et al., 2020). Indeed, amino acids were not a botanical marker candidate in unifloral and multifloral honeys (Hermosin et al., 2003; Carratu et al., 2011), they were not either entomological markers for the three meliponine genera studied here, but L-alanine was suggested as an entomological marker for *Tetrigona apicalis* from Malaysia (Razali et al., 2018). Besides the bee, the nectar and the pollen sources, the amino acid spectra of pot-honey can vary by the action of symbiotic microbiota in the stingless bee nest (Barbosa et al., 2017). The *Monoascus* sp. fungal mycelia growing on the food supply within the brood cells was essential for the larval development of *Scaptotrigona postica* (Menezes et al., 2015).

Residual proline of honey originates from salivary secretions in cephalic glands of stingless bee workers added to transformed nectar into honey. Proline content in honey of *Tetragonisca angustula* from Colombia was 770 mg/kg (Torres et al., 2004). Honeys of *Melipona scutellaris*, *Melipona quadrifasciata*, *Melipona subnitida*, and *Plebeia* sp. from Brazil varied between 201.6 and 924.6 mg/kg (Duarte et al., 2012). Similarly, honey of *Melipona beecheii* from Mexico varied from 264.5 to 1193.7 mg/kg (Moo-Huchin et al., 2015). Lower values were found in honey of *Melipona subnitida* and *Melipona scutellaris* from Brazil, ranging from

Table 4
Amino acid contents in Ecuadorian pot-honey quantified with ¹H NMR.

Amino acids (mg/kg honey)	Stingless bee genus		
	<i>Geotrigona</i> (n = 8)	<i>Melipona</i> (n = 4)	<i>Scaptotrigona</i> (n = 8)
Alanine	101.78 ± 20.84 ^b [44.11–237.31]	7.64 ± 1.27 ^a [5.35–10.16]	32.19 ± 5.61 ^a [6.1–59.6]
Aspartic acid	n.d.	n.d.	19.15 ± 19.15 [0.0–153.2]
Glutamine	43.59 ± 11.22 ^a [15.41–118.34]	8.40 ± 0.99 ^a [6.02–10.78]	35.36 ± 14.36 ^a [10.45–119.97]
Isoleucine	n.d.	n.d.	n.d.
Leucine	22.57 ± 12.31 ^a [0.0–93.73]	n.d.	27.66 ± 8.79 ^a [0.0–65.01]
Phenylalanine	415.86 ± 82.37 ^a [163.31–783.38]	125.37 ± 56.90 ^a [26.04–244.20]	312.53 ± 79.80 ^a [147.85–826.16]
Proline	416.58 ± 43.83 ^b [225.24–565.77]	81.37 ± 28.77 ^a [34.41–165.24]	286.91 ± 39.05 ^b [179.71–529.16]
Pyroglutamic acid	96.03 ± 31.61 ^a [0.00–218.26]	n.d.	73.90 ± 49.05 ^a [0.00–338.20]
Tyrosine	56.11 ± 11.99 ^a [25.62–124.62]	n.d.	99.81 ± 24.28 ^a [24.69–203.04]
Valine	30.58 ± 5.20 ^b [15.95–61.07]	0.31 ± 0.31 ^a [0.0–1.22]	15.67 ± 3.91 ^{ab} [3.72–37.17]
Total average amino acids	1183.10	277.82	903.18

Results are expressed as means ± standard error and [minimum - maximum] values, n.d. not detected. Different superscripts in a row show significant difference between G-M-S honeys with ANOVA (Scheffé, P < 0.05).

46.0 to 205.0 mg/kg (de Sousa et al., 2016). Biluca et al. (2019) analyzed 16 free amino acids (aspartic acid, glutamic acid, asparagine, glutamine, serine, arginine, glycine, threonine, alanine, proline, tyrosine, valine, leucine, isoleucine, phenylalanine, tryptophan) in honey of nine stingless bee species from Brazil, the two most concentrated and ubiquitous were phenylalanine (5.2–1231.0 mg/kg) and proline (12.1–762.0 mg/kg). Glutamine (1–605) was absent in half of the honeys, histidine was absent in all. Valine (16.1–85.9) was present in all honeys except *Melipona bicolor*, like tyrosine (8.5–156.0) which was not present also in one of the two *Melipona quadrifasciata* honeys, possibly indicating a non entomological origin. The quantification of 17 amino acids (arginine, histidine, lysine, phenylalanine, isoleucine, leucine, methionine, valine, threonine, tyrosine, proline, glutamic acid, aspartic acid, serine, alanine, glycine, and cysteine) was done for the honey produced by *Tetragonula laeviceps* in three locations of Indonesia (Agussalim et al., 2021). The average proline varied from 80.54 to 597 mg/kg, similar to the Ecuadorian pot-honeys ranges in Table 4 (34.4–565.8). Their higher concentrations were found for glutamic acid with 1238.74 (Sleman), 1922.98 (Lombok), and surprisingly it was ten fold lower in the third region (Gunungkidul), possibly related to the protein sources available. In the pot-honeys produced by twelve species of Ecuadorian stingless bees, proline concentration was higher than leucine, the other amino acid measured by (Villacrés-Granda et al., 2021). In their five *Melipona* spp. the average proline varied from 198.58 to 5132.71 mg/kg honey, in the *Scaptotrigona polysticta* it was 4179 mg/kg, and for the other species 84.56–1459.51 mg/kg. In this study comparing three genera of Ecuadorian stingless bees, the highest proline average was 416.58 mg/kg for *Geotrigona*.

Pyroglutamic acid is a less known and studied ubiquitous natural amino acid derivative in which the free amino group of glutamic acid or glutamine cyclizes to form a lactam. It was absent in *Melipona* honey and varied from 0.0 to 338.2 mg/kg with averages of *Geotrigona* (96.03) > *Scaptotrigona* (73.9).

3.5. Markers of plant origin

Volatile markers of monofloral honey from different regions may vary according to the flora. Putative markers were the volatile compounds most often reported in twenty selected monofloral honeys produced by *A. mellifera* (Machado et al., 2020).

Table 5
Plant markers in Ecuadorian pot-honey quantified with ¹H NMR.

Markers of botanical or entomological origin (mg/kg honey)	Stingless bee genus		
	<i>Geotrigona</i> (n = 8)	<i>Melipona</i> (n = 4)	<i>Scaptotrigona</i> (n = 8)
Acetoin	n.d.	17.10 ± 16.11 ^a [0.00–65.38]	2.06 ± 2.06 ^a [0.0–16.45]
2,3-Butanediol	502.31 ± 112.13 ^a [101.32–972.82]	264.29 ± 141.78 ^a [15.47–530.16]	127.59 ± 37.82 ^a [28.79–323.83]
Dihydroxyacetone	133.68 ± 13.41 ^b [90.89–183.83]	32.06 ± 7.39 ^a [23.01–54.15]	43.37 ± 7.63 ^a [14.76–85.64]
Kynurenic acid	n.d.	n.d.	n.d.
Methylglyoxal	n.d.	3.27 ± 3.27 ^a [0.0–13.18]	4.38 ± 3.01 ^a [0.0–22.37]
Methylglyoxal dihydrate	18.45 ± 9.14 ^a [0.0–58.15]	19.66 ± 5.38 ^a [7.85–43.51]	9.47 ± 2.19 ^a [0.00–20.64]
Methylglyoxal monohydrate	21.96 ± 2.94 ^b [14.14–39.29]	6.54 ± 1.12 ^a [3.56–8.59]	5.74 ± 1.42 ^a [1.92–12.76]
3-Phenyllactic acid	69.24 ± 34.80 [0.00–232.34]	n.d.	n.d.
Shikimic acid	11.52 ± 3.43 ^a [3.22–32.51]	10.47 ± 3.85 ^a [3.36–20.35]	30.46 ± 14.45 ^a [6.09–84.30]
Trigonelline	18.94 ± 3.61 ^a [5.64–32.47]	2.16 ± 1.55 ^a [0.00–6.59]	25.32 ± 8.84 ^a [6.09–84.30]

Results are expressed as means ± standard error and [minimum - maximum] values, n.d. not detected. Different superscripts in a row show significant difference between G-M-S honeys with ANOVA (Scheffé, P < 0.05).

In **Table 5**, the ten targeted botanical markers were quantified by ^1H NMR (mg/kg), with the most concentrated 2,3-butanediol and the absent kynurenic acid. The shikimic acid was included in this table.

3.5.1. Acetoin

Acetoin is 3-hydroxy-2-butanone, a natural volatile of the Australian *Eucalyptus leucoxylon* and *Eucalyptus melliodora* (D'Arcy et al., 1997). It is also present in heather honey from France (Radovic et al., 2000), Germany and Norway (Guyot et al., 1999). It was absent in *Geotrigona*. Only low concentrations were detected in *Melipona* (17.1) mg/kg and *Scaptotrigona* (2.1).

3.5.2. 2,3-Butanediol

The most concentrated botanical marker in the three Ecuadorian honeys was 2,3-butanediol, from 127.6 to 502.3 mg/kg honey being *Geotrigona* > *Melipona* > *Scaptotrigona*. This volatile was found in rosemary and thymus honeys from Spain (Pérez et al., 2002), and was reported (mg/kg) in a sensory study of the Austrian unifloral honey volatiloma for dandelion 0.02, robinia 0.50, rape 0.40, fir tree 0.05, chestnut 0.17, linden 0.11, orange 7.77, and lavender 0.03 (Siegmund et al., 2018), all less concentrated than the Ecuadorian honey.

3.5.3. Dihydroxyacetone

Dihydroxyacetone (DHA) is the precursor to methylglyoxal (Smallfield et al., 2018). Manuka phytochemicals including DHA were related with elements of the soil (Meister et al., 2021) Nectar chemical traits were different between sites, but plant-to-plant variation within sites were the largest (Icon et al., 2021). DHA varied from 32.1 to 133.7 mg/kg in the Ecuadorian pot-honeys, another botanical source is producing it because manuka is not growing in Ecuador. Glycerol is transformed to dihydroxyacetone by the bacterium *Acetobacter suboxidans* (Charney, 1978).

3.5.4. Kynurenic acid

Resonances of minor compounds also play a certain role such as quinoline alkaloids and kynurenic acid for chestnut honey (Truchado et al., 2009). Kynurenic acid was found in small concentrations in Alpine honey from Italy (Leoni et al., 2021). In Moroccan jujube *Ziziphus lotus* honey kynurenic acid contents were 17.7 mg/kg by HPLC-DAD-ESI-MS and 6.6 mg/kg by GC-EI-MS (Khallouki et al., 2021). In the Ecuadorian pot-honeys, kynurenic acid was non-detected, like in the Chilean honeys (Fuentes Molina et al., 2020).

3.5.5. Methylglyoxal, methylglyoxal dihydrate and methylglyoxal monohydrate

These are markers of manuka (*Leptospermum scoparium*) honey. Antibacterial activity of honey after hydrogen peroxide release is known as non-peroxide activity (NPA) in Australia, similar to the Unique Manuka Factor (UMF) in New Zealand, with a rating >10+ considered therapeutically active (Cokcetin et al., 2016). These authors explain DHA is converted to MGO during storage, indicating the potential of honey to increase bioactivity. Highest MGO >800 mg/kg corresponded to highest NPA (21.9–26.3%). Methylglyoxal (MGO) was not detected in *Geotrigona* honey but in *Melipona* and *Scaptotrigona* it was 3.3–4.4 mg/kg respectively. However, both MGO dihydrate (9.5–19.7) and MGO monohydrate (5.7–22.0) had no significant differences between the three honey types for MGO dihydrate, whereas MGO monohydrate was significantly highest in the *Geotrigona* honey.

3.5.6. 3-Phenyllactic acid

In thyme honey from Greece it varied from 1.5 to 28.1 g/kg (Alisandrakis et al., 2009). *Tetragonula carbonaria* raw honeys were harvested from "sugarbag" in Australia. The phenolic extracts averaged to 58.7 mg/kg and included 3-phenyllactic acid in a range from 3.7 to 12.7 mg of gallic acid equivalents/100 g honey (Massaro et al., 2014). Manuka (*Leptospermum scoparium*) and kanuka (*Kunzea ericoides*) nectars

and honeys from New Zealand contained 3-phenyllactic acid, 740 mg/kg manuka honey and 660 mg/kg kanuka honey (Bong et al., 2018), which is a broad spectrum antimicrobial compound. In **Table 5**, 3-phenyllactic acid was found only in *Geotrigona* honey, 69.2 mg/kg.

3.5.7. Shikimic acid

In microorganisms, shikimic acid is derived from phenolics, and the shikimate biosynthesis pathway produces precursors for the aromatic amino acids phenylalanine, tyrosine, and tryptophan. The physiology of the bee or the associated microbiota in the nest, also the bee diet, or a combination of them could explain the presence of shikimic acid in honey (Santos-Sánchez et al., 2019). A rare apple honey was characterized by high contents of shikimic acid-pathway derivatives (Kuš et al., 2013). In our study there is a wide variation from 3.22 to 84.30 mg/kg, as observed in **Table 5**, with a higher average in *Scaptotrigona* (30.5) than *Geotrigona* (11.5) and *Melipona* (10.5). Therefore it becomes interesting to describe the concentration of shikimic acid by *Scaptotrigona* in the honey they process, compared to that of *Geotrigona* and *Melipona*. However, variations were too high to separate these three honey types.

3.5.8. Trigonelline

This nucleobase has been identified as coffee origin. In a study of unifloral honey from Colombia and multifloral from Honduras, trigonelline ranged from 23 to 86 mg/kg honey, and was suggested as a marker of *Apis mellifera* unifloral honey of *Coffea arabica* (Schievano et al., 2015). It was present in the *Meliponula ferruginea* honey from Tanzania (Popova et al., 2021). Although the honey produced by the three stingless bee genera were not statistically different, averages here ranged from 2.2 to 25.3 mg/kg *Melipona* < *Geotrigona* < *Scaptotrigona*.

In the Chilean honey study (Fuentes Molina et al., 2020), a reduced set of markers excluding the aliphatic organic acids was named low molecular weight molecules, and they found it a quite heterogeneous group, like here. These markers did not separate the three genera of Ecuadorian honey.

In **Fig. 9**, a graphic comparison was presented for selected targeted NMR based concentrations (from **Tables 1, 3–5**) of sugars (fructose, glucose, raffinose), organic acids (acetic acid, lactic acid, citric acid X10), amino acids (proline, phenylalanine, tyrosine) and plant markers (2,3-butanediol, dihydroxyacetone, trigonelline) in Ecuadorian honey produced by *Geotrigona*, *Melipona* and *Scaptotrigona*. Statistical differences by ANOVA Scheffé post-hoc were detected in parameters from the four chemical groups, sugars, organic acids, amino acids and markers. Therefore it is worth to continue with them, other sugars and alcohols would be needed in the reference standard if possible for Bruker.

3.6. PCA multivariate analysis

In **Fig. 10**, the loading plots of active variables or honey components used in the PCA are organized for: 1. Sugars (10A), 2. Aliphatic organic acids (10B), 3. Amino acids (10C), and 4. Markers of plant or bee origin (10D). In these plots F1 x F2, the variability of honey was explained 67.87% by sugars, 67.52% by organic acids, 59.78% by amino acids, and 48.84% by plant-bee markers.

One set of sugars is organized in the upper left portion (fructose, glucose, mannose), and two in the lower portion, left (gentiobiose, maltotriose, raffinose) and right (maltose, melezitose, sucrose, turanose) quadrant (See **Fig. 10A**). The first dimension F1 explains 46.02% variations. F1 explains 21.85% of the variation useful to position *Geotrigona* below and *Scaptotrigona* on the top. However separations of the three honey types are not complete. An interesting fact is that HMF and ethanol accounted for by 100% of the variance in another diagram not shown here, but these two parameters did not discriminate the honey types studied here. The aliphatic organic acids were also distributed in three quadrants, in **Fig. 10B**. Fumaric acid in the upper left, the major acids (acetic, citric and lactic) to the right, and formic, malic, pyruvic




Fig. 9. Major sugars (g/100 g), aliphatic organic acids (g/100 g), amino acids (mg/kg) and plant markers (mg/kg) in honey produced by *Geotrigona* (Geo) n = 8, *Melipona* (Mel) n = 4 and *Scaptotrigona* (Sca) n = 8. Boxplots show the range of values (median thick line, lower 0.25 and upper 0.75 quartile white boxes, minimum and maximum values whiskers, and outliers of each dataset). Significant differences of honey between the three genera in each parameter were marked (Scheffé, *P < 0.05). Note that outliers in the three amino acids and trigonelline belong to different honey samples, one *Geotrigona* and three *Scaptotrigona*.




Fig. 10. PCA loading plots for NMR metabolites: Sugars (A), Aliphatic organic acids (B), Amino acids (C), and Markers (D) in pot-honeys produced by *Geotrigona* (Geo), *Melipona* (Mel) and *Scaptotrigona* (Sca) Ecuadorian stingless bees.

and succinic acids in the lower right. F1 explained 53.28% of the variance and F2 14.24%. A similar distribution for the amino acids in Fig. 10C. The aspartic acid is separated by the first dimension to the left from the remaining seven amino acids. Tyrosine, phenylalanine and leucine are in the upper right quadrant. Glutamine and a tight cluster of alanine, proline and valine are in the lower right quadrant. Looking at the amino acids, the first and second dimensions accounted for by 59.78% of the variance –almost 10% less than sugars and organic acids. It was possible to see that *Geotrigona* was separated from *Melipona* by the first dimension, whereas *Scaptotrigona* was widespread in all space (not shown). Therefore, the PCA of amino acids did not discriminate the bee genus that originated the honey. In Fig. 10D, the ten markers are distributed in all the space. Shikimic acid and methylglyoxal in the upper left, and acetoin and methylglyoxal dihydrate in the lower left. Trigonelline, pyroglutamic acid, 3-phenyllactic acid and dihydroxyacetone in the upper right and in the lower right methylglyoxal monohydrate and the most abundant 2,3-butanediol.

The consensus configuration of the three Ecuadorian pot-honeys, showed their position in biplots of sugars and organic acids in Fig. 11, with a similar resolution by sugars (11A) and organic acids (11B) to differentiate the bee genus of Ecuadorian honey. Both succeeded to cluster *Geotrigona* apart but *Melipona* and *Scaptotrigona* were not resolved. In the sugars biplot (Fig. 11A), F1 separated *Geotrigona* to the

left (gentiobiose, maltotriose and raffinose) from *Melipona* and *Scaptotrigona* to the right with the remaining seven sugars, F2 additionally positioned *Melipona* in the upper quadrant (fructose, glucose, mannose) and *Scaptotrigona* in the lower position (maltose, melezitose, sucrose, turanose). In the organic acids biplot (Fig. 11B), F1 separated *Melipona* and *Scaptotrigona* to the left with fumaric acid, and *Geotrigona* to the right with the other seven targeted NMR organic acids, but no further separation of *Melipona* and *Scaptotrigona* was achieved by F2. The first dimension explained 53.28% of the variance, and separated *Melipona* and *Scaptotrigona* honeys to the left, from *Geotrigona* to the right, where the major aliphatic organic acids (acetic, citric and lactic) were positioned in the upper right quadrant. The second component (F2 14.24%) limited the position of *Melipona* in the lower quadrant but no further discrimination was achieved from *Scaptotrigona* honey types.

Hierarchical Cluster Analysis (HCA) performed to verify the grouping of the pot-honeys according to the ten sugars contents in Fig. 11A formed two large clusters observed in the dendrogram, separating the *Geotrigona* (cluster 1) from the *Melipona* and *Scaptotrigona* (cluster 2) honeys. Similarly with the nine organic acid based dendrogram in Fig. 11B. This separation was not achieved with the two sets of eight amino acids and eight markers that were more related to the botanical origin. In the following section dendrograms will be compared with PCA.




Fig. 11. Biplots of all sugars (A), and all aliphatic organic acids (B) of pot-honeys produced by *Geotrigona* (Geo), *Melipona* (Mel) and *Scaptotrigona* (Sca) Ecuadorian stingless bees.

The relationships between the targeted metabolites and the three bee genera studied here had a descriptive and quantitative value, but did not replace missing classic quality factors such as enzyme activity, moisture, ash content and electrical conductivity, besides sensory parameters like instrumental color. Indeed, *Melipona* and *Scaptotrigona* honey clusters were separated using DA in Venezuela (Vit et al., 1998) and PCA in Brazil (Ávila et al., 2018), also the country of origin for *Geniotrigona* and *Heterotrigona* from Malaysia, and *Tetragonula* from Australia (Zawawi et al., 2022). Multiparametric analysis of these three collections of honey were not by NMR but included ash content for Venezuela and also mineral contents for the others. This fact points out the importance of inorganic elements and compounds to discriminate sets of honeys by multivariate analysis. Their origin in the nest was less studied than organic compounds. A reduction from ten to four variables was successful for DA separation of genus origin: Contents of nitrogen, reducing sugars, sucrose, and diastase activity. In contrast with the amino acids, not very discriminant in this work, total nitrogen was useful to separate *Melipona* and *Scaptotrigona* honeys from Venezuela (Vit et al., 1998). In targeted NMR, free acidity and pH were replaced by a set of organic acids, which besides the sugars were the two chemical groups able to

explain 70% of the variance. Antioxidant and antibacterial activity were also used besides the elemental mineral contents to improve the multifactorial performance to discriminate botanical, geographical and entomological origin.

3.7. Assignment of stingless bee genus to three problem honeys

Three honeys received as “abeja de tierra” the ethnic name for underground stingless bees, “bermejo” the ethnic name for *Melipona mimetica*, and “cushillomishki” a Kichwa name, were tested by PCA and HCA of all sugars biplot in our database (Fig. 11A) to explore their entomological origin.

The PCA solution in Fig. 12A positioned each problem honey in each of the regions of the three genera studied here. It is worth noting that being PCA a variable reduction approach based on correlations, inserting new honeys in the database affected the initial reference plot, as seen by comparing Fig. 6A with 20 honeys and Fig. 10A with 23, including the three honeys to be tested. Despite the allocated genus for each honey by PCA, in the HCA dendrogram with 20 reference pot-honeys + 3 problem samples (Fig. 10B), the separation was not achieved. The three




Fig. 12. Biplot (A) and dendrogram (B) of all ten sugars in 23 honeys produced by *Geotrigona* (Geo), *Melipona* (Mel) and *Scaptotrigona* (Sca) Ecuadorian stingless bees, (see Fig. 11 B biplot and 7A dendrogram for 20 pot-honeys). In the sugars biplot, suggested solutions of the three additional problem honeys are highlighted with a dot-circle: Ati is in the left side of the biplot with *Geotrigona*, Ber is not in the *Melipona* but in the *Scaptotrigona* region, positioned in the upper-right quadrant, and ★Cus could be a *Melipona* honey positioned in the lower right quadrant. However in the dendrogram 12B the three honeys were integrated in the *Melipona-Scaptotrigona* cluster. HCA by Euclidean distance using Ward’s method.

Table 6

Contents of sugars and organic acids (g/100 g honey) in three problem honeys.

Honey	Sugars										
	F	G	Ge	M	Ma	Me	Mt	R	S	T	Total
Ati	22.01	21.24	0.45	0.73	0.46	0.19	0.09	0.29	2.98	0.00	48.44
Ber	30.41	23.20	0.03	2.15	0.00	0.28	0.04	1.11	0.25	1.57	59.04
Cus	26.68	22.87	0.01	0.28	0.00	0.13	0.02	0.47	1.62	0.31	52.39

Honey	Organic acids										
	AcA	CitA	FoA	FuA	LacA	MalA	PygA	PyrA	QA	SucA	Total
Ati	0.65	0.07	0.05	0.00	0.93	0.01	0.01	0.00	0.00	0.00	1.72
Ber	0.17	0.00	0.00	0.00	0.17	0.00	0.00	0.00	0.00	0.00	0.34
Cus	0.21	0.02	0.00	0.00	0.23	0.00	0.00	0.00	0.00	0.00	0.46

Honey: **Ati** Abeja de tierra *Geotrigona*, **Ber** Bermejo *Melipona*, **Cus** Cushillomishki Kichwa name. **Sugars:** F fructose, G glucose, Ge Gentiobiose, M maltose, Ma mannose, Me melezitose, Mt maltotriose, R raffinose, S sucrose, T turanose. **Aliphatic organic acids:** AcA acetic acid, CitA citric acid, FoA formic acid, FuA fumaric acid, LacA lactic acid, MalA malic acid, PygA pyroglutamic acid, PyrA pyruvic acid, QA quinic acid, SucA succinic acid.

problem samples were located in the *Melipona-Scaptotrigona* cluster. This would mean the diagnostic for Ati is not a *Geotrigona* known as abeja de tierra. Ber and Cus are either *Melipona* or *Scaptotrigona*. The genus *Melipona* and the genus *Scaptotrigona* are too close for individual honey separations with the ten sugars or the AOAs measured in this work, that explained a 70% of the variance >60% by amino acids >50% by botanical markers (See Fig. 8). According to Fig. 12A, Ati could be a *Geotrigona*, and Ber could be a *Melipona*. But in 12B Ati is not in the *Geotrigona* cluster. The multifactorial outcome needs further analysis for a diagnostic. Therefore, individual contents of sugars and AOAs from NMR profiles of these three honeys were given in Table 6.

From Table 6, the F + G of Ati 43.25 g/100g is too high for a *Geotrigona* honey, see the range [13.08–24.50] in Table 1, and the total organic acids are too low (1.72) compared to Table 3b (4.44). Ber has 53.61g/100 g F + G, which is within the range of *Melipona* and *Scaptotrigona* [49.12–56.10], as well as 0.17 for each acetic acid and lactic acid lower than averages in Table 3a [1.2–1.6]. The maltose content (2.15) is in the *Scaptotrigona* range [0.92–2.75], higher than Geo and Mel. Except for the lower maltose (0.28) content, Cus is similar to Mel, with lower F + G (49.55), slightly higher acetic acid (0.21) and lactic

acid (0.23). Therefore, based in this descriptive analysis, Cus is more likely a *Melipona* honey, Ber could be a *Scaptotrigona* or a *Melipona* sp. with higher maltose content and if Ati is not a *Geotrigona*, sugars were possibly added to that honey, not cane sugar but another honey perhaps from *Apis mellifera*, able to increase fructose and glucose contents. The F + G content in Ati is too high for *Geotrigona*, and too low for *Melipona* or *Scaptotrigona*. The organic acids, particularly, the acetic acid and the lactic acid are too low for *Geotrigona* and too high for *Melipona* and *Scaptotrigona*. Also, if Ati is not a *Geotrigona* honey, could be from another genus not studied here.

3.8. The honey authenticity test by interphase emulsion (HATIE) revealed pot-honey biosurfactants in *Scaptotrigona vitorum*

A diagram in Fig. 13 shows two patterns for genuine pot-honey: 1. One phase for *Scaptotrigona vitorum* and three phases for *Geotrigona* and *Melipona*. The HATIE authenticity test used with genuine pot-honeys became a Honey Biosurfactant Test (HBT). *Scaptotrigona vitorum* honey had a suspected microbial origin surfactant. Thus, causing that distinctive pattern that makes it an innovative entomological origin fingerprint




Fig. 13. Diagram for genuine Honey Biosurfactant Test. *Scaptotrigona vitorum* one phase, *Geotrigona* and *Melipona* three phases. After: Vit (2022a).

for the studied species of *Scaptotrigona* genus, compared to *Geotrigona* and *Melipona*, in this set of pot-honeys.

3.9. Searching stingless bee markers in pot-honey

Bee-related lipophilic signals in the central region of honey ^{13}C -NMR spectra (3.5–4.5 ppm), are more likely to be entomological markers (Vit et al., 2015). However, considering the selectivity of food choices in different habitats, diverse chemical classes (amino acids, AOAs and sugars) were showing a functional role for entomological assignment of pot-honey. They would ideally indicate what kind of bee produced the honey –at least comparing bees from the same region. Their presence and concentrations represent different origins and functions. An unusual HPLC high maltose content was proposed as a predictor of entomological origin for Venezuelan pot-honeys using multivariate HCA, a cluster of *Melipona-Scaptotrigona* was separated from *Frieseomelitta-Nannotrigona-Tetragonisca* (Vit et al., 1998a). A further discriminant analysis (DA) with ten quality factors separated *Melipona*, *Scaptotrigona* from Trigonini into three clusters (Vit et al., 1998b). Phylogenetically, this pattern is interesting given that *Melipona* and *Scaptotrigona* are not closely related (Roubik, 1992; Rasmussen and Cameron, 2010). The maltose source is a minor floral nectar, more related to transglycosilation enzymes added by bees to make honey (Low et al., 1988). An NMR-LCMS-based metabolomics approach of stingless bee honey from Malaysia suggested discriminant metabolites were one amino acid, two organic acids and four sugars: These were D-fructofuranose in honey from *Heterotrigona itama*, α -D-glucose, β -D-glucose, D-xylose from *Geotrigona thoracica* and L-lactic acid, acetic acid, L-alanine from *Tetrigona apicalis* (Razali et al., 2018). In our data, raffinose was a sugar that discriminated *Geotrigona* (5.03 g/100 g) from a *Melipona* and *Scaptotrigona* mixed cluster. (See Fig. 7B). However, raffinose also was in the sugar spectra (0.01–0.17 g/100 g) of *A. mellifera* honey from Korea (Jang et al., 2016), (0.03–0.08 g/100 g) from Spain (Pascual-Mat ea et al., 2018), and (0.65 g/100 g) of *M. ferruginea* from Tanzania (Popova et al., 2021). Therefore, it was not an entomological marker specific for *Geotrigona*, but it was a discriminant sugar compared to *Melipona* and *Scaptotrigona* honeys from Ecuador in this study. A genetic bee-origin of chemical profiles in honey has not been approached as the vast literature on geographic and plant origin –bee diet– It is referred to as entomological origin mainly for comparative scopes of compositional (chemical, botanical, physical) and functional (antibacterial, anticancer, anti-inflammatory, antioxidant, cytotoxic, sensory) properties. This possible influence of stingless bee genus would help our understanding of environmental or other factors, if in fact honey production has a component driven by bee phylogeny or taxonomic group. Navigating tricky questions regarding stingless bee phylogeny is an expression used by Gr uter (2020) in the preface of his book, to anticipate the complexity of the topic.

Bees are known to be selective in their choices of nectar, pollen, and plant resin sources. *Geotrigona acapulconis* collected nectar from ‘Hass’ cultivars of avocado *Persea americana* in its native Mexico. Among the *Geotrigona argentina* floral resources studied in 146 pollen pots and combined pot-honey from four nests, major nectar sources overlapped major pollen sources (Vossler et al., 2010). Sympatric *Tetragonula carbonaria* and *Austroplebeia australis* use of resources may have phylogeographic origins to avoid or reduce competition: Firstly, *T. carbonaria* higher flight activity enabled faster sugar intake from a broader plant spectrum than *A. australis*, and secondly, the smaller *A. australis* colonies collected higher sugar concentrations from a narrower plant spectrum (Leonhardt et al., 2014). Accordingly, these distinctive behavioral traits could convey the chemical composition of honey. Thus, studying chemical traits of honey may reveal phylogenetic signals, and seems possible. Surprisingly, compositional profiles of honey produced by domesticated stingless bees were more similar to domesticated *Apis* spp. honey than to feral pot-honey by targeted protonic NMR (Noiset et al., 2022).

For example, after feeding sucrose solutions to *T. carbonaria*, a fake honey with (64–72) g trehalulose/100 g honey, (18–23) erlose, (9–12) fructose, and minor glucose was produced in contrast to a fructose: glucose:water (25:25:50) feeding that produced a 58 g/100 g total sugars were the F/G ratio fed remained unchanged. These authors hypothesized a major trehalulose honey from stingless bees feeding high sucrose floral nectar (Hungerford et al., 2021). A feral small stingless bee *Frieseomelitta* aff. *varia* from Venezuela, also had a surprisingly low 6.9 g glucose/100 g honey, possibly the very high (56.9) maltose-like HPLC sugar was trehalulose (Vit et al., 1998a). Being trehalulose a rare sucrose isomer α -(1 \rightarrow 1) glucose-fructose, bee-nest-enzymes involved in its anabolic/catabolic pathways could be a potential candidate for a valid long-term physiological trait, in agreement with behavioral observations by Leonhardt et al. (2014). Nectars colonized by microbial species affected bee foraging behavior based on chemical changes they caused to the nectar (Good et al., 2014), nectar volatiles influenced food preference and acceptability (Rering et al., 2018), pollen and yeast-based emissions were compared (Rering et al., 2021). A floral nectar colonizing yeast *Metschnikowia* that may reach the bee nest, was a common genus in honey bee guts (Kakumanu et al., 2016). Genes codifying for α -glucosidases from the nectar-yeast *M. gruessii* and *M. reukaufii* were expressed (Garcia-Gonzalez et al., 2019), and produced rare sugars present in honey (erlose, isomelezitose, trehalulose) after sucrose splitting activity (Garcia-Gonzalez et al., 2020).

A note on phylogenetic traits. They are diverse features essential to understanding the evolution of biodiversity, connecting microevolutionary processes of natural selection to macroevolutionary patterns of phenotypic evolution (Hansen and Martins, 1996). Not only morphological traits, but also long-term changes in physiology and behavior are traits of phylogeny –sharing features. A phylogenetic or family tree is a graphic indicator with branches (relationships) used to depict a history of evolution for a group of taxa (i.e., species, genus). The term “phylogeny” was coined in German by Ernst Haeckel in 1870 (the word is of Greek derivations where phylos φ υλον = race + geneia γενε a = birth or origin). Dendrograms of stingless bee taxa related by the chemical profiles of their honey can inform us on the skills to: 1. Select sugary materials available from the environment, 2. Apply bee-physiology for transformations in the nest, i.e. bee-enzymes 3. Select associated microbiota for further processing, defense, and preservation of a moist sugary substrate from spoilage in the warm nest. Phylogenetic and geographical signals of host-specific LAB in *Tetragonula* and *Austroplebeia* stingless bees contributed to the evolutionary history of the bee-LAB association (Leonhardt et al., 2014).

In a morphological trait such as the cuticular hydrocarbon profile, bee species may be distinctive: 1. Different drivers can affect the chemical profile based on functionality, 2. The ability to acquire environmental materials (chemical compounds) was strongly correlated with functionality, as well as the overall phylogenetic placement of the bee (Leonhardt et al., 2013). What physiological traits instead of a morphological trait would convey a similar correlation for honey? A multitask behavior to preserve the quality of honey processing for the health of bee colonies (Leonhardt, 2017). The nutritional choices made by stingless bees on food resources, may additionally become a solid source of information for long-term conservation plans (Leonhardt et al., 2020). Furthermore, a bee-derived marker, was that of associated microbiota-derived marker. More microbial-origin markers of the stingless bee nest are waiting to be discovered in the future, in addition to the overwhelming quantities of acetic acid and lactic acid they produce. A *Bacillus* marker or a related AAB-LAB, a yeast or fungal fingerprints? Arabitol and mannitol of cerumen extracts from stingless bee nests are sweet sugar alcohols (polyols) that may be of interest as fungal origin (Popova et al., 2021). Another answer now is a suspected *Starmerella* yeast associated with *Scaptotrigona vitorum*, one possibility is *Starmerella bombicola*, known for the extracellular sophorolipid biosynthesis with surfactant action. This fingerprint was not detected by ^1H -NMR but by the HATIE, honey authenticity test based on interphase

emulsion formed after shaking a honey water dilution with diethyl ether. The *S. vitorum* honey has a biosurfactant. Its chemical ID and the associated microbe synthesizing it were unknown, but a Honey Biosurfactant Test HBT was proposed (Vit, 2022a).

4. Conclusions

From the 41 metabolites studied here by targeted ¹H NMR, the amino acid isoleucine, the AOA quinic acid, and the botanical marker kynurenic acid were not detected in the Ecuadorian honeys. Therefore, the multivariate analysis was conducted for 10 sugars, 8 AOAs, 9 amino acids and 9 botanical markers. Sugars and organic acids were more discriminant of the bee genus than amino acids and markers. A test of three problem honeys revealed multifactorial analysis limitations, and the need of descriptive parameters for a preliminary fit of these honeys according to their bee-genus, with the incipient 20-honey database presented here. A solid database needs to grow for useful comparisons with similar honeys, and also to know the exceptions.

The targeted NMR chemical profiles were useful for a multi-purpose reference database to check an unknown pot-honey or suspected of adulteration, mislabeled or fake. Such a database is growing for botanical, entomological and geographical origins, and adulteration sources from various countries. With so many floral and entomological resources, the biodiversity of honeys in Ecuador is a treasure to be discovered. Our results with limited number of honeys deserve continuation, with few honey quality control data besides the ¹H-NMR. Considering that the pot-honey composition bears a relationship of taxa from three kingdoms represented by plant-bee-microbe interactions, facts not explained by the botanical and entomological origin, may have a microbiota answer in the nest. The discovery of biosurfactant activity in *Scaptotrigona vitorum* honey evidenced a microbial origin that needs identification as well as the chemical nature of the biosurfactant.

The scientific interest to investigate pot-honey composition is a slow cumulative effort to contribute to meliponine scientific research community for understanding the bee processes and for regulatory purposes, which eventually will support stingless bee keeping, agriculture, and consumers. A norm for the Ecuadorian honey produced by stingless bees is needed and the NMR data reported here is a support to establish required honey quality standards.

The term foodomics coined by Alejandro Cifuentes (2009) as “A discipline that studies the Food and Nutrition domains through the application of omics technologies” is needed for pot-honey research and discoveries by specialized teams on genomics, transcriptomics and proteomics, besides the metabolomics that has already started. The consistent influence of stingless bee genus in honey would help understanding environmental or other factors, if in fact stingless bee honey processing has a component driven by bee phylogeny and associations with microbes having multifaceted functions including nutritional traits. The biosurfactant activity of *Scaptotrigona vitorum* honey revealed by the test suggested microbial associations originating that visual behavior. Our conclusion is for this set of Ecuadorian pot-honeys. Echeverrigaray et al. (2021) found *Starmarella bombycolina* in the Brazilian honey of *Scaptotrigona bipunctata* and *S. depilis* but not in *S. tubiba*. Therefore, the microbial association with stingless bees was species specific for *Scaptotrigona*, not genus specific as in our Ecuadorian study with the *Scaptotrigona vitorum* species. HATIE and microbiota associated to other Ecuadorian *Scaptotrigona* species need to be studied.

Funding

Prometeo, Senescyt at Universidad Técnica de Machala from El Oro province, Ecuador scholarship to P. Vit. Quality Services International GmbH - QSI (Bremen, Germany) contributed with NMR analysis. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CRedit authorship contribution statement

Patricia Vit: Conceptualization, Data curation, Formal analysis, Writing – original draft, Writing – review & editing. **Jane van der Meulen:** Formal analysis. **Maria Diaz:** Data curation, Formal analysis, Writing – review & editing. **Silvia R.M. Pedro:** Writing - review and editing. **Isabelle Esperança:** Data curation, Formal analysis. **Rahimah Zakaria:** Data curation, Formal analysis. **Gudrun Beckh:** Resources. **Favian Maza:** Resources. **Gina Meccia:** Writing – review & editing. **Michael S. Engel:** Writing - review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

To stingless bee keepers from Ecuador. David W. Roubik from Smithsonian Tropical Research Institute in Ancon, Panama kindly edited the intricated final section. Martin Linkogel from Quality Services International GmbH, Bremen, Germany, for his professionalism to facilitate current institutional contacts. To Omar Malagón from Universidad Técnica Particular de Loja, for the map of Ecuador. To CDCHTA from Universidad de Los Andes, Mérida, Venezuela. To Universidad Politécnica delle Marche, Ancona, Italy.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.crfs.2022.11.005>.

References

- Aguilar, I., Herrera, E., Vit, P., 2013. Acciones para valorizar la miel de pote [on-line]. In: Vit, P., Roubik, D.W. (Eds.), *Stingless Bees Process Honey and Pollen in Cerumen Pots*. Facultad de Farmacia y Bioanálisis, Universidad de Los Andes, Mérida, Venezuela, pp. 1–13. accessed on 10 December 2021. <http://www.saber.ula.ve/handle/123456789/35292>.
- Agussalim, A., Umami, N., Nurliyani, Agus, A., 2021. The physicochemical composition of honey from Indonesian stingless bee (*Tetragonula laeviceps*). *Biodiversitas* 22, 3257–3263. <https://doi.org/10.13057/biodiv/d220820>.
- Alissandrakis, E., Tarantilis, P.A., Pappas, C., Harizanis, P.C., Polissiou, M., 2009. Ultrasound-assisted extraction gas chromatography–mass spectrometry analysis of volatile compounds in unifloral thyme honey from Greece. *Eur. Food Res. Technol.* 229, 365–373. <https://doi.org/10.1007/s00217-009-1046-8>.
- Ávila, S., Beuxa, M.R., Hoffmann Ribania, R., Zambiazib, R., 2018. Stingless bee honey: quality parameters, bioactive compounds, health promotion properties and modification detection strategies. *Trends Food Sci. Technol.* 81, 37–50. <https://doi.org/10.1016/j.tifs.2018.09.002>.
- Ávila, S., Lazzarotto, M., Hornung, P.S., Lopes Teixeira, G., Ito, V.C., Barba Bellettini, M., Regina Beux, M., Beta, T., Hoffmann Ribani, R., 2019. Influence of stingless bee genus (*Scaptotrigona* and *Melipona*) on the mineral content, physicochemical and microbiological properties of honey. *J. Food Sci. Technol.* 56, 4742–4748. <https://doi.org/10.1007/s13197-019-03939-8>.
- Barbosa, R.N., Leong, S.L., Vinnere-Pettersson, O., Chen, A.J., Souza-Motta, C.M., Frisvad, J.C., Houbraeken, J., 2017. Phylogenetic analysis of *Monascus* and new species from honey, pollen and nests of stingless bees. *Stud. Mycol.* 86, 29–51. <https://doi.org/10.1016/j.simyco.2017.04.001>.
- Baumung, R., Formato, G., Pettis, J., 2021. Introduction, 25. In: FAO, IZSLT, Apimondia, CAAS (Eds.), *Good beekeeping practices for sustainable apiculture*, FAO Animal Production and Health Guidelines No, pp. 1–4. Rome. <https://www.fao.org/3/cb5353en/cb5353en.pdf>.
- Biluca, F.C., Bernal, J., Valverde, S., Ares, A.M., Gonzaga, L.V., Costa, A.C.O., Fett, R., 2019. Determination of free amino acids in stingless bee (Meliponinae) honey. *Food Anal. Methods* 902–907. <https://doi.org/10.1007/s12161-018-01427-x>, 2019.
- Blum, M.S., Brand, J.A., 1972. *Social insect pheromones: their chemistry and function*. *Am. Zool.* 12, 553–576.
- Bogdanov, S., Vit, P., Kilchenman, V., 1996. Sugar profiles and conductivity of stingless bee honeys from Venezuela. *Apidologie* 27, 445–450.
- Bogdanov, S., Charrière, J.-D., Imdorf, A., Kilchenmann, V., Fluri, P., 2002. Determination of residues in honey after treatments with formic and oxalic acid under field conditions. *Apidologie* 33, 399–409.

- Bong, J., Loomes, K.M., Lin, B., Stephens, J.M., 2018. New approach: chemical and fluorescence profiling of NZ honeys. *Food Chem.* 267, 355–367. <https://doi.org/10.1016/j.foodchem.2017.07.065>.
- Braghini, F., Biluca, F.C., Schulz, M., Gonzaga, L.V., Costa, A.C.O., Fett, R., 2021. Stingless bee honey: a precious but unregulated product - reality and expectations. *Food Rev. Int.* <https://doi.org/10.1080/87559129.2021.1884875>.
- Brugnerotto, P., Della Betta, F., Gonzaga, L.V., Fett, R., Oliveira Costa, A.C., 2019. A capillary electrophoresis method to determine aliphatic organic acids in bracing honeydew honey and floral honey. *J. Food Compos. Anal.* 82, 103243 <https://doi.org/10.1016/j.jfca.2019.103243>.
- Camargo, J.M.F., Pedro, S.R.M., 2007. *Meliponini lepeletier 1836*. In: Moure, J.S., Urban, D., Melo, G.A.R. (Eds.), *Catalogue of Bees (Hymenoptera, Apoidea) in the Neotropical Region*. Sociedade Brasileira de Entomologia, pp. 272–578.
- Carratu, B., Ciarocchi, M., Mosca, M., Sanzini, E., 2011. Free amino acids, oxalate and sulphate for honey characterization. *J. ApiProduct ApiMed. Sci.* 3, 81–88.
- Charney, W., 1978. Dihydroxyacetone. US Patent 4, 589, 076.
- Cifuentes, A., 2009. Food analysis and foodomics. *J. Chromatogr. A* 1216, 7109. <https://doi.org/10.1016/j.chroma.2009.09.018> accessed on 18 December 2021.
- Coketin, N.N., Pappalardo, M., Campbell, L.T., Brooks, P., Carter, D.A., Blair, S., Harry, E.J., 2016. The antibacterial activity of Australian *Leptospermum* honey correlates with methylglyoxal levels. *PLoS One* 11, e0167780. <https://doi.org/10.1371/journal.pone.0167780>.
- Codex, S.T.A.N., 1987, 1987. In: Codex Standard for Honey (World-wide Standard) 12-1981 Rev. vol. 1. accessed on 18 December 2021. <https://www.fao.org/3/w0076e/w0076e30.htm>.
- Consonni, R., Cagliani, L.R., Cogliati, C., 2012. NMR characterization of saccharides in Italian honeys of different floral sources. *J. Agric. Food Chem.* 60, 4526–4534. <https://doi.org/10.1021/jf3008713>.
- D'Arcy, B.R., Rintoul, G.B., Rowland, C.Y., Blackman, A.J., 1997. Composition of Australian honey extractives. 1. Norisoprenoids, monoterpenes, and other natural volatiles from blue gum (*Eucalyptus leucocylon*) and yellow box (*Eucalyptus melliodora*) honeys. *J. Agric. Food Chem.* 45, 1834–1843. <https://doi.org/10.1021/jf960625>.
- Department of Standards Malaysia, 2017. Kelulut (Stingless Bee) Honey - Specification MS 2683, 2017. <https://es.scribd.com/document/398215369/Kelulut-Stingless-bee-honey-Specification>.
- Echeverrigaray, S., Scariot, F.J., Foresti, L., Schwarz, L.V., Ronaldo Kauê Mattos Rocha, R.K.M., da Silva, G.P., Moreira, J.P., Delamare, A.P.L., 2021. Yeast biodiversity in honey produced by stingless bees raised in the highlands of southern Brazil. *Int. J. Food Microbiol.* 347, 109200 <https://doi.org/10.1016/j.jfmmicro.2021.109200>.
- Engel, M.S., 2021. A key to the subgenera of the stingless bee genus *Melipona* (Hymenoptera: Apidae). *Entomol. Mon. Mag.* 157, 273–281.
- Engel, M.S., 2022. Notes on South American stingless bees of the genus *Scaptotrigona* (Hymenoptera: Apidae). Part IV: four new species of group B from the Andean region. *J. Melittology* 112, 1–13. <https://doi.org/10.17161/jom.v112.18128>.
- Fletcher, M., Hungerford, N.L., Webber, D., de Jesus, M.C., Zhang, J., Stone, I.S., Zawawi, N., 2020. Stingless bee honey, a novel source of trehalose: a biologically active disaccharide with health benefits. *Sci. Rep.* 10, 12128 <https://doi.org/10.1038/s41598-020-68940-0>.
- de la Fuente, E., Sanz, M.L., Martínez-Castro, I., Sanz, J., Ruiz-Matute, A.I., 2007. Volatile and carbohydrate composition of rare unifloral honeys from Spain, 2007 *Food Chem.* 105, 84–89. <https://doi.org/10.1016/j.foodchem.2007.07.023>.
- Duarte, A.W.F., Vasconcelos, M.R.S., de Menezes, A.P.D., da Silva, S.C., Oda-Souza, M., López, A.M.Q., 2012. Composition and antioxidant activity of honey from Africanized and stingless bees in Alagoas (Brazil): a multivariate analysis. *J. Apic. Res.* 51, 23–35. <https://doi.org/10.3896/JBRA.1.51.1.04>.
- Fuentes Molina, O., Alizadeh, K., Sergio, A., Bucareyd, S.A., Castaneza Zúñiga, E., Vásquez Quiral, P., 2020. Analysis of organic molecules, physicochemical parameters, and pollen as indicators for authenticity, botanical origin, type and quality of honey samples examined. *Int. J. Food Prop.* 23, 2242–2256. <https://doi.org/10.1080/10942912.2020.1850775>.
- García-González, M., Minguet-Lobato, M., Plou, F.J., Fernández-Lobato, M., 2020. Molecular characterization and heterologous expression of two α -glucosidases from *Metschnikowia* spp, both producers of honey sugars. *Microb. Cell Factories* 140.
- García-González, M., Plou, F.J., Cervantes, F.V., Remacha, M., Poveda, A., Jiménez-Barbero, J., Fernández-Lobato, M., 2019. Efficient production of isomelezitose by a glucosyltransferase activity in *Metschnikowia reukauffii* cell extracts. *Microb. Biotechnol.* 12, 1274–1285. <https://doi.org/10.1111/1751-7915.13490>.
- Gerginova, D., Simova, S., Popova, M., Stefova, M., Stanoeva, J.P., Bankova, V., 2020. NMR Profiling of North Macedonian and Bulgarian honeys for detection of botanical and geographical origin. *Molecules* 25, 4687. <https://doi.org/10.3390/molecules25204687>.
- Gilliam, M., 1997. Identification and roles of non-pathogenic micro-flora associated with honey bees. *FEMS Microbiol. Lett.* 155, 1–10. <https://doi.org/10.1111/j.1574-6968.1997.tb12678.x>.
- Gilliam, M., Roubik, D.W., Lorenz, J., 1990. Microorganisms associated with pollen, honey, and brood provisions in the nest of a stingless bee, *Melipona fasciata*. *Apidologie* 21, 89–97.
- Gonnet, M., Lavie, P., Nogueira-Neto, P., 1964. Étude de quelques caractéristiques des miels récoltés para certains *Meliponines* brésiliens. *C. R. Acad. Sci Paris* 258, 3107–3109.
- Good, A.P., Gauthier, M.-P.L., Vannette, R.L., Fukami, T., 2014. Honey bees avoid nectar colonized by three bacterial species, but not by a yeast species, isolated from the gut. *PLoS One* 9, e86494. <https://doi.org/10.1371/journal.pone.0086494>.
- Gresley, A.L., Kenny, J., Cassar, C., Kelly, A., Sinclair, A., Fielder, M.D., 2012. The application of high resolution diffusion NMR to the analysis of manuka honey. *Food Chem.* 135, 2879–2886. <https://doi.org/10.1016/j.foodchem.2012.07.072>.
- Grüter, C., 2020. *Stingless Bees. Their Behavior, Ecology and Evolution*. Fascinating Life Sciences. Springer Nature, Cham, Switzerland.
- Guyot, C., Scheirman, V., Collin, S., 1999. Floral origin markers of heather honeys: *Calluna vulgaris* and *Erica arborea*. *Food Chem.* 64, 3–11.
- Hansen, T.F., Martins, E.P., 1996. Translating between microevolutionary process and macroevolutionary patterns: the correlation structure of interspecific data. *Evolution* 50, 1404–1417. <https://doi.org/10.1111/j.1558-5646.1996.tb03914.x>.
- Heard, T.A., 2016. *The Australian Native Bee Book. Keeping Stingless Bee Hives for Pets, Pollination and Sugarbag Honey*. Sugarbag Bees, Brisbane.
- Hermosin, I., Chicon, R.M., Cabezudo, M.D., 2003. Free amino acid composition and botanical origin of honey. *Food Chem.* 83, 263–268.
- Hungerford, N.L., Zhang, Smith, T.J., Yates, H.S.A., Chowdhury, S.A., Carter, J.M., Carpinelli de Jesus, M., Fletcher, M.T., 2021. Feeding sugars to stingless bees: identifying the origin of trehalose-rich honey composition. *J. Agric. Food Chem.* 69, 10292–10300. <https://doi.org/10.1021/acs.jafc.1c02859>.
- IBM Corp., 2019. IBM SPSS Statistics for Windows, Version 26.0. IBM Corp. Armonk, NY.
- Icon, N.S., Manley-Harris, M., Clearwater, M.J., 2021. Floral nectar of wild manuka (*Leptospermum scoparium*) varies more among plants than among sites. *N. Z. J. Crop Hortic. Sci.* 47 <https://doi.org/10.1080/01140671.2019.1670681>.
- Jang, E.-S., Kim, I.-S., Lee, E.-J., Seo, H.-S., Lee, H.-J., Kim, E., Kim, K.-T., Kim, J.-B., 2016. Characterization of traditional Korean unifloral honey based on the mono-, di-, and trisaccharides. *Korean J. Food Sci. Technol.* 48, 1–8. <https://doi.org/10.9721/KJFST.2016.48.1.1>.
- Kakumani, M.L., Reeves, A.M., Anderson, T.D., Rodrigues, R.R., Williams, M.A., 2016. Honey bee gut microbiome is altered by in-hive pesticide exposures. *Front. Microbiol.* 7, 1255.
- Khalloubi, F., Akdad, M., Bouddine, T., Hajji, L., Owen, R.W., 2021. HPLC-ESI-MS and GC-EM Identification and quantitation of polyphenolics and alkaloids in Moroccan jujube honeys. *J. Apicult. Sci.* 64, 287–299.
- Khansariroeh, E., Salmaki, Y., Akbari Azirani, T., Henareh, F., Alizadeh, K., Ramezani, E., Zarre, S., Beckh, G., Behling, H., 2021. The sources and quality of Iranian honey. *Heliyon* 7, e06651. <https://doi.org/10.1016/j.heliyon.2021.e06651>.
- Kuš, P.M., Jerkovic, I., Tuberoso, C.I.G., Šarolic, M., 2013. The volatile profiles of a rare apple (*Malus domestica* Borkh.) honey: shikimic acid-pathway derivatives, terpenes, and others. *Chem. Biodivers.* 10, 1638–1652. <https://doi.org/10.1002/cbdv.201200404>.
- Kwong, W.K., Medina, L.A., Koch, H., Sing, K.-W., Soh, E.J.Y., Ascher, J.S., Jaffe, R., Moran, N.A., 2017. Dynamic microbiome evolution in social bees. *Sci. Adv.* 3, e1600513 <https://doi.org/10.1126/sciadv.1600513>.
- Leonhardt, S.D., 2017. Chemical ecology of stingless bees. *J. Chem. Ecol.* 43, 385–402. <https://doi.org/10.1007/s10886-017-0837-9>.
- Leonhardt, S.D., Lihoreau, M., Spaethe, J., 2020. Mechanisms of nutritional resource exploitation by insects. *Insects* 11, 570. <https://doi.org/10.3390/insects11090570>.
- Leonhardt, S.D., Heard, T.A., Wallace, H.M., 2014. Differences in the resource intake of two sympatric Australian stingless bee species. *Apidologie* 45, 514–527. <https://doi.org/10.1007/s13592-013-0266-x>.
- Leonhardt, S.D., Kaltenpoth, M., 2014. Microbial communities of three sympatric Australian stingless bee species. *PLoS One* 9, e105718. <https://doi.org/10.1371/journal.pone.0105718>.
- Leonhardt, S.D., Rasmussen, C., Schmitt, T., 2013. Genes versus environment: geography and phylogenetic relationships shape the chemical profiles of stingless bees on a global scale. *Proc. R. Soc. B* 280, 20130680 <https://doi.org/10.1098/rspb.2013.0680>.
- Leoni, V., Giupponi, L., Pavlovic, R., Gianoncelli, C., Cecati, F., Ranzato, E., Martinotti, S., Pedrali, D., Giorgi, A., Panseri, S., 2021. Multidisciplinary analysis of Italian Alpine wildflower honey reveals criticalities, diversity and value. *Sci. Rep.* 11, 19316 <https://doi.org/10.1038/s41598-021-98876-y>.
- Lolli, M., Bertelli, D., Plessi, M., Sabatini, A.G., Restani, C., 2008. Classification of Italian honeys by 2D HR-NMR. *J. Agric. Food Chem.* 56, 1298–1304. <https://doi.org/10.1021/jf072763c>.
- Low, N.H., Nelson, D.L., Sporns, P., 1988. Carbohydrate analysis of Western Canadian honeys and their nectar sources to determine the origin of nectar oligosaccharides. *J. Apic. Res.* 27, 245–251.
- Machado, A.M., Miguel, M.G., Vilas-Boas, M., Figueiredo, A.C., 2020. Honey volatiles as a fingerprint for botanical origin - A review on their occurrence on monofloral honeys. *Molecules* 25, 374. <https://doi.org/10.3390/molecules25020374>.
- Massaro, C.F., Shelley, D., Heard, T.A., Brooks, P., 2014. In vitro antibacterial phenolic extracts from "sugarbag" pot-honeys of Australian stingless bees (*Tetragonula carbonaria*). *J. Agric. Food Chem.* 62, 12209–12217. <https://doi.org/10.1021/jf5051848>.
- Mato, I., Huidobro, J.F., Simal-Lozano, J., Sancho, M.T., 2006. Capillary zone electrophoresis method for the determination of inorganic anions and formic acid in honey. *J. Agric. Food Chem.* 54, 9292–9296. <https://doi.org/10.1021/jf061536s>.
- Meister, A., Gutierrez-Gines, M.J., Maxfield, A., Gaw, S., Dickinson, N., Horswell, J., Robinson, B., 2021. Chemical elements and the quality of manuka (*Leptospermum scoparium*) honey. *Foods* 10, 1670. <https://doi.org/10.3390/foods10071670>.
- Menezes, C., Vollet-Neto, A., Marsaioli, A.J., Zamplieri, D., Fontoura, I.C., Luchessi, A.D., Imperatriz-Fonseca, V.L., 2015. A Brazilian social bee must cultivate fungus to survive. *Curr. Biol.* 25, 2851–2855. <https://doi.org/10.1016/j.cub.2015.09.028>.
- Michener, C.D., 2000. *The Bees of the World*. Johns Hopkins University Press, Baltimore.
- Michener, C.D., 2013. *The Meliponini*. In: Vit, P., Pedro, S.R.M., Roubik, D. (Eds.), *Pot-honey. A Legacy of Stingless Bees*. Springer, New York, pp. 3–17.

- Moo-Huchin, V.M., Gonzalez-Aguilar, G.A., Lira-Maas, J.D., Perez-Pacheco, E., Estrada-Leon, R., MooHuchin, M.I., Sauri-Duch, E., Sauri-Duch, E., 2015. Physicochemical properties of *Melipona beecheii* honey of the Yucatan Peninsula. *J. Food Res.* 4, 25. <https://doi.org/10.5539/jfr.v4n5p25>.
- Morais, P.B., Sao Thiago Calaca, P.S., Rosa, C.A., 2013. Microorganisms associated with stingless bees. In: Vit, P., Pedro, S.R.M., Roubik, D. (Eds.), *Pot-honey. A Legacy of Stingless Bees*. Springer, New York, pp. 173–186.
- Ng, W.-J., Sit, N.-W., Ooi, P.A.-C., Ee, K.-Y., Lim, T.-M., 2021. Botanical origin differentiation of Malaysian stingless bee honey produced by *Heterotrigona itama* and *Geniotrigona thoracica* using chemometrics. *Molecules* 26, 7628. <https://doi.org/10.3390/molecules26247628>.
- Ngalimat, M.S., Raja, A., Rahman, R.N.Z., Yusof, M.T., Syahir, A., Sabri, S., 2019. Characterisation of bacteria isolated from the stingless bee *Heterotrigona itama* honey, bee bread and propolis. *PeerJ* 7, e7478. <https://doi.org/10.7717/peerj.7478>.
- Noiset, P., Cabriol, N., Rojas-Oropeza, M., Warrit, N., Nkoba, K., Vereecken, N.J., 2022. Honey compositional convergence and the parallel domestication of social bees. *Sci. Rep.* 12, 18280. <https://doi.org/10.1038/s41598-022-23310-w>.
- Ohmenhaeuser, M., Monakhova, Y.B., Kuballa, T., Lachenmeier, D.W., 2013. Qualitative and quantitative control of honeys using NMR spectroscopy and chemometrics. *Anal. Chem.*, 825318 <https://doi.org/10.1155/2013/825318>, 2013.
- Olofsson, T.C., Butler, E., Markowicz, P., Lindholm, C., Larsson, L., Vásquez, A., 2016. Lactic acid bacterial symbionts in honeybees – an unknown key to honey's antimicrobial and therapeutic activities. *Int. Wound J.* 13, 668–679. <https://doi.org/10.1111/iwj.12345>, 2016.
- Pascual-Mateá, A., Osésa, S.M., Marazzan, G.L., Gardini, S., Fernández Muñio, M.A., Sancho, M.T., 2018. Sugar composition and sugar-related parameters of honeys from the northern Iberian Plateau. *J. Food Compos. Anal.* 74, 34–43.
- de Paula, G.T., Menezes, C.C., Pupo, M.T., Rosa, C.A., 2021. Stingless bees and microbial interactions. *Curr. Opin. Insect Sci.* 44, 41–47. <https://doi.org/10.1016/j.cois.2020.11.006>.
- Pérez, R.A., Sánchez-Brunete, C., Calvo, R.M., Tadeo, J.L., 2002. Analysis of volatiles from Spanish honeys by solid-phase microextraction and gas chromatography–mass spectrometry. *J. Agric. Food Chem.* 50, 2633–2637. <https://doi.org/10.1021/jf011551r>.
- Pérez-Pérez, E.M., Suárez, E., Peña-Vera, M.J., González, A.C., Vit, P., 2013. Antioxidant activity and microorganisms in nest products of *Tetragonisca angustula* Latreille, 1811 from Mérida, Venezuela [on-line]. In: Vit, P., Roubik, D.W. (Eds.), *Stingless Bees Process Honey and Pollen in Cerumen Pots*. Facultad de Farmacia y Bioanálisis, Universidad de Los Andes, Mérida, Venezuela, pp. 1–8. accessed on 10 December 2021. <http://www.saber.ula.ve/handle/123456789/35292>.
- Persano-Oddo, L.P., Heard, T.A., Rodríguez-Malaver, A., Pérez, R.A., Fernández-Muñio, M., Sancho, M.T., Sesta, G., Lusco, L., Vit, P., 2008. Composition and antioxidant activity of *Trigona carbonaria* honey from Australia. *J. Med. Food* 11, 789–794. <https://doi.org/10.1089/jmf.2007.0724>.
- Persano-Oddo, L., Piro, R., 2004. Main European unifloral honeys: descriptive sheets. *Apidologie* 35 (1), S81–S38.
- Pimentel, T.C., Rosset, M., de Sousa, J.M.B., Gomes de Oliveira, L.I., Meireles Mafaldo, I., Estevez Pintado, M.M., de Souza, E.L., Magnani, M., 2021. Stingless bee honey: an overview of health benefits and main market challenges. *J. Food Biochem.* e13883 (in press). <https://onlinelibrary.wiley.com/doi/epdf/10.1111/jfbc.13883>.
- Popova, M., Gerginova, D., Trusheva, B., Simova, S., Tamfu, A.N., Ceylan, O., Clark, K., Bankova, V.A., 2021. Preliminary study of chemical profiles of honey, cerumen, and propolis of the African stingless bee *Meliponula ferruginea*. *Foods* 10 (997), 1–17. <https://doi.org/10.3390/foods10050997>.
- Radovic, B.S., Careri, M., Mangia, A., Musci, M., Gerboles, M., Anklam, E., 2000. Contribution of dynamic headspace GC–MS analysis of aroma compounds to authenticity testing of honey. *Food Chem.* 72, 511–520, 2001.
- Rasmussen, C., Cameron, S.A., 2010. Global stingless bee phylogeny supports ancient divergence, vicariance, and long distance dispersal. *Biol. J. Linn. Soc.* 99, 206–232. <https://doi.org/10.1111/j.1095-8312.2009.01341.x>.
- Razali, M.T.A., Zainal, Z.A., Maulidiani, M., Shaari, K., Zamri, Z., Idrus, M.Z.M., Khatib, A., Abas, F., Ling, Y.S., Rui, L.L., Ismail, I.S., 2018. Classification of raw stingless bee honeys by bee species origins using the NMR and LC-MS-based metabolomics approach. *Molecules* 23. <https://doi.org/10.3390/molecules23092160>, 2018.
- Rering, C.C., Beck, J.J., Hall, G.W., McCartney, M.M., Vannette, R.L., 2018. Nectar-inhabiting microorganisms influence nectar volatile composition and attractiveness to a generalist pollinator. *New Phytol.* 220, 750–759. <https://doi.org/10.1111/nph.14809>.
- Rering, C.C., Rudolph, A.B., Beck, J.J., 2021. Pollen and yeast change nectar aroma and nutritional content alone and together, but honey bee foraging reflects only the avoidance of yeast. *Environ. Microbiol.* 23, 4141–4150. <https://doi.org/10.1111/1462-2920.15528>.
- Roubik, D.W., 1989. *Ecology and Natural History of Tropical Bees*. Cambridge University Press, New York, p. 514.
- Roubik, D.W., 1992. Stingless bees: a guide to Panamanian and Mesoamerican species and their nests (Hymenoptera: Apidae: Meliponinae). In: Quintero, D., Aiello, A. (Eds.), *Insects of Panama and Mesoamerica: Selected Studies*. Oxford University Press, Oxford, pp. 495–524.
- Roubik, D.W., 2018. 100 Species of meliponines (Apidae: Meliponini) in a parcel of Werster Amazonian Forest at Yasuní Biosphere Reserve, Ecuador. In: Vit, P., Pedro, S.R.M., Roubik, D.W. (Eds.), *Pot-pollen in Stingless Bee Melittology*. Springer, Cham, Switzerland, pp. 189–206.
- Roubik, D.W., Vergara, C., 2021. Geographical distribution of bees: a history and an update. In: FAO, IZSLT, Apimondia, CAAS (Eds.), *Good Beekeeping Practices for Sustainable Apiculture*, FAO Animal Production and Health Guidelines No. 25, pp. 11–13. Rome. <https://www.fao.org/3/cb5353en/cb5353en.pdf>.
- Salvador, L., Guijarro, M., Rubio, D., Aucatoma, B., Guillén, T., Vargas Jentzsch, P., Ciobota, V., Stolkler, L., Ulic, S., Vásquez, L., Garrido, P., Bravo, J., Ramos Guerrero, L., 2019. Exploratory monitoring of the quality and authenticity of commercial honey in Ecuador. *Foods* 8, 105. <https://doi.org/10.3390/foods8030105>. Available online.
- Sancho, M.T., Mato, I., Huidobro, J.F., Fernández-Muñio, M.A., Pascual-Maté, M.A., 2013. Nonaromatic organic acids of honeys. pp 447–458. In: Vit, P., Pedro, S.R.M., Roubik, D. (Eds.), *Pot-honey. A Legacy of Stingless Bees*. Springer, New York. https://doi.org/10.1007/978-1-4614-4960-7_32.
- Santos-Sánchez, N.F., Salas-Coronado, R., Hernández-Carlos, B., Villanueva-Cañongo, C., 2019. Shikimic acid pathway in biosynthesis of phenolic compounds. In: Soto-Hernández, M., García-Mateos, R., Palma-Tenango (Eds.), *Plant Physiological Aspects of Phenolic Compounds*. M. IntechOpen. <https://doi.org/10.5772/intechopen.83815>.
- Schievano, E., Finotello, C., Mammi, S., Illy Belci, A., Colombari, S., Navarini, L., 2015. Preliminary characterization of monofloral Coffea spp. honey: correlation between potential biomarkers and pollen content. *J. Agric. Food Chem.* 63, 5858–5863. <https://doi.org/10.1021/jf506359u>.
- Schwarz, H.F., 1948. Stingless bees (Meliponidae) of the Western Hemisphere. *Bull. Am. Mus. Nat. Hist.* 90, 1–546. <http://hdl.handle.net/2246/1231>.
- Secretaría de Regulación y Gestión Sanitaria y Secretaría de Alimentos y Bioeconomía, 2019. Miel de *Tetragonisca fiebrigi* (yatef). Resolución Conjunta 17/2019 RESFC-2019-17-APN-SRYGS#MSYDS 02/05/2019 N° 29258/19 v. 02/05/2019. <https://www.boletinoficial.gob.ar/detalleAviso/primera/206764/20190502>.
- Seraglio, S.K.T., Bergamo, G., Brugnerotto, P., Gonzaga, L.V., Fett, R., Costa, A.C.O., 2021. Aliphatic organic acids as promising authenticity markers of bractacacia honeydew honey. *Food Chem.* 343, 128449 <https://doi.org/10.1016/j.foodchem.2020.128449>.
- Shamsudin, S., Selamat, J., Sanny, M., Abd, S.-B., Jambari, N.N., Mian, Z., Khatib, A.A., 2019. Influence of origins and bee species on physicochemical, antioxidant properties and botanical discrimination of stingless bee honey. *Int. J. Food Prop.* 22, 238–263. <https://doi.org/10.1080/10942912.2019.1576730>.
- Siegmund, B., Urdl, K., Jurek, A., Leitner, E., 2018. More than honey: investigation on volatiles from monovarietal honeys using new analytical and sensory approaches. *J. Agric. Food Chem.* 66, 2432–2442. <https://doi.org/10.1021/acs.jafc.6b05009>.
- Smallfield, B.M., Joyce, N.I., van Klink, J.W., 2018. Developmental and compositional changes in *Leptospermum scoparium* nectar and their relevance to mānuka honey bioactives and markers. *N. Z. J. Bot.* 56, 183–197. <https://doi.org/10.1080/0028825X.2018.1446450>.
- Smith, F., 1863. Descriptions of Brazilian honey bees belonging to the genera *Melipona* and *Trigona*, which were exhibited, together with samples of their honey and wax, in the Brazilian Court of the International Exhibition of 1862. *Trans. R. Entomol. Soc. Lond.* 11, 497–512. <https://doi.org/10.1111/j.1365-2311.1863.tb01298.x>.
- de Sousa, L.P., 2021. Bacterial communities of indoor surface of stingless bee nests. *PLoS One* 16, e0252933. <https://doi.org/10.1371/journal.pone.0252933>.
- de Sousa, J.M.B., Souza, E.L., Marques, G., Benassi, M.T., Gullón, B., Pintado, M.M., Magnani, M., 2016. Sugar profile, physicochemical and sensory aspects of monofloral honeys produced by different stingless bee species in Brazilian semi-arid eegion. *LWT–Food Sci. Technol.* 65, 645–651. <https://doi.org/10.1016/j.lwt.2015.08.058>.
- Spraul, M., Schütz, B., Rinke, P., Koswig, S., Humpfer, E., Schäfer, H., Mortter, M., Fang, F., Marx, U.C., Minoja, A., 2009. NMR-based multi parametric quality control of fruit juices: SGF profiling. *Nutrients* 1, 148–155. <https://doi.org/10.3390/nu1020148>.
- Suarez-Luque, S., Mato, I., Huidobro, J.F., Simal-Lozano, J., Sancho, M.T., 2006. Capillary zone electrophoresis method for the determination of inorganic anions and formic acid in honey. *J. Agric. Food Chem.* 54, 9292–9296. <https://doi.org/10.1021/jf061536s>.
- Suto, M., Kawashima, H., Nakamura, Y., 2020. Determination of organic acids in honey by liquid chromatography with tandem mass spectrometry. *Food Anal. Methods* 13, 2249–2257. <https://doi.org/10.1007/s12161-020-01845-w>.
- Torres, A., Garedew, A., Schmolz, E., Lamprecht, I., 2004. Calorimetric investigation of the antimicrobial action and insight into the chemical properties of “angelita” honey - a product of the stingless bee *Tetragonisca angustula* from Colombia. *Thermochim. Acta* 415, 107–113. <https://doi.org/10.1016/j.tca.2003.06.005>.
- Truchado, P., Martos, I., Bortolotti, L., Sabatini, A.G., Ferreres, F., Tomas-Barberan, F.A., 2009. Use of quinoline alkaloids as markers of the floral origin of chestnut honey. *J. Agric. Food Chem.* 57, 5680–5686. <https://doi.org/10.1021/jf900766v>.
- Villacrés-Granda, I., Coello, D., Proaño, A., Ballesteros, I., Roubik, D.W., Jijón, G., Granda-Albujá, G., Granda-Albujá, S., Abreu-Naranjo, R., Maza, F., Tejerah, E., González-Paramás, A.M., Bullón, P., Alvarez-Suarez, J.M., 2021. Honey quality parameters, chemical composition and antimicrobial activity in twelve Ecuadorian stingless bees (Apidae: Apinae: Meliponini) tested against multiresistant human pathogens. *LWT–Food Sci. Technol.* 140, 110737 <https://doi.org/10.1016/j.lwt.2020.110737>.
- Vit, P., 1995. Tubo pistón volumétrico para análisis (diseño industrial). Resolución 1286. Boletín Servicio Autónomo de la Propiedad Industrial. Caracas, 05 de Mayo de 1995. 184° y 136°. Venezuela. Cilindro adaptado para medir, agitar miel con reactivos, y observar separación de fases fundamentales para veredicto miel genuina o miel falsa.
- Vit, P., 1998. A test to detect cane sugar honey. *Arch. Latinoam. Nutr.* 48, 62–64. <https://www.alanrevista.org/ediciones/1998/1/art-12/>.
- Vit, P., 1999. Dispositivo per l'analisi del miele. AR93U002, Roma, 1999.
- Vit, P., 2022a. A Honey Authenticity Test by Interphase Emulsion Reveals Biosurfactant Activity and Biotechnology in the Stingless Bee Nest of *Scaptotrigona vitorum*

- 'Catiana' from Ecuador. *Interciencia* 47, 416–425. https://www.interciencia.net/wp-content/uploads/2022/10/01_6935_A_Vit_v47n10_10.pdf.
- Vit, P., 2022b. Sour honeys from 57 species of stingless bees in 18 countries. *Bee World* 99, 1–8. <https://doi.org/10.1080/0005772X.2022.2079842>.
- Vit, P., Chuttong, B., Zawawi, N., Diaz, M., van der Meulen, J., Ahmad, H.F., Tomas-Barberan, F.A., Meccia, G., Danmek, K., Moreno, J.E., Roubik, D.W., Barth, O.M., Lachenmeier, D.W., Engel, M.S., 2022. A novel integrative methodology for research on pot-honey variations during post-harvest. *Sociobiology* 69, e8251. <https://doi.org/10.13102/sociobiology.v69i4.8251>.
- Vit, P., Fernández-Maeso, M.C., Ortiz-Valbuena, A., 1998a. Potential use of the three frequently occurring sugars in honey to predict stingless bee entomological origin. *J. Appl. Entomol.* 122, 5–8. <https://doi.org/10.1111/j.1439-0418.1998.tb01452.x>.
- Vit, P., Medina, M., Enriquez, M.E., 2004. Quality standards for medicinal uses of Meliponinae honey in Guatemala, Mexico and Venezuela. *Bee World* 85, 2–5. <https://doi.org/10.1080/0005772X.2004.11099603>.
- Vit, P., Pedro, S.R.M., Meléndez, V., Frisone, V., 2018. Diversity of stingless bees in Ecuador, meliponiculture and fostering a living museum for Meliponini of the world. In: Vit, P., Pedro, S.R.M., Roubik, D.W. (Eds.), *Pot-pollen in Stingless Bee Melittology*. Springer, Cham, Switzerland, pp. 207–227.
- Vit, P., Pedro, S.R.M., Roubik, D. (Eds.), 2013. *Pot-honey. A Legacy of Stingless Bees*. Springer, New York. https://doi.org/10.1007/978-1-4614-4960-7_11.
- Vit, P., Persano Oddo, L., Marano, M.L., Salas de Mejías, E., 1998b. Venezuelan stingless bee honeys characterised by multivariate analysis of compositional factors. *Apidologie* 29, 377–389.
- Vit, P., Rojas, L.B., Usubillaga, A., Aparicio, R., Meccia, G., Fernando Muiño, M.A., Sancho, M.T., 2011. Presencia de ácido láctico y otros compuestos semivolátiles en mieles de Meliponini. *Rev. Inst. Nac. Hig. Rafael Rangel* 42, 58–63. <http://www.scielo.org.ve/pdf/inhrr/v42n1/art08.pdf>.
- Vit, P., Uddin, J., Zuccato, V., Maza, F., Schievano, E., 2015. Entomological origin of honey discriminated by NMR chloroform extracts in Ecuadorian honey. *Int. J. Nutr. Food Eng.* 9, 494–497 accessed on 10 December 2021. <https://publications.waset.org/10001263/entomological-origin-of-honey-discriminated-by-nmr-chloroform-extracts-in-ecuadorian-honey>.
- Vossler, F.G., Telleria, M.C., Cunningham, M., 2010. Floral resources foraged by *Geotrigona argentina* (Apidae, Meliponini) in the Argentine dry chaco forest. *Grana* 49, 142–153. <https://doi.org/10.1080/00173131003694274>.
- Walker, M.J., Cowen, S., Gray, K., Hancock, P., Burns, D.T., 2022. Honey authenticity: the opacity of analytical reports - part 1 defining the problem. *NPJ Sci Food* 6, 11. <https://doi.org/10.1038/s41538-022-00126-6>.
- Weaver, N., Weaver, E.C., 1981. Beekeeping with the stingless bee *Melipona beecheii* by the yucatecan Maya. *Bee World* 62, 7–19. <https://doi.org/10.1080/0005772X.1981.11097806>.
- XLSTAT Software Premium, 2021. *XLSTAT Statistical and Data Analysis Solution*. Addinsoft, New York, USA.
- Yong, C.-H., Muhammad, S.A., Aziz, F.A., Nasir, F.I., Mustafa, M.Z., Ibrahim, B., Kelly, S. D., Cannavan, A., Seow, E.-K., 2022. Detecting adulteration of stingless bee honey using untargeted ¹H NMR metabolomics with chemometrics. *Food Chem.* 368, 130808 <https://doi.org/10.1016/j.foodchem.2021.130808>.
- Zawawi, N., Zhang, J., Hungerford, N.L., Yates, H.S.A., Webber, D.C., Farrell, M., Tinggi, U., Bhandari, B., Fletcher, M.T., 2022. Unique physicochemical properties and rare reducing sugar trehalulose mandate new international regulation for stingless bee honey. *Food Chem.* 373, 131566 <https://doi.org/10.1016/j.foodchem.2021.131566>.